

# QUANTUM REPRESENTATIONS AND MONODROMIES OF FIBERED LINKS

RENAUD DETCHERRY AND EFSTRATIA KALFAGIANNI

ABSTRACT. Andersen, Masbaum and Ueno conjectured that certain quantum representations of surface mapping class groups should send pseudo-Anosov mapping classes to elements of infinite order (for large enough level  $r$ ). In this paper, we relate the AMU conjecture to a question about the growth of the Turaev-Viro invariants  $TV_r$  of hyperbolic 3-manifolds. We show that if the  $r$ -growth of  $|TV_r(M)|$  for a hyperbolic 3-manifold  $M$  that fibers over the circle is exponential, then the monodromy of the fibration of  $M$  satisfies the AMU conjecture. Building on earlier work [9] we give broad constructions of (oriented) hyperbolic fibered links, of arbitrarily high genus, whose  $SO(3)$ -Turaev-Viro invariants have exponential  $r$ -growth. As a result, for any  $g > n \geq 2$ , we obtain infinite families of non-conjugate pseudo-Anosov mapping classes, acting on surfaces of genus  $g$  and  $n$  boundary components, that satisfy the AMU conjecture.

## 1. INTRODUCTION

Given a compact oriented surface  $\Sigma$ , possibly with boundary, the mapping class group  $\text{Mod}(\Sigma)$  is the group of isotopy classes of orientation preserving homeomorphisms of  $\Sigma$  that fix the boundary. The Witten-Reshetikhin-Turaev Topological Quantum Field Theories [18, 25] provide infinite families of finite dimensional projective representations of mapping class groups. For each semi-simple Lie algebra, there is an associated theory and an infinite family of such representations. In this article we are concerned with the  $SO(3)$ -theory and we will follow the skein-theoretic framework given by Blanchet, Habegger, Masbaum and Vogel [7]: For each odd integer  $r \geq 3$ , let  $U_r = \{0, 2, 4, \dots, r-3\}$  be the set of even integers smaller than  $r-2$ . Given a primitive  $2r$ -th root of unity  $\zeta_r$ , a compact oriented surface  $\Sigma$ , and a coloring  $c$  of the components of  $\partial\Sigma$  by elements of  $U_r$ , a finite dimensional  $\mathbb{C}$ -vector space  $RT_r(\Sigma, c)$  is constructed in [7], as well as a projective representation:

$$\rho_{r,c} : \text{Mod}(\Sigma) \rightarrow \mathbb{P}\text{Aut}(RT_r(\Sigma, c)).$$

For different choices of root of unity the representations  $\rho_{r,c}$  are related by Galois group actions. In this paper, we will always choose  $\zeta_r = e^{\frac{i\pi}{r}}$ , which is important for us in order to apply results from [9, 10].

The representation  $\rho_{r,c}$  is called the  $SO(3)$ -quantum representation of  $\text{Mod}(\Sigma)$  at level  $r$ . These representations are known to be asymptotically faithful by the work of Andersen [1]. However the question of how well these representations reflect the geometry of the mapping class groups remains wide open.

---

*Date:* November 9, 2017.

Research supported by NSF Grants DMS-1404754 and DMS-1708249.

By the Nielsen-Thurston classification, mapping classes  $f \in \text{Mod}(\Sigma)$  are divided into three types: periodic, reducible and pseudo-Anosov. Furthermore, the type of  $f$  determines the geometric structure, in the sense of Thurston, of the 3-manifold obtained as mapping torus of  $f$ . In [2] Andersen, Masbaum and Ueno formulated the following conjecture and proved it when  $\Sigma$  is the four-holed sphere.

**Conjecture 1.1.** (AMU conjecture [2]) *Let  $\phi \in \text{Mod}(\Sigma)$  be a pseudo-Anosov mapping class. Then for any big enough level  $r$ , there is a choice of colors  $c$  of the components of  $\partial\Sigma$ , such that  $\rho_{r,c}(\phi)$  has infinite order.*

Note that it is known that the representations  $\rho_{r,c}$  send Dehn twists to elements of finite order and criteria for recognizing reducible mapping classes from their images under  $\rho_{r,c}$  are given in [3].

The results of [2] were extended by Egsgaard and Jorgensen in [11] and by Santharoubane in [21] to prove Conjecture 1.1 for some mapping classes of spheres with  $n \geq 5$  holes. In [20], Santharoubane proved the conjecture for the one-holed torus. However, until recently there were no known cases of the AMU conjecture for mapping classes of surfaces of genus at least 2. In [16], Marché and Santharoubane used skein theoretic techniques in  $\Sigma \times S^1$  to obtain such examples of mapping classes in arbitrary high genus. As explained by Koberda and Santharoubane [13], by means of Birman exact sequences of mapping class groups, one extracts representations of  $\pi_1(\Sigma)$  from the representations  $\rho_{r,c}$ . Elements in  $\pi_1(\Sigma)$  that correspond to pseudo-Anosov mappings classes via Birman exact sequences are characterized by a result of Kra [14]. Marché and Santharoubane used this approach to obtain their examples of pseudo-Anosov mappings classes satisfying the AMU conjecture by exhibiting elements in  $\pi_1(\Sigma)$  satisfying an additional technical condition they called Euler incompressibility. However, they informed us that they suspect their construction yields only finitely many mapping classes in any surface of fixed genus, up to mapping class group action.

The purpose of the present paper is to describe an alternative method for constructing mapping classes, acting on surfaces of arbitrary high genus, that satisfy the AMU conjecture. Our approach is to relate the conjecture with a question on the growth rate, with respect to  $r$ , of the  $SO(3)$ -Turaev-Viro 3-manifold invariants  $TV_r$ .

For  $M$  a compact orientable 3-manifold, closed or with boundary, the invariants  $TV_r(M)$  are real-valued topological invariants of  $M$ , that can be computed from state sums over triangulations of  $M$  and are closely related to the  $SO(3)$ -Witten-Reshetikhin-Turaev TQFTs. For a compact 3-manifold  $M$  (closed or with boundary) we define:

$$lTV(M) = \liminf_{r \rightarrow \infty, r \text{ odd}} \frac{2\pi}{r} \log |TV_r(M, q = e^{\frac{2i\pi}{r}})|.$$

Let  $f \in \text{Mod}(\Sigma)$  be a mapping class represented by a pseudo-Anosov homeomorphism of  $\Sigma$  and let  $M_f = F \times [0, 1] /_{(x,1) \sim (f(x),0)}$  be the mapping torus of  $f$ .

**Theorem 1.2.** *Let  $f \in \text{Mod}(\Sigma)$  be a pseudo-Anosov mapping class and let  $M_f$  be the mapping torus of  $f$ . If  $lTV(M_f) > 0$ , then  $f$  satisfies the conclusion of the AMU conjecture.*

The proof of the theorem relies having on the properties of TQFT underlying the Witten-Reshetikhin-Turaev  $SO(3)$ -theory as developed in [7].

As a consequence of Theorem 1.2 whenever we have a hyperbolic 3-manifold  $M$  with  $lTV(M) > 0$  that fibers over the circle, then the monodromy of the fibration represents a mapping class that satisfies the AMU conjecture.

By a theorem of Thurston, a mapping class  $f \in \text{Mod}(\Sigma)$  is represented by a pseudo-Anosov homeomorphism of  $\Sigma$  if and only if the mapping torus  $M_f$  is hyperbolic. In [8] Chen and Yang conjectured that for any hyperbolic 3-manifold with finite volume  $M$  we should have  $lTV(M) = \text{vol}(M)$ . Their conjecture implies, in particular, that the aforementioned technical condition  $lTV(M_f) > 0$  is true for all pseudo-Anosov mapping classes  $f \in \text{Mod}(\Sigma)$ . Hence, the Chen-Yang conjecture implies the AMU conjecture.

In this paper we will be concerned with surfaces with boundary and mapping classes that appear as monodromies of fibered links in  $S^3$ . We show the following.

**Theorem 1.3.** *Let  $L \subset S^3$  be a link with  $lTV(S^3 \setminus L) > 0$ . Then there are fibered hyperbolic links  $L'$ , with  $L \subset L'$  and  $lTV(S^3 \setminus L') > 0$ , and such that the complement of  $L'$  fibers over  $S^1$  with fiber a surface of arbitrarily large genus. In particular, the monodromy of such a fibration gives a mapping class in  $\text{Mod}(\Sigma)$  that satisfies the AMU conjecture.*

In [9] the authors gave criteria for constructing 3-manifolds, and in particular link complements, whose  $SO(3)$ -Turaev-Viro invariants satisfy the condition  $lTV > 0$ . Starting from these links, and applying Theorem 1.3, we obtain fibered links whose monodromies give examples of mapping classes that satisfy Conjecture 1.1. However, the construction yields only finitely many mapping classes in the mapping class groups of fixed surfaces. This is because the links  $L'$  obtained by Theorem 1.3 are represented by closed homogeneous braids and it is known that there are only finitely many links of fixed genus and number of components represented that way. To obtain infinitely many mapping classes for surfaces of fixed genus and number of boundary components, we need to refine our construction. We do this by using Stallings twists and appealing to a result of Long and Morton [15] on compositions of pseudo-Anosov maps with powers of a Dehn twist. The general process is given in Theorem 5.4. As an application we have the following.

**Theorem 1.4.** *Let  $\Sigma$  denote an orientable surface of genus  $g$  and with  $n$ -boundary components. Suppose that either  $n = 2$  and  $g \geq 3$  or  $g \geq n \geq 3$ . Then there are infinitely many non-conjugate pseudo-Anosov mapping classes in  $\text{Mod}(\Sigma)$  that satisfy the AMU conjecture.*

The paper is organized as follows: In Section 2, we summarize results from the  $SO(3)$ -Witten-Reshetikhin-Turaev TQFT and their relation to Turaev-Viro invariants that we need in this paper. In Section 3.2, we discuss how to construct families of links whose  $SO(3)$ -Turaev-Viro invariants have exponential growth (i.e.  $lTV > 0$ ) and then we prove Theorem 1.2 that explains how this exponential growth relates to the AMU Conjecture. In Section 4, we describe a method to get hyperbolic fibered links with any given sublink and we prove Theorem 1.3. In Section 5, we explain how to refine the construction of Section 4 to get infinite families of mapping classes on fixed genus surfaces that satisfy the AMU Conjecture (see Theorem 1.4). We also provide an explicit construction that leads to Theorem 1.4.

**1.1. Acknowledgement.** We would like to thank Matthew Hedden, Julien Marché, Gregor Masbaum, Ramanujan Santharoubane and Tian Yang for helpful discussions. We also thank Tian Yang for bringing to our attention the connection between the volume conjecture of [8] and the AMU conjecture.

## 2. TQFT PROPERTIES AND QUANTUM REPRESENTATIONS

In this section, we summarize some properties of the  $SO(3)$ -Witten-Reshetikhin-Turaev TQFTs, which we introduce in the skein-theoretic framework of [7], and briefly discuss their relation to the  $SO(3)$ -Turaev-Viro invariants.

**2.1. Witten-Reshetikhin-Turaev  $SO(3)$ -TQFTs.** Given an odd  $r \geq 3$ , let  $U_r$  denote the set of even integers less than  $r - 2$ . A banded link in a manifold  $M$  is an embedding of a disjoint union of annuli  $S^1 \times [0, 1]$  in  $M$ , and a  $U_r$ -colored banded link  $(L, c)$  is a banded link whose components are colored by elements of  $U_r$ . For a closed, oriented 3-manifold  $M$ , the Reshetikhin-Turaev invariants  $RT_r(M)$  are complex valued topological invariants. They also extend to invariants  $RT_r(M, (L, c))$  of manifolds containing colored banded links. These invariants are part of a compatible set of invariants of compact surfaces and compact 3-manifolds, which is called a TQFT. Below we summarize the main properties of the theory that will be useful to us in this paper, referring the reader to [6, 7] for the precise definitions and details.

**Theorem 2.1.** ([7, Theorem 1.4]) *For any odd integer  $r \geq 3$  and any primitive  $2r$ -th root of unity  $\zeta_r$ , there is a TQFT functor  $RT_r$  with the following properties:*

- (1) *For  $\Sigma$  a compact oriented surface, and if  $\partial\Sigma \neq \emptyset$  a coloring  $c$  of  $\partial\Sigma$  by elements of  $U_r$ , there is a finite dimensional  $\mathbb{C}$ -vector space  $RT_r(\Sigma, c)$ , with a Hermitian form  $\langle, \rangle$ . Moreover for disjoint unions, we have*

$$RT_r(\Sigma \amalg \Sigma') = RT_r(\Sigma) \otimes RT_r(\Sigma').$$

- (2) *For  $M$  a closed compact oriented 3-manifold, containing a  $U_r$ -colored banded link  $(L, c)$ , the value  $RT_r(M, (L, c), \zeta_r) \in \mathbb{C}$  is the  $SO(3)$ -Reshetikhin-Turaev invariant at level  $r$ .*
- (3) *For  $M$  a compact oriented 3-manifold with  $\partial M = \Sigma$ , and  $(L, c)$  a  $U_r$ -colored banded link in  $M$ , the invariant  $RT_r(M, (L, c))$  is a vector in  $RT_r(\Sigma)$ . Moreover, for compact oriented 3-manifolds  $M_1, M_2$  with  $\partial M_1 = -\partial M_2 = \Sigma$ , we have*

$$RT_r(M_1 \cup_{\Sigma} M_2) = \langle RT_r(M_1), RT_r(M_2) \rangle.$$

Finally, for disjoint unions  $M = M_1 \amalg M_2$ , we have

$$RT_r(M) = RT_r(M_1) \otimes RT_r(M_2).$$

- (4) *For a cobordism  $M$  with  $\partial M = -\Sigma_0 \cup \Sigma_1$ , there is a map*

$$RT_r(M) \in \text{End}(RT_r(\Sigma_0), RT_r(\Sigma_1)).$$

- (5) *The composition of cobordisms is sent by  $RT_r$  to the composition of linear maps, up to a power of  $\zeta_r$ .*

In [7] the authors construct some explicit orthogonal basis  $E_r$  for  $RT_r(\Sigma, c)$ : Let  $\Sigma$  be a compact, oriented surface that is not the 2-torus or the 2-sphere with less than four holes. Let  $P$  be a collection of simple closed curves on  $\Sigma$  that contains the boundary  $\partial\Sigma$  and gives a pants decomposition of  $\Sigma$ . The elements of  $E_r$  are in one-to-one correspondence with colorings  $\hat{c} : P \rightarrow U_r$ , such that  $\hat{c}$  agrees with  $c$  on  $\partial\Sigma$  and for each pant the colors of the three boundary components satisfy certain admissibility conditions. We will not make use of the general construction. What we need is the following:

**Theorem 2.2.** ([7, Theorem 4.11, Corollary 4.10])

- (1) For  $\Sigma$  a compact, oriented surface, with genus  $g$  and  $n$  boundary components, such that  $(g, n) \neq (1, 0), (0, 0), (0, 1), (0, 2), (0, 3)$ , we have

$$\dim(RT_r(\Sigma, c)) \leq r^{3g-3+n}.$$

- (2) If  $\Sigma = T$  is the 2-torus we actually have an orthonormal basis for  $RT_r(T)$ . It consists of the elements  $e_0, e_2, \dots, e_{r-3}$ , where

$$e_i = RT_r(D^2 \times S^1, ([0, \frac{1}{2}] \times S^1, i))$$

is the Reshetikhin-Turaev vector of the solid torus with the core viewed as banded link and colored by  $i$ .

**2.2.  $SO(3)$ -quantum representations of the mapping class groups.** For any odd integer  $r \geq 3$ , any choice of a primitive  $2r$ -root of unity  $\zeta_r$  and a coloring  $c$  of the boundary components of  $\Sigma$  by elements of  $U_r$ , we have a finite dimensional projective representation,

$$\rho_{r,c} : \text{Mod}(\Sigma) \rightarrow \mathbb{P}\text{Aut}(RT_r(\Sigma, c)).$$

If  $\Sigma$  is a closed surface and  $f \in \text{Mod}(\Sigma)$ , we simply have  $\rho_r(f) = RT_r(C_\phi)$ , where the cobordism  $C_\phi$  is the mapping cylinder of  $f$  :

$$C_f = \Sigma \times [0, 1] \coprod_{(1,x) \sim f(x)} \Sigma.$$

The fact that this gives a projective representation of  $\text{Mod}(\Sigma)$  is a consequence of points (4) and (5) of Theorem 2.1.

For  $\Sigma$  with non-empty boundary, giving the precise definition of the quantum representations would require us to discuss the functor  $RT_r$  for cobordisms containing colored tangles (see [7]). Since in this paper we will only be interested in the traces of the quantum representations, we will not recall the definition of the quantum representations in its full generality. We will use the following theorem:

**Theorem 2.3.** For  $r \geq 3$  odd, let  $\Sigma$  be a compact oriented surface with  $c$  a  $U_r$ -coloring on the components of  $\partial\Sigma$ . Let  $\tilde{\Sigma}$  be the surface obtained from  $\Sigma$  by capping the components of  $\partial\Sigma$  with disks. For  $f \in \text{Mod}(\Sigma)$ , let  $\tilde{f} \in \text{Mod}(\tilde{\Sigma})$ , denote the mapping class of the extension of  $f$  on the capping disks by the identity. Let  $M_{\tilde{f}} = F \times [0, 1] /_{(x,1) \sim (\tilde{f}(x),0)}$  be the mapping torus of  $\tilde{f}$  and let  $L \subset M_{\tilde{f}}$  denote the link whose components consist of the cores of the solid tori in  $M_{\tilde{f}}$  over the capping disks. Then, we have

$$\text{Tr}(\rho_{r,c}(f)) = RT_r(M_{\tilde{f}}, (L, c)).$$

**2.3.  $SO(3)$ -Turaev-Viro invariants.** In [26], Turaev and Viro introduced invariants of compact oriented 3-manifolds as state sums on triangulations of 3-manifolds. The triangulations are colored by representations of a semi-simple quantum Lie algebra. In this paper, we are only concerned with the  $SO(3)$ -theory: Given a compact 3-manifold  $M$ , an odd integer  $r \geq 3$ , and a primitive  $2r$ -root of unity, there is an  $\mathbb{R}$ -valued invariant  $TV_r$ . We refer to [10] for the precise flavor of Turaev-Viro invariants we are using here, and to [26] and for the original definitions and proofs of invariance. We will make use the following theorem, which relates the

Turaev-Viro invariants  $TV_r(M)$  of a 3-manifold  $M$  with the Witten-Reshetikhin-Turaev TQFT  $RT_r$ . For closed 3-manifolds it was proved by Roberts [19], and was extended to manifolds with boundary by Benedetti and Petronio [5]. In fact, as Benedetti and Petronio formulated their theorem in the case of  $SU_2$ -TQFT, the adaptation of the proof in the setting of  $SO(3)$ -TQFT we use here can be found in [10].

**Theorem 2.4.** ([5, Theorem 3.2]) *For  $M$  an oriented compact 3-manifold with empty or toroidal boundary and  $r \geq 3$  an odd integer, we have:*

$$TV_r(M, q = e^{\frac{2i\pi}{r}}) = \|RT_r(M, \zeta = e^{\frac{i\pi}{r}})\|^2.$$

### 3. GROWTH OF TURAEV-VIRO INVARIANTS AND THE AMU CONJECTURE

In this section, first we explain how the growth of the  $SO(3)$ -Turaev-Viro invariants is related to the AMU conjecture. Then we give examples of link complements  $M$  for which the  $SO(3)$ -Turaev-Viro invariants have exponential growth with respect to  $r$ ; that is, we have  $lTV(M) > 0$ .

**3.1. Exponential growth implies the AMU conjecture.** Let  $\Sigma$  denote a compact orientable surface with or without boundary and, as before, let  $\text{Mod}(\Sigma)$  denote the mapping class group of  $\Sigma$  fixing the boundary.

**Theorem 1.2.** *Let  $f \in \text{Mod}(\Sigma)$  be a pseudo-Anosov mapping class and let  $M_f$  be the mapping torus of  $f$ . If  $lTV(M_f) > 0$ , then  $f$  satisfies the conclusion of the AMU conjecture.*

The proof of Theorem 1.2 relies on the following elementary lemma:

**Lemma 3.1.** *If  $A \in \text{GL}_n(\mathbb{C})$  is such that  $|\text{Tr}(A)| > n$ , then  $A$  has infinite order.*

*Proof.* Up to conjugation we can assume that  $A$  is upper triangular. If the sum of the  $n$  diagonal entries has modulus bigger than  $n$ , one of these entries must have modulus bigger than 1. This implies that  $A$  has infinite order.  $\square$

*Proof of Theorem 1.2.* Suppose that for the mapping torus  $M_f$  of some  $f \in \text{Mod}(\Sigma)$ , we have  $lTV(M_f) > 0$ . We will prove Theorem 1.2 by relating  $TV_r(M_f)$  to traces of the quantum representations of  $\text{Mod}(\Sigma)$ . By Theorem 2.4, we have

$$TV_r(M_f) = \|RT_r(M_f)\|^2 = \langle RT_r(M_f), RT_r(M_f) \rangle,$$

where, with the notation of Theorem 2.1,  $\langle, \rangle$  is the Hermitian form on  $RT_r(\Sigma, c)$ .

Suppose that  $\Sigma$  has genus  $g$  and  $n$  boundary components. Now  $\partial M_f$  is a disjoint union of  $n$  tori. Note that by Theorem 2.2-(2) and Theorem 2.1-(1),  $RT_r(\partial M_f)$  admits an orthonormal basis given by vectors

$$\mathbf{e}_c = e_{c_1} \otimes e_{c_2} \otimes \dots \otimes e_{c_n},$$

where  $c = (c_1, c_2, \dots, c_n)$  runs over all  $n$ -tuples of colors in  $U_r$ , one for each boundary component. By Theorem 2.1-(3) and Theorem 2.2-(2), this vector is also the  $RT_r$ -vector of the cobordism consisting of  $n$  solid tori, with the  $i$ -th solid torus containing the core colored by  $c_i$ .

We can write  $RT_r(M_f) = \sum_c \lambda_c \mathbf{e}_c$  where  $\lambda_c = \langle RT_r(M_f), \mathbf{e}_c \rangle$ . Thus we have

$$TV_r(M_f) = \sum_c |\lambda_c|^2 = \sum_c |\langle RT_r(M_f), \mathbf{e}_c \rangle|^2,$$

where  $\mathbf{e}_c$  is the above orthonormal basis of  $RT_r(\partial M_f)$  and the sum runs over  $n$ -tuples of colors in  $U_r$ . By Theorem 2.1-(3), the pairing  $\langle RT_r(M_f), \mathbf{e}_c \rangle$  is obtained by filling the boundary components of  $M_f$  by solid tori and adding a link  $L$  which is the union of the cores and the core of the  $i$ -th component is colored by  $c_i$ . Thus by Theorem 2.3, we have

$$\langle RT_r(M_f), \mathbf{e}_c \rangle = RT_r(M_{\tilde{f}}, (L, c)) = \text{Tr}(\rho_{r,c}(f)),$$

and thus

$$TV_r(M_f) = \sum_c |\text{Tr} \rho_{r,c}(f)|^2,$$

where the sum ranges over all colorings of the boundary components of  $M_f$  by elements of  $U_r$ . Now, on the one hand, since  $lTV(M_f) > 0$ , the sequence  $\{TV_r(M_f)\}_r$  is bounded below by a sequence that is exponentially growing in  $r$  as  $r \rightarrow \infty$ . On the other hand, by Theorem 2.2-(1), the sequence  $\sum_c \dim(RT_r(\Sigma, c))$  only grows polynomially in  $r$ . For big enough  $r$ , there will be at least one  $c$  such that  $|\text{Tr} \rho_{r,c}(f)| > \dim(RT_r(\Sigma, c))$ . Thus by Lemma 3.1,  $\rho_{r,c}(\phi)$  will have infinite order.  $\square$

By a theorem of Thurston [23], a mapping class  $f \in \text{Mod}(\Sigma)$  is represented by a pseudo-Anosov homeomorphism of  $\Sigma$  if and only if the mapping torus  $M_f$  is hyperbolic.

As a consequence of Theorem 1.2, whenever a hyperbolic 3-manifold  $M$  that fibers over the circle has  $lTV(M) > 0$ , the monodromy of the fibration represents a mapping class that satisfies the AMU Conjecture.

In the remaining of this paper we will be concerned with surfaces with boundary and mapping classes that appear as monodromies of fibered links in  $S^3$ .

**3.2. Link complements with  $lTV > 0$ .** Links with exponentially growing Turaev-Viro invariants will be the fundamental building block of our construction of examples of pseudo-Anosov mapping classes satisfying the AMU conjecture. We will need the following result proved by the authors in [9].

**Theorem 3.2.** ([9, Corollary 5.3]) *Assume that  $M$  and  $M'$  are oriented compact 3-manifolds with empty or toroidal boundaries and such that  $M$  is obtained by a Dehn-filling of  $M'$ . Then we have:*

$$lTV(M) \leq lTV(M').$$

Note that for a link  $L \subset S^3$ , and a sublink  $K \subset L$ , the complement of  $K$  is obtained from that of  $L$  by Dehn-filling. Thus Theorem 3.2 implies that if  $K$  is a sublink of a link  $L \subset S^3$  and  $lTV(S^3 \setminus K) > 0$ , then we have  $lTV(S^3 \setminus L) > 0$ .

**Corollary 3.3.** *Let  $K \subset S^3$  be the knot  $4_1$  or a link with complement homeomorphic to that of the Borromean links or the Whitehead link. If  $L$  is any link containing  $K$  as a sublink then  $lTV(S^3 \setminus L) > 0$ .*

*Proof.* Denote by  $B$  the Borromean rings. By [10],  $lTV(S^3 \setminus 4_1) = 2v_3 \simeq 2.02988$  and  $lTV(S^3 \setminus B) = 2v_8 \simeq 7.32772$ ; and hence the conclusion holds for  $B$  and  $4_1$ . The complement of  $K = 4_1$  is obtained by Dehn filling along one of the components of the Whitehead link  $W$ . Thus, by Theorem 3.2,  $lTV(S^3 \setminus W) \geq 2v_3 > 0$ . For links with homeomorphic complements the conclusion follows since the Turaev-Viro invariants are homeomorphism invariants

of the link complement; that is, they will not distinguish different links with homeomorphic complements.  $\square$

**Remark 3.4.** Additional classes of links with  $lTV > 0$  are given by the authors in [10] and [9]. Some of these examples are non-hyperbolic. However it is known that any link is a sublink of a hyperbolic link [4]. Thus one can start with any link  $K$  with  $lTV(S^3 \setminus K) > 0$  and construct hyperbolic links  $L$  containing  $K$  as sublink; by Theorem 3.2 these will still have  $lTV(S^3 \setminus L) > 0$ .

#### 4. A HYPERBOLIC VERSION OF STALLINGS'S HOMOGENIZATION

A classical result of Stallings [22] states that every link  $L$  is a sublink of fibered links with fibers of arbitrarily large genera. Our purpose in this section is to prove the following hyperbolic version of this result.

**Theorem 4.1.** *Given a link  $L \subset S^3$ , there are hyperbolic links  $L'$ , with  $L \subset L'$  and such that the complement of  $L'$  fibers over  $S^1$  with fiber a surface of arbitrarily large genus.*

**4.1. Homogeneous braids.** Let  $\sigma_1, \dots, \sigma_{n-1}$  denote the standard braid generators of the  $n$ -strings braid group  $B_n$ . We recall that a braid  $\sigma \in B_n$  is said to be *homogeneous* if each standard generator  $\sigma_i$  appearing in  $\sigma$  always appears with exponents of the same sign. In [22], Stallings studied relations between closed homogeneous braids and fibered links. We summarize his results as follows:

- (1) The closure of any homogeneous braid  $\sigma \in B_n$  is a fibered link: The complement fibers over  $S^1$  with fiber the surface  $F$  obtained by Seifert's algorithm from the homogeneous closed braid diagram. The Euler characteristic of  $F$  is  $\chi(F) = n - c(\sigma)$  where  $c(\sigma)$  is the number of crossings of  $\sigma$ .
- (2) Given a link  $L = \hat{\sigma}$  represented as the closure of a braid  $\sigma \in B_n$ , one can add additional strands to obtain a homogeneous braid  $\sigma' \in B_{n+k}$  so that the closure of  $\sigma'$  is a link  $L \cup K$ , where  $K$ , the closure of the additional  $k$ -strands, represents the unknot. Furthermore, we can arrange  $\sigma'$  so that the linking numbers of  $K$  with the components of  $\hat{\sigma}$  are any arbitrary numbers. The link  $L \cup K$ , as a closed homogeneous braid, is fibered.

Throughout the paper we will refer to the component  $K$  of  $L \cup K$ , as the Stallings component.

In order to prove Theorem 4.1, given a hyperbolic link  $L$ , we want to apply Stallings' homogenizing method in a way such that the resulting link is still hyperbolic.

Let  $L$  be a hyperbolic link with  $n$  components  $L_1, \dots, L_n$ . The complement  $M_L := S^3 \setminus L$  is a hyperbolic 3-manifold with  $n$  cusps; one for each component. For each cusp, corresponding to some component  $L_i$ , there is a conjugacy class of a rank two abelian subgroup of  $\pi_1(M_L)$ . We will refer to this as the *peripheral group* of  $L_i$ .

**Definition 4.2.** Let  $L$  be a hyperbolic link with  $n$  components  $L_1, \dots, L_n$ . We say that an unknotted circle  $K$  embedded in  $S^3 \setminus L$  satisfies condition ( $\clubsuit$ ) if (i) the free homotopy class  $[K]$  does not lie in a peripheral group of any component of  $L$ ; and (ii) we have

$$\gcd(lk(K, L_1), lk(K, L_2), \dots, lk(K, L_n)) = 1.$$



The rest of this subsection is devoted to the proof of the following proposition that is needed for the proof of Theorem 4.1.

**Proposition 4.3.** *Given a hyperbolic link  $L$ , one can choose the Stallings component  $K$  so that (i)  $K$  satisfies condition  $(\clubsuit)$ ; and (ii) the fiber of the complement of  $L \cup K$  has arbitrarily high genus.*

Since  $L$  is hyperbolic, we have the discrete faithful representation

$$\rho : \pi_1(S^3 \setminus L) \longrightarrow \mathrm{PSL}_2(\mathbb{C}).$$

We recall that an element of  $A \in \mathrm{PSL}_2(\mathbb{C})$  is called *parabolic* if  $\mathrm{Tr}(A) = \pm 2$ , and that  $\rho$  takes elements in the peripheral subgroups of  $\pi_1(S^3 \setminus L)$  to parabolic elements in  $\mathrm{PSL}_2(\mathbb{C})$ . Since matrix trace is invariant under conjugation, in the discussion below we will not make distinction between elements in  $\pi_1(M_L)$  and their conjugacy classes. With this understanding we recall that if an element  $\gamma \in \pi_1(S^3 \setminus L)$  satisfies  $\mathrm{Tr}(\rho(\gamma)) \neq \pm 2$ , then it does not lie in any peripheral subgroup [24, Chapter 5].

**Lemma 4.4.** *Let  $A$  and  $B$  be elements in  $\mathrm{PSL}_2(\mathbb{C})$ .*

- (1) *If  $A$  and  $B$  are non commuting parabolic elements then  $|\mathrm{Tr}(A^l B^{-l})| > 2$  for some  $l$ .*
- (2) *If  $|\mathrm{Tr}(A)| > 2$ , then  $|\mathrm{Tr}(A^k B)| \neq 2$  for all  $k$  big enough.*

*Proof.* For (1), note that after conjugation we can take  $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$ . Then

$$|\mathrm{Tr}(A^l B^{-l})| = |1 - l^2 x| \xrightarrow{l \rightarrow \infty} \infty.$$

For (2), after conjugation take  $A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$  where  $|\lambda| > 1$  and write  $B = \begin{pmatrix} u & v \\ w & x \end{pmatrix}$ . Then

$$\mathrm{Tr}(A^k B) = \lambda^k u + \lambda^{-k} x,$$

which as  $k \rightarrow +\infty$  tends either to infinity if  $u \neq 0$  or to 0 else. In the first case we will have  $|\mathrm{Tr}(A^k B)| > 2$  for  $k$  big enough; in the second case we will have  $|\mathrm{Tr}(A^k B)| < 2$ . In both cases we have  $|\mathrm{Tr}(A^k B)| \neq 2$  as desired.  $\square$

Next we will consider the Wirtinger presentation of  $\pi_1(S^3 \setminus L)$  corresponding to a link diagram representing a hyperbolic link  $L$  as a closed braid  $\hat{\sigma}$ . The Wirtinger generators are conjugates of meridians of the components of  $L$  and are mapped to parabolic elements of  $\mathrm{PSL}_2(\mathbb{C})$  by  $\rho$ . A key point in the proof of Proposition 4.3 is to choose the Stallings component  $K$  so that the word it represents in  $\pi_1(S^3 \setminus L)$  is conjugate to one that begins with a sub-word  $(a^l b^{-l})^k$ , where  $a$  and  $b$  are Wirtinger generators mapped to non-commuting elements under  $\rho$ . Then we will use Lemma 4.4 to prove that the free homotopy class  $[K]$  is not in a peripheral subgroup of any component of  $L$ . We first need the following lemma:

**Lemma 4.5.** *Let  $L$  be a hyperbolic link in  $S^3$ , with a link diagram of a closed braid  $\hat{\sigma}$ . We can find two strands of  $\sigma$  meeting at a crossing so that if  $a$  and  $b$  are the Wirtinger generators corresponding to an under-strand and the over-strand of the crossing respectively, then  $\rho(a)$  and  $\rho(b)$  don't commute.*




FIGURE 1. The four step homogenization process.

*Proof.* Suppose that for any pair of Wirtinger generators  $a, b$  corresponding to a crossing as above,  $\rho(a)$  and  $\rho(b)$  commute. Since  $\rho(a)$  and  $\rho(b)$  are commuting parabolic elements of infinite order in  $\mathrm{PSL}_2(\mathbb{C})$ , elementary linear algebra shows that they share their unique eigenline. Then step by step, we get that the images under  $\rho$  of all Wirtinger generators share an eigenline. But this would imply that  $\rho(\pi_1(S^3 \setminus L))$  is abelian which is a contradiction.  $\square$

We can now turn to the proof of Proposition 4.3, which we will prove by tweaking Stallings homogenization procedure.

*Proof of Proposition 4.3.* Let  $L$  be a hyperbolic link, with components,  $L_1, \dots, L_n$ , represented as a braid closure  $\hat{\sigma}$ . Let  $a, b$  be Wirtinger generators of  $\pi_1(S^3 \setminus L)$  chosen as in Lemma 4.5.

Starting with the projection of  $\hat{\sigma}$ , we proceed in the following way:

We arrange the crossings of  $\hat{\sigma}$  to occur at different verticals on the projection plane.

- (1) Begin drawing the Stallings component so that near the strands where above chosen Wirtinger generators  $a, b$  occur, we create the pattern shown the left of Figure 2.
- (2) We deform the strands of  $\sigma$  to create “zigzags” as shown in the second drawing of Figure 1.
- (3) We fill the empty spaces in verticals with new braid strands and choose the new crossings so that the resulting braid is homogeneous and so that the new strands meet the strands of  $\sigma$  both in positive and negative crossings. Adding enough “zigzags” at the previous step will ensure that there is enough freedom in choosing the crossings to make this second condition possible.
- (4) At this stage, we have turned the braid  $\sigma$  into a homogeneous braid, say  $\sigma_h$ . The closure  $\hat{\sigma}_h$  contains  $L$  as a sublink and some number  $s \geq 1$  of unknotted components. To reduce the number of components added, we connect the new components with a single crossing between each pair of neighboring new components. Doing so we may have to create new crossings with the components of  $L$ , but we can always choose them to preserve homogeneity. Thus we homogenized  $L$  by adding a single unknotted component  $K$  to it.

The four step process described above is illustrated in Figure 1.




FIGURE 2. Changing  $[K]$  from a conjugate of  $w \in \pi_1(S^3 \setminus L)$  (left), to a conjugate of  $w(a^{-l}b^l)^k$  (right), for any non-negative  $l$  and  $k$ . Here  $l = k = 2$ .

Now, because we have positive and negative crossings of  $K$  with each component of  $L$ , we can set the linking numbers as we want just by adding an even number of positive or negative crossings between  $K$  and a component of  $L$  locally. If the strands  $a$  and  $b$  correspond to the same component  $L_1$ , we simply ask that  $lk(K, L_1) = 1$ . If they correspond to two distinct components  $L_1$  and  $L_2$ , we choose  $(lk(K, L_1), lk(K, L_2)) = (1, 0)$ .

Recall that we have chosen  $a, b$  to be Wirtinger generators of  $\pi_1(S^3 \setminus L)$ , as in Lemma 4.5, and so that  $K$  is added to  $L$  so that the pattern shown on the left hand side of Figure 2 occurs near the corresponding crossing. Assume that  $[K]$  is conjugate to a word  $w \in \pi_1(S^3 \setminus L)$ . Now one may modify the diagram of  $L \cup K$  locally, as shown in the right hand side of Figure 2, to make  $[K]$  conjugate to  $(a^{-l}b^l)^k w$  for any non-negative  $k$  and  $l$ . Notice also this move leaves  $K$  unknotted and that  $L \cup K$  is still a closed homogeneous braid. Also notice that doing so, we left  $lk(K, L_1)$  unchanged if  $a$  and  $b$  were part of the same component  $L_1$ , and we turned  $(lk(K, L_1), lk(K, L_2))$  into  $(1 - kl, kl)$  if they correspond to different components  $L_1$  and  $L_2$ . In both cases, we preserved the fact that

$$\gcd(lk(K, L_1), lk(K, L_2), \dots, lk(K, L_n)) = 1$$

and  $K$  satisfies part (ii) of Condition ( $\clubsuit$ ).

To ensure that part (i) of the condition is satisfied, note that since  $\rho(a)$  and  $\rho(b)$  are non-commuting, Lemma 4.4 (1) implies  $|\text{Tr}(\rho(a)^{-l}\rho(b)^l)| > 2$  for  $l \gg 0$ . Thus by choosing  $k \gg 0$ , and using Lemma 4.4 (2), we may assume that  $|\text{Tr}(A^k B)| \neq 2$ , where  $A = \rho(a)^{-l}\rho(b)^l$  and  $B = \rho(w)$ . Then  $[K] = A^k B$  is not in a peripheral subgroup of  $\pi_1(S^3 \setminus L)$ .

Now notice that as above mentioned positive integers  $k, l$  become arbitrarily large, the crossing number of the resulting homogeneous braid projections becomes arbitrarily large while the braid index remains unchanged. Since the fiber of the fibration of a closed homogeneous braid is the Seifert surface of the closed braid projection it follows that as  $k, l \rightarrow \infty$ , the genus of the fiber becomes arbitrarily large.  $\square$

**4.2. Ensuring hyperbolicity.** In this subsection we will finish the proof of Theorem 4.1. For this we need the following:

**Proposition 4.6.** *Suppose that  $L$  is a hyperbolic link and let  $L \cup K$  be a homogeneous closed braid obtained from  $L$  by adding a Stallings component  $K$  that satisfies condition  $(\clubsuit)$ . Then  $L \cup K$  is a hyperbolic link.*

Before we can proceed with the proof of Proposition 4.6 we need some preparation: We recall that when an oriented link  $L$  is embedded in a solid torus, the total winding number of  $L$  is the non-negative integer  $n$  such that  $L$  represents  $n$  times a generator of  $H_1(V, \mathbb{Z})$ . When convenient we will consider  $M_{L \cup K}$  to be the compact 3-manifold obtained by removing the interiors of neighborhoods of the components of  $L \cup K$ ; the interior of  $M_{L \cup K}$  is homeomorphic to  $S^3 \setminus (L \cup K)$ . In the course of the proof of the proposition we will see that condition  $(\clubsuit)$  ensures that the complement of  $L \cup K$  cannot contain embedded tori that are not boundary parallel or compressible (i.e.  $M_{L \cup K}$  is *atoroidal*). We need the following lemma that provides restrictions on winding numbers of satellite fibered links.

**Lemma 4.7.** *We have the following:*

- (1) *Suppose that  $L$  is a oriented fibered link in  $S^3$  that is embedded in a solid torus  $V$  with boundary  $T$  incompressible in  $S^3 \setminus L$ . Then, some component of  $L$  must have non-zero winding number.*
- (2) *Suppose that  $L$  is an oriented fibered link in  $S^3$  such that only one component  $K$  is embedded inside a solid torus  $V$ . If  $K$  has winding number 1, then  $K$  is isotopic to the core of  $V$ .*

Though this statement is fairly classical in the context of fibered knots [12], we include a proof as we are working with fibered links.

*Proof.* The complement  $M_L = S^3 \setminus N(L)$  fibers over  $S^1$  with fiber a surface  $(F, \partial F) \subset (M_L, \partial M_L)$ . Then  $S^3 \setminus L$  cut along  $F = F \times \{0\} = F \times \{1\}$  is homeomorphic to  $F \times [0, 1]$ . It is known that  $F$  maximizes the Euler characteristic in its homology class in  $H_2(M_L, \partial M_L)$  and thus  $F$  is incompressible and  $\partial$ -incompressible.

(1) Assume that the winding number of every component of  $L$  is zero, and consider the intersection of  $F$  with  $T$ , the boundary of the solid torus containing  $L$ . Since  $F \times [0, 1]$  is irreducible, and  $F$  is incompressible in the complement of  $L$ , up to isotopy, one can assume that the intersection  $T \cap F$  consists of a collection of parallel curves in  $T$ , each of which is homotopically essential in  $T$ . The hypothesis on the winding number implies that the intersection  $F \cap T$  is null-homologous in  $T$ , where each component of  $F \cap T$  is given the orientation inherited by the surface  $V \cap F$ . Thus the curves in  $F \cap T$  can be partitioned in pairs of parallel curves with opposite orientations in  $T \cap F$ . Each such pair bounds an annulus in  $T$  and in  $F \times (0, 1)$  each of these annuli has both ends on  $F \times \{0\}$  or on  $F \times \{1\}$ . This implies that we can find  $0 < t < 1$  such that  $F_t = F \times \{t\}$  misses the torus  $T$ . This in turn implies that  $T$  must be an essential torus in the manifold obtained by cutting  $S^3 \setminus L$  along the fiber  $F_t$ . But this is impossible since the later manifold is  $F_t \times I$  which is a handlebody and cannot contain essential tori; contradiction.

(2) By an argument similar to that used in case (1) above, we can simplify the intersection of the fiber surface  $F$  with  $T$  until it consists of one curve only. This curve, say  $\gamma$ , cuts  $T$  into

an essential annulus embedded in  $F \times (0, 1)$  with one boundary component on  $F \times \{0\}$  and the other on  $F \times \{1\}$ . As the annulus closes up, the curve  $\gamma$  must be fixed by the monodromy of the fibration and one can isotope  $T$  to make it compatible with the fibration. Then one has that  $K$  is fibered in  $V$ , and as the winding number of  $K$  is 1, by Corollary 1 in [12],  $K$  must be isotopic to the core of  $V$ .  $\square$

We are now ready to give the proof of Proposition 4.6.

*Proof of Proposition 4.6.* First we remark that  $S^3 \setminus (L \cup K)$  is non-split as  $S^3 \setminus L$  is and  $K$  represents a non-trivial element in  $\pi_1(S^3 \setminus L)$ .

Next we argue that  $S^3 \setminus (L \cup K)$  is atoroidal: Assume that we have an essential torus, say  $T$ , in  $M_{L \cup K} = S^3 \setminus (L \cup K)$ . Since  $L$  is hyperbolic, in  $M_L = S^3 \setminus L$  the torus  $T$  becomes either boundary parallel or compressible. Moreover, the torus  $T$  bounds a solid torus  $V$  in  $S^3$ .

Suppose that  $T$  becomes boundary parallel in the complement of  $L$ . Then, we may assume that  $V$  is a tubular neighborhood of a component  $L_i$  of  $V$ . Then  $K$  must lie inside  $V$ ; for otherwise  $T$  would still be boundary parallel in  $M_{L \cup K}$ . Then the free homotopy class  $[K]$  would represent a conjugacy class in the peripheral subgroup of  $\pi_1(M_L)$  corresponding to  $L_i$ . However this contradicts condition  $(\clubsuit)$ ; thus this case cannot happen.

Suppose now that we know that  $T$  becomes compressible in  $M_L$ . In  $S^3$ , the torus  $T$  bounds a solid torus  $V$  that contains a compressing disk of  $T$  in  $M_L$ . If  $V$  contains no component of  $L \cup K$ , the torus  $T$  is still compressible in  $M_{L \cup K}$ . Otherwise, there are again two cases:

*Case 1:* The solid torus  $V$  contains some components of  $L$ . We claim that  $V$  actually contains all the components of  $L$ . Otherwise, after compressing  $T$  in  $M_L$ , one would get a sphere that separates the components of  $L$ , which can not happen as  $L$  is non-split. Moreover, as the compressing disk is inside  $V$ , all components of  $L$  have winding number zero in  $V$ .

Since  $T$  is incompressible in the complement of  $M_{L \cup K}$ , the component  $K$  must also lie inside  $V$ . Note that  $V$  has to be knotted since otherwise  $T$  would compress outside  $V$  and thus in  $M_{L \cup K}$ . But then since  $K$  is unknotted, it must have winding number zero in  $V$ . Thus we have the fibered link  $L \cup K$  lying inside  $V$  so that each component has winding number zero. But then  $T$  can not be incompressible in  $M_{L \cup K}$  by Lemma 4.7-(1); contradiction. Thus this case will not happen.

*Case 2:* The solid torus  $V$  contains only  $K$ . Since  $T$  is incompressible in  $M_{L \cup K}$ ,  $K$  must be geometrically essential in  $V$ ; that is it doesn't lie in a 3-ball inside  $V$ . Since  $K$  is unknotted, it follows that  $V$  is unknotted. For each component  $L_i$  of  $L$ , we have

$$lk(K, L_i) = w \cdot lk(c, L_i),$$

where  $c$  is the core of  $V$ , and  $w$  denote the winding number of  $K$  in  $V$ . Since  $K$  satisfies condition  $(\clubsuit)$ , we know that

$$\gcd(lk(K, L_1), lk(K, L_2), \dots, lk(K, L_n)) = 1,$$

which implies that we must have  $w = 1$ . Thus by Lemma 4.7-(2),  $K$  is isotopic to the core of  $V$  and  $T$  is boundary parallel, contradicting the assumption that  $T$  is essential in  $M_{L \cup K}$ . This finishes the proof that  $M_{L \cup K}$  is atoroidal.

Since  $M_{L \cup K}$  contains no essential spheres or tori, and has toroidal boundary, it is either a Seifert fibered space or a hyperbolic manifold. But  $M_L$  is a Dehn-filling of  $M_{L \cup K}$  which is hyperbolic. Since the Gromov norm  $\|\cdot\|$  does not increase under Dehn filling [24] we get

$\|M_{L \cup K}\| \geq \|M_L\| > 0$ . The Gromov norm of Seifert 3-manifolds is zero, thus  $L \cup K$  must be hyperbolic.  $\square$

We can now finish the proof of Theorem 4.1 and the proof of Theorem 1.3 stated in the Introduction.

*Proof of Theorem 4.1.* Let  $L$  be any link. If  $L$  is not hyperbolic, then we can find a hyperbolic link  $L'$ , that contains  $L$  as a sub-link. See for example [4]. If  $L$  is hyperbolic then set  $L = L'$ . Then apply Proposition 4.3 to  $L'$  to get links  $L' \cup K$  that are closed homogeneous braids with arbitrarily high crossing numbers and fixed braid index. By [22], the links  $L' \cup K$  are fibered and the fibers have arbitrarily large genus and by Proposition 4.6 they are hyperbolic.  $\square$

*Proof of Theorem 1.3.* Suppose that  $L$  is a link with  $lTV(S^3 \setminus L) > 0$ . By Theorem 4.1 we have fibered hyperbolic links  $L'$  that contain  $L$  as sublink and whose fibers have arbitrarily large genus. By Theorem 3.2 we have  $lTV(S^3 \setminus L) > 0$ .  $\square$

## 5. STALLINGS TWISTS AND THE AMU CONJECTURE

By our results in the previous sections, starting from a hyperbolic link  $L \subset S^3$  with  $lTV(S^3 \setminus L) > 0$ , one can add an unknotted component  $K$  to obtain a hyperbolic fibered link  $L \cup K$ , with  $lTV(S^3 \setminus (L \cup K)) > 0$ . The monodromy of a fibration of  $L \cup K$  provides a pseudo-Anosov mapping class on the surface  $\Sigma = \Sigma_{g,n}$ , where  $g$  is the genus of the fiber and  $n$  is the number of components of  $L \cup K$ .

One can always increase the number of boundary components  $n$  by adding more components to  $L$  and appealing to Theorem 3.2. However since  $L \cup K$  is a closed homogeneous braid this construction alone will not provide infinite families of examples for fixed genus and number of boundary components.

In this section we show how to address this problem and prove Theorem 1.4 stated in the introduction and which, for the convenience of the reader we restate here.

**Theorem 1.4.** *Let  $\Sigma$  denote an orientable surface of genus  $g$  and with  $n$ -boundary components. Suppose that either  $n = 2$  and  $g \geq 3$  or  $g \geq n \geq 3$ . Then there are infinitely many non-conjugate pseudo-Anosov mapping classes in  $\text{Mod}(\Sigma)$  that satisfy the AMU conjecture.*

**5.1. Stallings twists and pseudo-Anosov mappings.** Stallings [22] introduced an operation that transforms a fibered link into a fibered link with a fiber of the same genus: Let  $L$  be a fibered link with fiber  $F$  and let  $c$  be a simple closed curve on  $F$  that is unknotted in  $S^3$  and such that  $lk(c, c^+) = 0$ , where  $c^+$  is the curve  $c$  pushed along the normal of  $F$  in the positive direction. The curve  $c$  bounds a disk  $D \subset S^3$  that is transverse to  $F$ . Let  $L_m$  denote the link obtained from  $L$  by a full twist of order  $m$  along  $D$ . This operation is known as Stallings twist of order  $m$ . Alternatively, one can think the Stallings twist operation as performing  $1/m$  surgery on  $c$ , where the framing of  $c$  is induced by the normal vector on  $F$ .

**Theorem 5.1.** ([22, Theorem 4]) *Let  $L$  be a link whose complement fibers over  $S^1$  with fiber  $F$  and monodromy  $f$ . Let  $L_m$  denote a link obtained by a Stallings twist of order  $m$  along a curve  $c$  on  $F$ . Then, the complement of  $L_m$  fibers over  $S^1$  with fiber  $F$  and the monodromy is  $f \circ \tau_c^m$ , where  $\tau_c$  is the Dehn-twist on  $F$  along  $c$ .*




FIGURE 3. A crossing and its  $A$  and  $B$  resolutions.




FIGURE 4. Left: Pattern in the state graph exhibiting a non-trivial Stallings twist. Right: The curve  $c$  obtained as a connected sum the curves  $c_1, c_2$  with self linking  $+2$  and  $-2$ .

Note that when  $c$  is parallel to a component of  $L$ , then such an operation does not change the homeomorphism class of the link complement; we call these Stallings twists trivial.

To facilitate the identification of non-trivial Stallings twists on link fibers, we recall the notion of *state graphs*:

Recall that the fiber for the complement of a homogeneous closed braid  $\hat{\sigma}$  is obtained as follows: Resolve all the crossings in the projection of  $\hat{\sigma}$  in a way consistent with the braid orientation. The result is a collection of nested embedded circles (Seifert circles) each bounding a disk on the projection plane; the disks can be made disjoint by pushing them slightly above the projection plane. Then we construct the fiber  $F$  by attaching a half twisted band for each crossing. The state graph consists of the collection of the Seifert circles together with an edge for each crossing of  $\hat{\sigma}$ . We will label each edge by  $A$  or  $B$  according to whether the resolution of the corresponding crossing during the construction of  $F$  is of type  $A$  or  $B$  shown in Figure 3, if viewed as unoriented resolution.

**Remark 5.2.** As the homogeneous braids get more complicated the fiber is more likely to admit a non-trivial Stallings twist. Indeed, if the state graph of  $L = \hat{\sigma}$  exhibits the local pattern shown in the left hand side of Figure 4, we can perform a non-trivial Stallings twist along the curve  $c$  which corresponds to the connected sum of the two curves  $c_1$  and  $c_2$  shown in the Figure. We can see that  $lk(c_1, c_1^+) = +2$  and  $lk(c_2, c_2^+) = -2$ , and the mixed linkings are zero. In the end,  $lk(c, c^+) = 2 - 2 = 0$ .

We will need the following theorem, stated and proved by Long and Morton [15] for closed surfaces. Here we state the bounded version and for completeness we sketch the slight adaptation of their argument in this setting.

**Theorem 5.3.** ([15, Theorem A]) *Let  $F$  be a compact oriented surface with  $\partial F \neq \emptyset$ . Let  $f$  be a pseudo-Anosov homeomorphism on  $F$  and let  $c$  be a non-trivial, non-boundary parallel simple closed curve on  $F$ . Let  $\tau_c$  denote the Dehn-twist along  $c$ . Then, the family  $\{f \circ \tau_c^m\}_m$  contains infinitely many non-conjugate pseudo-Anosov homeomorphisms.*

*Proof.* The proof rests on the fact that the mapping torus of  $f_m = f \circ \tau_c^m$  is obtained from  $M_f$  by performing  $1/m$ -surgery on the curve  $c$  with framing induced by a normal vector in  $F$ . Once we prove that  $M_f \setminus c$  is hyperbolic, Thurston's hyperbolic Dehn surgery theorem implies, for  $m$  big enough, that the mapping tori  $M_{f_m}$  are hyperbolic and all pairwise non-homeomorphic (as their hyperbolic volumes differ). Since conjugate maps have homeomorphic mapping tori the non-finiteness statement follows.

We will consider the curve  $c$  as embedded on the fiber  $F \times \{1/2\} \subset M_f$ . Notice that  $M_f \setminus c$  is irreducible, as  $c$  is non-trivial in  $\pi_1(F)$  and thus in  $\pi_1(M_f)$ .

We need to show that  $M_f \setminus c$  contains no essential embedded tori: Let  $T$  be a torus embedded in  $M_f \setminus c$ . If  $T$  is boundary parallel in  $M_f$ , it will also be in  $M_f \setminus c$ , otherwise one would be able to isotope  $c$  onto the boundary of  $M_f$ , and as  $c$  is actually a curve on  $F \times \{1/2\}$ ,  $c$  would be conjugate in  $\pi_1(F)$  to a boundary component. As we chose  $c$  non-boundary parallel in  $F$ , this does not happen.

Now, assume that  $T$  is non-boundary parallel in  $M_f$ . Then we can put  $T$  in general position and consider of  $T \cap F \times \{1/2\}$ . If this intersection is empty then  $T$  is compressible as  $F \times [0, 1]$  does not contain any essential tori. After isotopy we can assume that  $T \cap F \times [0, 1]$  is a collection of properly embedded annuli in  $F \times [0, 1]$ , each of which either misses a fiber  $F$  or is vertical with respect to the  $I$ -bundle structure. Now note that if one of these annuli misses a fiber then we can remove it by isotopy in  $M_f \setminus c$ , unless if it connects to curves parallel to  $c$  on opposite sides of  $c$  on  $F \times \{1/2\}$ . Also observe that we cannot have annuli that connect a non-boundary parallel curve  $c \subset F \times \{1/2\}$  to  $f(c)$ : For, since  $f$  is pseudo-Anosov, the curves  $f^k(c)$  and  $f^l(c)$  are freely homotopic on the fiber if and only if  $k = l$ ; and thus the annuli would never close up to give  $T$ . In the end, and since  $M_f$  is hyperbolic, we are left with two annuli connecting both sides of  $c$  and  $T$  is boundary parallel in  $M_f \setminus c$ .

Finally,  $M_f \setminus c$  is irreducible and atoroidal and since its Gromov norm satisfies  $\|M_f \setminus c\| > \|M_f\| > 0$ , it is hyperbolic.  $\square$

**5.2. Infinite families of mapping classes.** We are now ready to present our examples of infinite families of non-conjugate pseudo-Anosov mapping classes of fixed surfaces that satisfy the AMU conjecture. The following theorem gives the general process of the construction.

**Theorem 5.4.** *Let  $L$  be a hyperbolic fibered link with fiber  $\Sigma$  and monodromy  $f$ . Suppose that  $L$  contains a sublink  $K$  with  $lTV(S^3 \setminus K) > 0$ . Suppose, moreover, that the fiber  $\Sigma$  admits a non-trivial Stallings twist along a curve  $c \subset \Sigma$  such that the interior of the twisting disc  $D$  intersects  $K$  at most once geometrically. Let  $\tau_c$  denote the Dehn twist of  $\Sigma$  along  $c$ . Then the family  $\{f \circ \tau_c^m\}_m$  of homeomorphisms gives infinitely many non-conjugate pseudo-Anosov mappings classes in  $\text{Mod}(\Sigma)$  that satisfy the AMU conjecture.*






FIGURE 5. The link  $L_{4,m}$  and the state graph for  $L_{4,2}$ .

*Proof.* Since  $L$  contains  $K$  as sublink we have  $lTV(S^3 \setminus L) \geq lTV(S^3 \setminus K) > 0$ . Since  $D$  intersects  $K$  at most once, each of the links  $L'$  obtained by Stalling twists along  $c$ , will also contain a sublink isotopic to  $K$  and hence  $lTV(S^3 \setminus L') \geq lTV(S^3 \setminus K) > 0$ . The conclusion follows by Theorems 5.1, 5.3 and 1.2.  $\square$

We finish the section with concrete constructions of infinite families obtained by applying Theorem 5.4. Start with  $K_1 = 4_1$  represented as the closure of the homogeneous braid  $\sigma_2^{-1}\sigma_1\sigma_2^{-1}\sigma_1$ . We construction a 2-parameter family of links  $L_{n,m}$  where  $n \geq 2, m \geq 1$ , defined as follows:

The link  $L_{4,m}$  is shown in the left panel of Figure 5, where the box shown contains  $2m$  crossings. It is obtained from  $K_1$  by adding three unknotted components.

The link  $L_{3,m}$  is obtained from  $L_{4,m}$  by removing the unknotted component corresponding to the outermost string of the braid.

The link  $L_{n+1,m}$  for  $n \geq 4$  is obtained from  $L_{n,m}$  adding one strand in the following way: denote by  $K_1, \dots, K_n$  the components of  $L_{n,m}$  from innermost to outermost,  $K_1$  being the  $4_1$  component. To get  $L_{n+1,m}$ , we add one strand  $K_{n+1}$  to  $L_{n,m}$ , so that traveling along  $K_n$  one finds 2 crossings with  $K_{n-1}$ , then 2 crossings with  $K_{n+1}$ , then 2 crossings with  $K_{n-1}$ , then 2 crossings with  $K_{n+1}$ , and, moreover, the crossings with  $K_{n-1}$  and  $K_{n+1}$  have opposite signs. There is only one way to chose this new strand, and doing so we added one unknotted component to  $L_{n,m}$ , thus  $L_{n+1,m}$  has  $n + 1$  components and 4 more crossings than  $L_{n,m}$ .

In the special case  $n = 2$ , the link  $L_{2,m}$  is obtained from the link  $L_{3,m}$  by replacing the box with  $2m$  crossings with a box with  $2m - 1$  crossings. The links  $L_{2,m}$  are then 2-components links. We note that all the links  $L_{n,m}$  contain the component  $K_1$  we started with.

**Proposition 5.5.** *The link  $L_{n,m}$  is hyperbolic, fibered and satisfies the hypotheses of Theorem 5.4. The fiber has genus  $g = m + 2$  if  $n = 2$ ,  $g = n + m - 1$  otherwise.*

*Proof.* For every  $n \geq 2$  and any  $m \geq 1$ , the link  $L_{n,m}$  contains the knot  $K_1 = 4_1$  as sublink and as said earlier we have  $lTV(S^3 \setminus K_1) > 0$ . Since  $L_{n,m}$  is alternating, hyperbolicity follows from Menasco's criterion [17]: any prime non-split alternating diagram of a link that is not the standard diagram of the  $T(2, q)$  torus link, represents a hyperbolic link. Since  $L_{n,m}$  is represented by an alternating (and thus homogeneous) closed braid, fiberedness follows from Stallings's criterion. For  $n \geq 3$ , the resulting closed braid diagram has braid index  $2 + n$  and  $2 + 4n + 2m$  crossings. Hence the Euler characteristic is  $-3n - 2m$  and the genus is  $m + n - 1$ , as the fiber has  $n$  boundary components. In the case  $n = 2$ , the braid index is 5, number of crossings  $9 + 2m$ , thus the Euler characteristic of the fiber is  $-4 - 2m$  and the genus is  $m + 2$ . Using Remark 5.2 and the state graph given in Figure 5 we can easily locate a simple closed curve  $c$  on the fiber with the properties in the statement of Theorem 5.4.  $\square$

Now Proposition 5.5 and Theorem 5.4 immediately give Theorem 1.4 stated in the beginning of the section.

**Remark 5.6.** Note that if we restrict ourselves to closed homogeneous braids to get examples of links satisfying the hypotheses of Theorem 5.4, it seems necessary that the genus will grow with the number of components. However, one could consider other methods, that can increase the number of components, while keeping the fiber genus low, to produce links satisfying the hypotheses of Theorem 5.4. One way would be to take a Murasugi sum of the link  $L_{2,m}$  with links of arbitrarily large number of components and whose complement fibers over  $S^1$  with fiber of genus 0. This should be done so that the Murasugi sum operation leaves the component  $4_1 \subset L_{2,m}$  unaffected and it produces hyperbolic links. It seems plausible that combining homogeneous braids and Murasugi sums should give explicit examples of infinite families of mapping classes that satisfy the AMU Conjecture, for all surfaces  $\Sigma$  with genus  $g \geq 2$  and  $n \geq 2$  boundary components.

## REFERENCES

- [1] J. E. Andersen, *Asymptotic faithfulness of the quantum  $SU(n)$  representations of the mapping class groups*, Ann. of Math. **163** (2006), no. 1, 347–368.
- [2] J. E. Andersen, G. Masbaum, and K. Ueno, *Topological quantum field theory and the Nielsen-Thurston classification of  $M(0,4)$* , Math Proc. Cambridge Philos. Soc. **141** (2006), 477–488.
- [3] Andersen, J. E., *The Nielsen-Thurston classification of mapping classes is determined by TQFT*, J. Math. Kyoto Univ. **48** (2008), no. 2, 323–338.
- [4] M.D. Baker, *All links are sublinks of arithmetic links*, Pacific J. Math **203** (2002), no. 2, 257–263.
- [5] R. Benedetti and C. Petronio, *On Roberts' proof of the Turaev-Walker theorem*, J. Knot Theory Ramifications **5** (1996), no. 4, 427–439.
- [6] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Three-manifold invariants derived from the Kauffman bracket*, Topology **31** (1992), no. 4, 685–699.
- [7] ———, *Topological quantum field theories derived from the Kauffman bracket*, Topology **34** (1995), no. 4, 883–927.
- [8] Q. Chen and T. Yang, *Volume conjectures for the Reshetikhin-Turaev and the Turaev-Viro invariants*, arXiv:1503.02547, To appear in Quantum Topology.
- [9] R. Detcherry and E. Kalfagianni, *Gromov norm and Turaev-Viro invariants of 3-manifolds*, Arxiv 1705.09964.
- [10] R. Detcherry, E. Kalfagianni, and T. Yang, *Turaev Viro knot invariants and the colored Jones polynomials*, Arxiv 1701.07818.
- [11] J.K. Egsgaard and S.F. Jorgensen, *The homological content of the jones representations at  $q=-1$* , Journal of Knot Theory and its ramifications. **25** (2016), no. 11, 25pp.

- [12] M. Hirasawa, K. Murasugi, and D.S. Silver, *When does a satellite knot fiber?*, Hiroshima Math. J. **38** (2007), 411–423.
- [13] T. Koberda and R. Santharoubane, *Quotients of surface groups and homology of finite covers via quantum representations.*, Invent. Math. **206** (2016), no. 2, 269–292.
- [14] I. Kra, *On the Nielsen-Thurston-Bers type of some self-maps of Riemann surfaces*, Acta Math. **146** (1981), no. 3-4, 231–270.
- [15] D.D. Long and H.R. Morton, *Hyperbolic 3-manifolds and surface automorphisms*, Topology **25** (1986), no. 4, 575–583.
- [16] J. Marché and R. Santharoubane, *Asymptotics of the quantum representations of surface groups*, arXiv:1607.00664.
- [17] W. Menasco, *Closed incompressible surfaces in alternating knot and link complements*, Topology **23** (1984), 37–44.
- [18] N. Reshetikhin and V. G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), no. 3, 547–597.
- [19] J. Roberts, *Skein theory and Turaev-Viro invariants*, Topology **34** (1995), no. 4, 771–787.
- [20] R. Santharoubane, *Limits of the quantum  $SO(3)$  representations for the one-holed torus*, Journal of Knot Theory and its ramifications. **21** (2012), no. 11, 13pp.
- [21] ———, *Action of  $M(0,2n)$  on some kernel spaces coming from  $SU(2)$ -TQFT*, J. Lon. Math. Soc. **95** (2017), no. 3, 785–803.
- [22] J. R. Stallings, *Constructions of fibered knots and links*, Proc. of Symposia in Pure Mathematics **32** (1978), 55–60.
- [23] W. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces.*, Bull. Amer. Math. Soc. **19** (1988), no. 2, 417–431.
- [24] W. P. Thurston, *The geometry and topology of three-manifolds*, Princeton Univ. Math. Dept. Notes, 1979.
- [25] V. G. Turaev, *Quantum invariants of knots and 3-manifolds*, de Gruyter Studies in Mathematics, vol. 18, Walter de Gruyter & Co., Berlin, 1994.
- [26] V. G. Turaev and O. Y. Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, Topology **4** (1992), 865–902.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI, 48824, USA

*E-mail address:* [kalfagia@math.msu.edu](mailto:kalfagia@math.msu.edu)

*E-mail address:* [detcherry@math.msu.edu](mailto:detcherry@math.msu.edu)