

Review for Exam 2

- ▶ Sections 13.1-13.3. 14.1-14.6.
- ▶ 50 minutes.
- ▶ 5 problems, similar to homework problems.
- ▶ No calculators, no notes, no books, no phones.
- ▶ No green book needed.

Section 14.6

Example

- (a) Find the linear approximation $L(x, y)$ of the function $f(x, y) = \sin(2x + 3y) + 1$ at the point $(-3, 2)$.
- (b) Use the approximation above to estimate the value of $f(-2.9, 2.1)$.

Solution:

(a) $L(x, y) = f_x(-3, 2)(x + 3) + f_y(-3, 2)(y - 2) + f(-3, 2)$.

Since $f_x(x, y) = 2 \cos(2x + 3y)$ and $f_y(x, y) = 3 \cos(2x + 3y)$,

$$f_x(-3, 2) = 2 \cos(-6 + 6) = 2, \quad f_y(-3, 2) = 3 \cos(-6 + 6) = 3,$$

$$f(-3, 2) = \sin(-6 + 6) + 1 = 1.$$

the linear approximation is $L(x, y) = 2(x + 3) + 3(y - 2) + 1$.

Section 14.6

Example

- (a) Find the linear approximation $L(x, y)$ of the function $f(x, y) = \sin(2x + 3y) + 1$ at the point $(-3, 2)$.
- (b) Use the approximation above to estimate the value of $f(-2.9, 2.1)$.

Solution: Recall: $L(x, y) = 2(x + 3) + 3(y - 2) + 1$.

- (b) We use L to find the a linear approximation to $f(-2.9, 2.1)$.

We need to compute $L(-2.9, 2.1)$.

$$L(-2.9, 2.1) = 2(-2.9 + 3) + 3(2.1 - 2) + 1$$

$$L(-2.9, 2.1) = 2(0.1) + 3(0.1) + 1 \Rightarrow L(-2.9, 2.1) = 1.5. \triangleleft$$

Exact value is close to 1.479.

Section 14.5

Example

- (a) Find the gradient of $f(x, y, z) = \sqrt{x + 2yz}$.
- (b) Find the directional derivative of f at $(0, 2, 1)$ in the direction given by $\langle 0, 3, 4 \rangle$.
- (c) Find the maximum rate of change of f at the point $(0, 2, 1)$.

Solution:

(a) $\nabla f(x, y, z) = \frac{1}{2\sqrt{x + 2yz}} \langle 1, 2z, 2y \rangle$.

- (b) We evaluate the gradient above at $(0, 2, 1)$,

$$\nabla f(0, 2, 1) = \frac{1}{2\sqrt{0 + 4}} \langle 1, 2, 4 \rangle = \frac{1}{4} \langle 1, 2, 4 \rangle.$$

Section 14.5

Example

- (a) Find the gradient of $f(x, y, z) = \sqrt{x + 2yz}$.
- (b) Find the directional derivative of f at $(0, 2, 1)$ in the direction given by $\langle 0, 3, 4 \rangle$.
- (c) Find the maximum rate of change of f at the point $(0, 2, 1)$.

Solution: (b) Recall: $\nabla f(0, 2, 1) = \frac{1}{4}\langle 1, 2, 4 \rangle$.

We now need a unit vector parallel to $\langle 0, 3, 4 \rangle$,

$$\mathbf{u} = \frac{1}{\sqrt{9 + 16}}\langle 0, 3, 4 \rangle = \frac{1}{5}\langle 0, 3, 4 \rangle.$$

Then, $(D_{\mathbf{u}}f)(0, 2, 1) = \frac{1}{4}\langle 1, 2, 4 \rangle \cdot \frac{1}{5}\langle 0, 3, 4 \rangle = \frac{1}{20}(6 + 16) = \frac{11}{10}$.

We obtain, $(D_{\mathbf{u}}f)(0, 2, 1) = \frac{11}{10}$.

Section 14.5

Example

- (a) Find the gradient of $f(x, y, z) = \sqrt{x + 2yz}$.
- (b) Find the directional derivative of f at $(0, 2, 1)$ in the direction given by $\langle 0, 3, 4 \rangle$.
- (c) Find the maximum rate of change of f at the point $(0, 2, 1)$.

Solution:

(c) The maximum rate of change of f at a point is the magnitude of its gradient at that point, that is,

$$|\nabla f(0, 2, 1)| = \frac{1}{4}|\langle 1, 2, 4 \rangle| = \frac{1}{4}\sqrt{1 + 4 + 16} = \frac{\sqrt{21}}{4}.$$

The maximum rate of change of f at $(0, 2, 1)$ is

$$|\nabla f(0, 2, 1)| = \sqrt{21}/4.$$

◁

Section 14.4

Example

Find $\partial_{xy}(e^{-xy} \sin(x + yz))$. (Do not simplify your answer.)

Solution: We first compute the x -derivative,

$$\partial_x(e^{-xy} \sin(x + yz)) = -ye^{-xy} \sin(x + yz) + e^{-xy} \cos(x + yz).$$

The second derivative is

$$\partial_{xy}(e^{-xy} \sin(x + yz)) = \partial_y(-ye^{-xy} \sin(x + yz) + e^{-xy} \cos(x + yz)),$$

$$= -e^{-xy} \sin(x + yz) + xye^{-xy} \sin(x + yz) - ye^{-xy} \cos(x + yz)z$$

$$-xe^{-xy} \cos(x + yz) - e^{-xy} \sin(x + yz)z. \quad \triangleleft$$

Section 14.3

Example

Find any value of the constant a such that the function $f(x, y) = e^{-ax} \cos(y) - e^{-y} \cos(x)$ is solution of Laplace's equation $f_{xx} + f_{yy} = 0$.

Solution:

$$f_x = -ae^{-ax} \cos(y) + e^{-y} \sin(x), \quad f_{xx} = a^2 e^{-ax} \cos(y) + e^{-y} \cos(x).$$

$$f_y = -e^{-ax} \sin(y) + e^{-y} \cos(x), \quad f_{yy} = -e^{-ax} \cos(y) - e^{-y} \cos(x).$$

$$\begin{aligned} f_{xx} + f_{yy} &= [a^2 e^{-ax} \cos(y) + e^{-y} \cos(x)] \\ &\quad + [-e^{-ax} \cos(y) - e^{-y} \cos(x)], \end{aligned}$$

$$f_{xx} + f_{yy} = (a^2 - 1)e^{-ax} \cos(y).$$

Function f is solution of $f_{xx} + f_{yy} = 0$ iff $a = \pm 1$. \triangleleft

Section 14.2

Example

Compute the limit $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 \sin^2(y)}{2x^2 + 3y^2}$.

Solution:

Since $x^2 \leq 2x^2 + 3y^2$, that is, $\frac{x^2}{2x^2 + 3y^2} \leq 1$, the non-negative function $f(x, y) = \frac{x^2 \sin^2(y)}{2x^2 + 3y^2}$ satisfies the bounds

$$0 \leq f(x, y) \leq \sin^2(y).$$

Since $\lim_{y \rightarrow 0} \sin^2(y) = 0$, the Sandwich Theorem implies that

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 \sin^2(y)}{2x^2 + 3y^2} = 0. \quad \triangleleft$$

Section 13.3

Example

Reparametrize the curve $\mathbf{r}(t) = \left\langle \frac{3}{2} \sin(t^2), 2t^2, \frac{3}{2} \cos(t^2) \right\rangle$ with respect to its arc length measured from $t = 1$ in the direction of increasing t .

Solution:

We first compute the arc length function. We start with the derivative

$$\mathbf{r}'(t) = \langle 3t \cos(t^2), 4t, -3 \sin(t^2) \rangle,$$

We now need its magnitude,

$$|\mathbf{r}'(t)| = \sqrt{9t^2 \cos^2(t^2) + 16t^2 + 9 \sin^2(t^2)},$$

$$|\mathbf{r}'(t)| = \sqrt{9t^2 + 16t^2} = (\sqrt{9 + 16}) t \Rightarrow |\mathbf{r}'(t)| = 5t.$$

Section 13.3

Example

Reparametrize the curve $\mathbf{r}(t) = \left\langle \frac{3}{2} \sin(t^2), 2t^2, \frac{3}{2} \cos(t^2) \right\rangle$ with respect to its arc length measured from $t = 1$ in the direction of increasing t .

Solution: Recall: $|\mathbf{r}'(t)| = 5t$. The arc length function is

$$s(t) = \int_1^t 5\tau \, d\tau = \frac{5}{2} \left(\tau^2 \Big|_1^t \right) = \frac{5}{2} (t^2 - 1).$$

Inverting this function for t^2 , we obtain $t^2 = \frac{2}{5}s + 1$.

The reparametrization of $\mathbf{r}(t)$ is given by

$$\hat{\mathbf{r}}(s) = \left\langle \frac{3}{2} \sin\left(\frac{2}{5}s + 1\right), 2\left(\frac{2}{5}s + 1\right), \frac{3}{2} \cos\left(\frac{2}{5}s + 1\right) \right\rangle. \quad \triangleleft$$