

## Review for Exam 4

- ▶ Sections 16.1-16.6.
- ▶ 50 minutes.
- ▶ 5 to 10 problems, similar to homework problems.
- ▶ No calculators, no notes, no books, no phones.
- ▶ No green book needed.

## Review for Exam 4


- ▶ **(16.6) Surface integrals.**
- ▶ (16.5) Surface area.
- ▶ (16.4) The Green Theorem in a plane.
- ▶ (16.3) Conservative fields, potential functions.
- ▶ (16.2) Vector fields, work, circulation, flux (plane).
- ▶ (16.1) Line integrals.

## Surface integrals (16.6): Scalar fields

### Example

Integrate the function  $g(x, y, z) = x\sqrt{4 + y^2}$  over the surface cut from the parabolic cylinder  $z = 4 - y^2/4$  by the planes  $x = 0$ ,  $x = 1$  and  $z = 0$ .

### Solution:



We must compute:  $I = \iint_S g \, d\sigma$ .

Recall  $d\sigma = \frac{|\nabla f|}{|\nabla f \cdot \mathbf{k}|} dx \, dy$ , with  $\mathbf{k} \perp R$

and in this case  $f(x, y, z) = y^2 + 4z - 16$ .

$$\nabla f = \langle 0, 2y, 4 \rangle \Rightarrow |\nabla f| = \sqrt{16 + 4y^2} = 2\sqrt{4 + y^2}.$$

Since  $R = [0, 1] \times [-4, 4]$ , its normal vector is  $\mathbf{k}$  and  $|\nabla f \cdot \mathbf{k}| = 4$ .

Then,

$$\iint_S g \, d\sigma = \iint_R (x\sqrt{4 + y^2}) \frac{2\sqrt{4 + y^2}}{4} dx \, dy.$$

## Surface integrals (16.6)

### Example

Integrate the function  $g(x, y, z) = x\sqrt{4 + y^2}$  over the surface cut from the parabolic cylinder  $z = 4 - y^2/4$  by the planes  $x = 0$ ,  $x = 1$  and  $z = 0$ .

$$\text{Solution: } \iint_S g \, d\sigma = \iint_R (x\sqrt{4 + y^2}) \frac{2\sqrt{4 + y^2}}{4} dx \, dy.$$

$$\iint_S g \, d\sigma = \frac{1}{2} \iint_R x(4 + y^2) dx \, dy = \frac{1}{2} \int_{-4}^4 \int_0^1 x(4 + y^2) dx \, dy$$

$$\iint_S g \, d\sigma = \frac{1}{2} \left[ \int_{-4}^4 (4 + y^2) dy \right] \left[ \int_0^1 x dx \right] = \frac{1}{2} \left( 4y + \frac{y^3}{3} \right) \Big|_{-4}^4 \left( \frac{x^2}{2} \right) \Big|_0^1$$

$$\iint_S g \, d\sigma = \frac{1}{2} 2 \left( 4^2 + \frac{4^3}{3} \right) \frac{1}{2} = 8 \left( 1 + \frac{4}{3} \right) \Rightarrow \iint_S g \, d\sigma = \frac{56}{3}.$$

## Review for Exam 4


- ▶ (16.6) Surface integrals.
- ▶ **(16.5) Surface area.**
- ▶ (16.4) The Green Theorem in a plane.
- ▶ (16.3) Conservative fields, potential functions.
- ▶ (16.2) Vector fields, work, circulation, flux (plane).
- ▶ (16.1) Line integrals.

## Surface area (16.5)

### Example

Set up the integral for the area of the surface cut from the parabolic cylinder  $z = 4 - y^2/4$  by the planes  $x = 0$ ,  $x = 1$ ,  $z = 0$ .

### Solution:



We must compute:  $A(S) = \iint_S d\sigma$ .

Recall  $d\sigma = \frac{|\nabla f|}{|\nabla f \cdot \mathbf{k}|} dx dy$ , with  $\mathbf{k} \perp R$ .

Recall:  $f(x, y, z) = y^2 + 4z - 16$ .

$$\nabla f = \langle 0, 2y, 4 \rangle \Rightarrow |\nabla f| = \sqrt{16 + 4y^2} = 2\sqrt{4 + y^2}.$$

Since  $R = [0, 1] \times [-4, 4]$ , its normal vector is  $\mathbf{k}$  and  $|\nabla f \cdot \mathbf{k}| = 4$ .  
Then,

$$A(S) = \iint_R \frac{2\sqrt{4 + y^2}}{4} dx dy \Rightarrow A(S) = \int_0^1 \int_{-4}^4 \frac{2\sqrt{4 + y^2}}{4} dy dx. \quad \triangleleft$$

## Review for Exam 4

- ▶ (16.6) Surface integrals.
- ▶ (16.5) Surface area.
- ▶ **(16.4) The Green Theorem in a plane.**
- ▶ (16.3) Conservative fields, potential functions.
- ▶ (16.2) Vector fields, work, circulation, flux (plane).
- ▶ (16.1) Line integrals.

## The Green Theorem in a plane (16.4)

### Example

Use the Green Theorem in the plane to find the flux of  $\mathbf{F} = (x - y^2)\mathbf{i} + (x^2 + y)\mathbf{j}$  through the ellipse  $9x^2 + 4y^2 = 36$ .

Solution: Recall:  $\oint_C \mathbf{F} \cdot \mathbf{n} \, ds = \iint_S \operatorname{div} \mathbf{F} \, dx \, dy$ .

Recall:  $\operatorname{div} \mathbf{F} = \partial_x F_x + \partial_y F_y$ . Here is simpler to compute the right-hand side than the left-hand side.  $\operatorname{div} \mathbf{F} = 1 + 1 = 2$ .

Green's Theorem implies

$$\oint_C \mathbf{F} \cdot \mathbf{n} \, ds = \iint_R (2) \, dx \, dy. = 2A(R).$$

Since  $R$  is the ellipse  $x^2/4 + y^2/9 = 1$ , its area is  $A(R) = (2)(3)\pi$ . We conclude

$$\oint_C \mathbf{F} \cdot \mathbf{n} \, ds = 12\pi. \quad \triangleleft$$

## Review for Exam 4

- ▶ (16.6) Surface integrals.
- ▶ (16.5) Surface area.
- ▶ (16.4) The Green Theorem in a plane.
- ▶ **(16.3) Conservative fields, potential functions.**
- ▶ (16.2) Vector fields, work, circulation, flux (plane).
- ▶ (16.1) Line integrals.

## Conservative fields, potential functions (16.3)

### Example

Compute  $I = \int_{(0,0,0)}^{(1,-1,0)} 2x \cos(z) dx + z dy + (y - x^2 \sin(z)) dz$ .

**Solution:** The integral is specified by the path end points. That suggests that the vector field is a gradient field.

$$\mathbf{F} = \langle 2x \cos(z), z, [y - x^2 \sin(z)] \rangle = \nabla f = \langle \partial_x f, \partial_y f, \partial_z f \rangle.$$

$$\partial_x f = 2x \cos(z) \Rightarrow f = x^2 \cos(z) + g(y, z).$$

$$\partial_y f = z = \partial_y g \Rightarrow g = yz + h(z) \Rightarrow f = x^2 \cos(z) + yz + h(z).$$

$$\partial_z f = y - x^2 \sin(z) = -x^2 \sin(z) + y + h' \Rightarrow h' = 0$$

Since  $f = x^2 \cos(z) + yz + c$ , we obtain

$$I = \int_{(0,0,0)}^{(1,-1,0)} \nabla f \cdot d\mathbf{r} = f(1, -1, 0) - f(0, 0, 0) \Rightarrow I = 1. \triangleleft$$

## Review for Exam 4

- ▶ (16.6) Surface integrals.
- ▶ (16.5) Surface area.
- ▶ (16.4) The Green Theorem in a plane.
- ▶ (16.3) Conservative fields, potential functions.
- ▶ **(16.2) Vector fields, work, circulation, flux (plane).**
- ▶ (16.1) Line integrals.

## Vector fields, work, circulation, flux (plane) (16.2)

### Example

Find the flow of the velocity field  $\mathbf{F} = \langle xy, y^2, -yz \rangle$  from the point  $(0, 0, 0)$  to the point  $(1, 1, 1)$  along the curve of intersection of the cylinder  $y = x^2$  with the plane  $z = x$ .

**Solution:** The flow (also called circulation) of the field  $\mathbf{F}$  along a curve  $C$  parametrized by  $\mathbf{r}(t)$  for  $t \in [t_0, t_1]$  is given by

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{t_0}^{t_1} \mathbf{F}(t) \cdot \mathbf{r}'(t) dt.$$

We use  $t = x$  as the parameter of the curve  $\mathbf{r}$ , so we obtain

$$\mathbf{r}(t) = \langle t, t^2, t \rangle, \quad t \in [0, 1] \quad \Rightarrow \quad \mathbf{r}'(t) = \langle 1, 2t, 1 \rangle.$$

$$\mathbf{F}(t) = \langle t(t^2), (t^2)^2, -t^2(t) \rangle \quad \Rightarrow \quad \mathbf{F}(t) = \langle t^3, t^4, -t^3 \rangle.$$

## Vector fields, work, circulation, flux (plane) (16.2)

### Example

Find the flow of the velocity field  $\mathbf{F} = \langle xy, y^2, -yz \rangle$  from the point  $(0, 0, 0)$  to the point  $(1, 1, 1)$  along the curve of intersection of the cylinder  $y = x^2$  with the plane  $z = x$ .

Solution:  $\mathbf{r}'(t) = \langle 1, 2t, 1 \rangle$  for  $t \in [0, 1]$  and  $\mathbf{F}(t) = \langle t^3, t^4, -t^3 \rangle$ .

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{t_0}^{t_1} \mathbf{F}(t) \cdot \mathbf{r}'(t) dt = \int_0^1 \langle t^3, t^4, -t^3 \rangle \cdot \langle 1, 2t, 1 \rangle dt,$$

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 (t^3 + 2t^5 - t^3) dt = \int_0^1 2t^5 dt = \left. \frac{2}{6} t^6 \right|_0^1.$$

We conclude that  $\int_C \mathbf{F} \cdot d\mathbf{r} = \frac{1}{3}$ . ◁

## Review for Exam 4

- ▶ (16.6) Surface integrals.
- ▶ (16.5) Surface area.
- ▶ (16.4) The Green Theorem in a plane.
- ▶ (16.3) Conservative fields, potential functions.
- ▶ (16.2) Vector fields, work, circulation, flux (plane).
- ▶ **(16.1) Line integrals.**

## Line integrals (16.1)

### Example

Integrate the function  $f(x, y) = x^3/y$  along the plane curve  $C$  given by  $y = x^2/2$  for  $x \in [0, 2]$ , from the point  $(0, 0)$  to  $(2, 2)$ .

**Solution:** We have to compute  $I = \int_C f ds$ , by that we mean

$$\int_C f ds = \int_{t_0}^{t_1} f(x(t), y(t)) |\mathbf{r}'(t)| dt,$$

where  $\mathbf{r}(t) = \langle x(t), y(t) \rangle$  for  $t \in [t_0, t_1]$  is a parametrization of the path  $C$ . In this case the path is given by the parabola  $y = x^2/2$ , so a simple parametrization is to use  $x = t$ , that is,

$$\mathbf{r}(t) = \left\langle t, \frac{t^2}{2} \right\rangle, \quad t \in [0, 2] \quad \Rightarrow \quad \mathbf{r}'(t) = \langle 1, t \rangle.$$

## Line integrals (16.1)

### Example

Integrate the function  $f(x, y) = x^3/y$  along the plane curve  $C$  given by  $y = x^2/2$  for  $x \in [0, 2]$ , from the point  $(0, 0)$  to  $(2, 2)$ .

**Solution:**  $\mathbf{r}(t) = \left\langle t, \frac{t^2}{2} \right\rangle$  for  $t \in [0, 2]$ , and  $\mathbf{r}'(t) = \langle 1, t \rangle$ .

$$\int_C f ds = \int_{t_0}^{t_1} f(x(t), y(t)) |\mathbf{r}'(t)| dt = \int_0^2 \frac{t^3}{t^2/2} \sqrt{1+t^2} dt,$$

$$\int_C f ds = \int_0^2 2t \sqrt{1+t^2} dt, \quad u = 1+t^2, \quad du = 2t dt.$$

$$\int_C f ds = \int_1^5 u^{1/2} du = \frac{2}{3} u^{3/2} \Big|_1^5 = \frac{2}{3} (5^{3/2} - 1).$$

We conclude that  $\int_C f ds = \frac{2}{3} (5\sqrt{5} - 1)$ .

◁