

## Separable differential equations (Sect. 1.3).

- ▶ Separable ODE.
- ▶ Solutions to separable ODE.
- ▶ Explicit and implicit solutions.
- ▶ Euler homogeneous equations.

## Separable ODE.

### Definition

A *separable* differential equation on the function  $y$  has the form

$$h(y) y'(t) = g(t),$$

where  $h, g$  are given scalar functions.

### Remark:

A differential equation  $y'(t) = f(t, y(t))$  is separable iff

$$y' = \frac{g(t)}{h(y)} \quad \Leftrightarrow \quad f(t, y) = \frac{g(t)}{h(y)}.$$

### Example

$$y'(t) = \frac{t^2}{1 - y^2(t)}, \quad y'(t) + y^2(t) \cos(2t) = 0.$$

## Separable ODE.

### Example

Determine whether the differential equation below is separable,

$$y'(t) = \frac{t^2}{1 - y^2(t)}.$$

**Solution:** The differential equation is separable, since it is equivalent to

$$(1 - y^2) y' = t^2 \quad \Rightarrow \quad \begin{cases} g(t) = t^2, \\ h(y) = 1 - y^2. \end{cases}$$

◁

**Remark:** The functions  $g$  and  $h$  are not uniquely defined. Another choice here is:

$$g(t) = c t^2, \quad h(y) = c(1 - y^2), \quad c \in \mathbb{R}.$$

## Separable ODE.

### Example

Determine whether The differential equation below is separable,

$$y'(t) + y^2(t) \cos(2t) = 0$$

**Solution:** The differential equation is separable, since it is equivalent to

$$\frac{1}{y^2} y' = -\cos(2t) \quad \Rightarrow \quad \begin{cases} g(t) = -\cos(2t), \\ h(y) = \frac{1}{y^2}. \end{cases}$$

◁

**Remark:** The functions  $g$  and  $h$  are not uniquely defined. Another choice here is:

$$g(t) = \cos(2t), \quad h(y) = -\frac{1}{y^2}.$$

## Separable ODE.

**Remark:** Not every first order ODE is separable.

### Example

- ▶ The differential equation  $y'(t) = e^{y(t)} + \cos(t)$  is not separable.
- ▶ The linear differential equation  $y'(t) = -\frac{2}{t}y(t) + 4t$  is not separable.
- ▶ The linear differential equation  $y'(t) = -a(t)y(t) + b(t)$ , with  $b(t)$  non-constant, is not separable.

## Separable differential equations (Sect. 1.3).

- ▶ Separable ODE.
- ▶ **Solutions to separable ODE.**
- ▶ Explicit and implicit solutions.
- ▶ Euler homogeneous equations.

## Solutions to separable ODE.

### Theorem (Separable equations)

If the functions  $h, g$  are continuous,  $h \neq 0$ , then the separable ODE

$$h(y) y' = g(t)$$

has infinitely many solutions  $y$  satisfying the algebraic equation

$$H(y(t)) = G(t) + c,$$

where  $c \in \mathbb{R}$  is arbitrary, and where  $H$  and  $G$  are primitives (antiderivatives) of  $h$  and  $g$ , respectively; that is,

$$H'(u) = h(u), \quad G'(t) = g(t).$$

**Remark:** Given functions  $g, h$ , find their primitives  $G, H$ .

## Solutions to separable ODE.

### Example

Find all solutions  $y$  to the equation  $y'(t) = \frac{t^2}{1 - y^2(t)}$ .

**Solution:** The equation is equivalent to

$$(1 - y^2) y'(t) = t^2 \quad \Rightarrow \quad g(t) = t^2, \quad h(y) = 1 - y^2.$$

Integrate on both sides of the equation,

$$\int [1 - y^2(t)] y'(t) dt = \int t^2 dt + c.$$

The substitution  $u = y(t)$ ,  $du = y'(t) dt$ , implies that

$$\int (1 - u^2) du = \int t^2 dt + c \quad \Leftrightarrow \quad \left(u - \frac{u^3}{3}\right) = \frac{t^3}{3} + c.$$

## Solutions to separable ODE.

### Example

Find all solutions  $y$  to the equation  $y'(t) = \frac{t^2}{1 - y^2(t)}$ .

**Solution:** Recall:  $\left(u - \frac{u^3}{3}\right) = \frac{t^3}{3} + c$ .

Substitute the unknown function  $y$  back in the equation above,

$$\left(y - \frac{y^3}{3}\right) = \frac{t^3}{3} + c, \quad c \in \mathbb{R}.$$

**Remark:** Recall the notation in the Theorem:

$$g(t) = t^2 \quad \Rightarrow \quad G(t) = \frac{t^3}{3},$$

$$h(y) = 1 - y^2 \quad \Rightarrow \quad H(y) = y - \frac{y^3}{3}.$$

Hence we recover the Theorem expression:  $H(y(t)) = G(t) + c$ .

## Solutions to separable ODE.

### Remarks:

- ▶ The equation  $y(t) - \frac{y^3(t)}{3} = \frac{t^3}{3} + c$  is algebraic in  $y$ , since there is no  $y'$  in the equation.
- ▶ Every function  $y$  satisfying the algebraic equation

$$y(t) - \frac{y^3(t)}{3} = \frac{t^3}{3} + c,$$

is a solution of the differential equation above.

- ▶ We now verify the previous statement: Differentiate on both sides with respect to  $t$ , that is,

$$y'(t) - 3\left(\frac{y^2(t)}{3}\right)y'(t) = 3\frac{t^2}{3} \quad \Rightarrow \quad (1 - y^2)y' = t^2.$$

## Solutions to separable ODE.

### Example

Find all solutions  $y$  to the equation  $y'(t) + y^2(t) \cos(2t) = 0$ .

**Solution:** The differential equation is separable,

$$\frac{y'(t)}{y^2(t)} = -\cos(2t) \Rightarrow g(t) = -\cos(2t), \quad h(y) = \frac{1}{y^2}.$$

Integrate on both sides of the equation,

$$\int \frac{y'(t)}{y^2(t)} dt = - \int \cos(2t) dt + c.$$

The substitution  $u = y(t)$ ,  $du = y'(t) dt$ , implies that

$$\int \frac{du}{u^2} = - \int \cos(2t) dt + c \Leftrightarrow -\frac{1}{u} = -\frac{1}{2} \sin(2t) + c.$$

## Solutions to separable ODE.

### Example

Find all solutions  $y$  to the equation  $y'(t) + y^2(t) \cos(2t) = 0$ .

**Solution:** Recall:  $-\frac{1}{u} = -\frac{1}{2} \sin(2t) + c$ .

Substitute the unknown function  $y$  back in the equation above,

$$-\frac{1}{y(t)} = -\frac{1}{2} \sin(2t) + c, \quad c \in \mathbb{R}.$$

**Remark:** Recall the notation in the Theorem:

$$g(t) = -\cos(2t) \Rightarrow G(t) = -\frac{1}{2} \sin(2t).$$

$$h(y) = \frac{1}{y^2} \Rightarrow H(y) = -\frac{1}{y}.$$

Hence we recover the Theorem expression:  $H(y(t)) = G(t) + c$ .

## Separable differential equations (Sect. 1.3).

- ▶ Separable ODE.
- ▶ Solutions to separable ODE.
- ▶ **Explicit and implicit solutions.**
- ▶ Euler homogeneous equations.

## Explicit and implicit solutions.

### Definition

Assume the notation in the Theorem above. The solution  $y$  of a separable ODE is given in *implicit form* iff function  $y$  is given by

$$H(y(t)) = G(t) + c,$$

The solution is given in *explicit form* iff function  $H$  is invertible and

$$y(t) = H^{-1}(G(t) + c).$$

### Example

(a)  $y(t) - \frac{y^3(t)}{3} = \frac{t^3}{3} + c$  is in implicit form.

(b)  $-\frac{1}{y(t)} = -\frac{1}{2}\sin(2t) + c$  is in implicit form.

(c)  $y(t) = \frac{2}{\sin(2t) - 2c}$  is in explicit form.

## Separable differential equations (Sect. 1.3).

- ▶ Separable ODE.
- ▶ Solutions to separable ODE.
- ▶ Explicit and implicit solutions.
- ▶ **Euler homogeneous equations.**

## Euler homogeneous equations.

### Definition

The first order ODE  $y'(t) = f(t, y(t))$  is called *Euler homogeneous* iff for every numbers  $c, t, u \in \mathbb{R}$  the function  $f$  satisfies

$$f(ct, cu) = f(t, u).$$

### Remark:

- ▶ The function  $f$  is invariant under the change of scale of its arguments.
- ▶ If  $f(t, u)$  has the property above, it must depend only on  $u/t$ .
- ▶ Therefore, there exists  $F : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(t, u) = F\left(\frac{u}{t}\right)$ .
- ▶ So, a first order ODE is Euler homogeneous iff it has the form

$$y'(t) = F\left(\frac{y(t)}{t}\right).$$



## Euler homogeneous equations.

### Example

Show that the equation below is Euler homogeneous,

$$(t - y)y' - 2y + 3t + \frac{y^2}{t} = 0.$$

**Solution:** Rewrite the equation in the standard form

$$(t - y)y' = 2y - 3t - \frac{y^2}{t} \Rightarrow y' = \frac{\left(2y - 3t - \frac{y^2}{t}\right)}{(t - y)}.$$

Divide numerator and denominator by  $t$ . We get,

$$y' = \frac{\left(2y - 3t - \frac{y^2}{t}\right) \left(\frac{1}{t}\right)}{(t - y) \left(\frac{1}{t}\right)} \Rightarrow y' = \frac{2\left(\frac{y}{t}\right) - 3 - \left(\frac{y}{t}\right)^2}{\left[1 - \left(\frac{y}{t}\right)\right]}.$$

## Euler homogeneous equations.

### Example

Show that the equation below is Euler homogeneous,

$$(t - y)y' - 2y + 3t + \frac{y^2}{t} = 0.$$

**Solution:** Recall:  $y' = \frac{2\left(\frac{y}{t}\right) - 3 - \left(\frac{y}{t}\right)^2}{\left[1 - \left(\frac{y}{t}\right)\right]}.$

We conclude that the ODE is Euler homogeneous, because the right-hand side of the equation above depends only on  $y/t$ .

Indeed, in our case:

$$f(t, y) = \frac{2y - 3t - (y^2/t)}{t - y}, \quad F(x) = \frac{2x - 3 - x^2}{1 - x},$$

and  $f(t, y) = F(y/t)$ .

◁

## Euler homogeneous equations.

### Example

Determine whether the equation below is Euler homogeneous,

$$y' = \frac{t^2}{1 - y^3}.$$

### Solution:

Divide numerator and denominator by  $t^3$ , we obtain

$$y' = \frac{t^2}{(1 - y^3)} \frac{\left(\frac{1}{t^3}\right)}{\left(\frac{1}{t^3}\right)} \Rightarrow y' = \frac{\left(\frac{1}{t}\right)}{\left(\frac{1}{t^3}\right) - \left(\frac{y}{t}\right)^3}.$$

Then, the differential equation is **not Euler homogeneous**.  $\triangleleft$

## Euler homogeneous equations.

### Theorem

If the equation  $y'(t) = f(t, y(t))$  is Euler homogeneous, then the differential equation for the unknown  $v(t) = \frac{y(t)}{t}$  is separable.

**Remark:** Euler homogeneous equations can be transformed into separable equations.

**Proof:** If  $y' = f(t, y)$  is Euler homogeneous, then it can be written as  $y' = F(y/t)$  for some function  $F$ . Introducing  $v = y/t$ ,

$$y(t) = t v(t) \Rightarrow y'(t) = v(t) + t v'(t).$$

Introduce all these changes into the ODE, then

$$v + t v' = F(v) \Rightarrow v' = \frac{(F(v) - v)}{t}.$$

This last equation is separable.  $\square$

## Euler homogeneous equations.

### Example

Find all solutions  $y$  of the equation  $y' = \frac{t^2 + 3y^2}{2ty}$ .

**Solution:** The equation is Euler homogeneous, since

$$y' = \frac{t^2 + 3y^2}{2ty} \frac{\left(\frac{1}{t^2}\right)}{\left(\frac{1}{t^2}\right)} \Rightarrow y' = \frac{1 + 3\left(\frac{y}{t}\right)^2}{2\left(\frac{y}{t}\right)}.$$

Therefore, we introduce the change of unknown  $v = y/t$ , so  $y = tv$  and  $y' = v + tv'$ . Hence

$$v + tv' = \frac{1 + 3v^2}{2v} \Rightarrow tv' = \frac{1 + 3v^2}{2v} - v = \frac{1 + 3v^2 - 2v^2}{2v}$$

We obtain the **separable** equation  $v' = \frac{1}{t} \left( \frac{1 + v^2}{2v} \right)$ .

## Euler homogeneous equations.

### Example

Find all solutions  $y$  of the equation  $y' = \frac{t^2 + 3y^2}{2ty}$ .

**Solution:** Recall:  $v' = \frac{1}{t} \left( \frac{1 + v^2}{2v} \right)$ . We rewrite and integrate it,

$$\frac{2v}{1 + v^2} v' = \frac{1}{t} \Rightarrow \int \frac{2v}{1 + v^2} v' dt = \int \frac{1}{t} dt + c_0.$$

The substitution  $u = 1 + v^2(t)$  implies  $du = 2v(t) v'(t) dt$ , so

$$\int \frac{du}{u} = \int \frac{dt}{t} + c_0 \Rightarrow \ln(u) = \ln(t) + c_0 \Rightarrow u = e^{\ln(t) + c_0}.$$

But  $u = e^{\ln(t)} e^{c_0}$ , so denoting  $c_1 = e^{c_0}$ , then  $u = c_1 t$ . Hence

$$1 + v^2 = c_1 t \Rightarrow 1 + \left(\frac{y}{t}\right)^2 = c_1 t \Rightarrow y(t) = \pm t \sqrt{c_1 t - 1}.$$