

Linear Variable coefficient equations (Sect. 1.2)

- ▶ Review: Linear constant coefficient equations.
- ▶ The Initial Value Problem.
- ▶ Linear variable coefficients equations.
- ▶ The Bernoulli equation: A nonlinear equation.

Review: Linear constant coefficient equations

Definition

Given functions a, b , a *first order, linear ODE* in the unknown function y is the equation

$$y'(t) = a(t)y(t) + b(t).$$

Theorem (Constant coefficients)

Given constants $a, b \in \mathbb{R}$ with $a \neq 0$, the linear differential equation

$$y'(t) = ay(t) + b$$

has infinitely many solutions, one for each value of $c \in \mathbb{R}$, given by

$$y(t) = ce^{at} - \frac{b}{a}.$$

Review: Linear constant coefficient equations

Example


Find all functions y solution of the ODE $y' = 2y + 3$.

Solution:

We concluded that the ODE has infinitely many solutions, given by

$$y(t) = c e^{2t} - \frac{3}{2}, \quad c \in \mathbb{R}.$$

Since we did one integration, it is reasonable that the solution contains a constant of integration, $c \in \mathbb{R}$.



Verification: $y' = 2c e^{2t}$, but we know that $2c e^{2t} = 2y + 3$, therefore we conclude that y satisfies the ODE $y' = 2y + 3$. \triangleleft

Linear Variable coefficient equations (Sect. 1.2)

- ▶ Review: Linear constant coefficient equations.
- ▶ **The Initial Value Problem.**
- ▶ Linear variable coefficients equations.
- ▶ The Bernoulli equation: A nonlinear equation.

The Initial Value Problem

Definition

The *Initial Value Problem* (IVP) for a linear ODE is the following:
Given functions a, b and constants t_0, y_0 , find a function y solving

$$y' = a(t)y + b(t), \quad y(t_0) = y_0.$$

Remark: The initial condition selects one solution of the ODE.

Theorem (Constant coefficients)

Given constants a, b, t_0, y_0 , with $a \neq 0$, the initial value problem

$$y' = ay + b, \quad y(t_0) = y_0$$

has the unique solution

$$y(t) = \left(y_0 + \frac{b}{a} \right) e^{a(t-t_0)} - \frac{b}{a}.$$

The Initial Value Problem

Example

Find the solution to the initial value problem

$$y' = 2y + 3, \quad y(0) = 1.$$

Solution: Every solution of the ODE above is given by

$$y(t) = c e^{2t} - \frac{3}{2}, \quad c \in \mathbb{R}.$$

The initial condition $y(0) = 1$ selects only one solution:

$$1 = y(0) = c - \frac{3}{2} \Rightarrow c = \frac{5}{2}.$$

We conclude that $y(t) = \frac{5}{2} e^{2t} - \frac{3}{2}$.

◁

Linear Variable coefficient equations (Sect. 1.2)

- ▶ Review: Linear constant coefficient equations.
- ▶ The Initial Value Problem.
- ▶ **Linear variable coefficients equations.**
- ▶ The Bernoulli equation: A nonlinear equation.

Linear variable coefficients equations

Theorem (Variable coefficients)

Given continuous functions a, b and constants t_0, y_0 , all solutions to the equation

$$y' = a(t)y + b(t)$$

are given by

$$y(t) = c e^{A(t)} + e^{A(t)} \int e^{-A(t)} b(t) dt,$$

where we have introduced the function $A(t) = \int a(t) dt$.

Remarks:

- The function $\mu(t) = e^{-A(t)}$ is called an **integrating factor**.
- See the proof in the Lecture Notes.

Linear variable coefficients equations

Example

Find all solutions y to the equation

$$t y' = -2y + 4t^2,$$

Solution: We first express the equation as in the Theorem,

$$y' = -\frac{2}{t}y + 4t \Rightarrow y' + \frac{2}{t}y = 4.$$

$$e^{f(t)} y' + \frac{2}{t} e^{f(t)} y = 4t e^{f(t)}, \quad f'(t) = \frac{2}{t}.$$

This function $\mu = e^{f(t)}$ is an integrating factor.

$$f(t) = \int \frac{2}{t} dt = 2 \ln(t) = \ln(t^2)$$

Therefore, $\mu(t) = e^{f(t)} = t^2$.

Linear variable coefficients equations

Example

Find the solution y to the IVP

$$t y' + 2y = 4t^2, \quad y(1) = 2.$$

Solution: The integrating factor is $\mu(t) = t^2$. Hence,

$$t^2 \left(y' + \frac{2}{t} y \right) = t^2(4t) \Leftrightarrow t^2 y' + 2t y = 4t^3$$

$$(t^2 y)' = 4t^3 \Leftrightarrow t^2 y = t^4 + c \Leftrightarrow y = t^2 + \frac{c}{t^2}.$$

We conclude $y(t) = t^2 + \frac{c}{t^2}$ for any $c \in \mathbb{R}$.

◁

Linear variable coefficients equations

Theorem (Variable coefficients)

Given continuous functions $a, b : \mathbb{R} \rightarrow \mathbb{R}$ and given constants $t_0, y_0 \in \mathbb{R}$, the IVP

$$y' = a(t)y + b(t) \quad y(t_0) = y_0$$

has the unique solution

$$y(t) = e^{A(t)} \left(y_0 + \int_{t_0}^t e^{-A(s)} b(s) ds \right),$$

where we have introduced the function $A(t) = \int_{t_0}^t a(s) ds$.

Remarks:

- (a) The function $\mu(t) = e^{-A(t)}$ is called an **integrating factor**.
- (b) See the proof in the Lecture Notes.

Linear variable coefficients equations

Example

Find the solution y to the IVP

$$t y' + 2y = 4t^2, \quad y(1) = 2.$$

Solution: We already know that all solutions to the differential equation are

$$y = t^2 + \frac{c}{t^2}.$$

The initial condition implies

$$2 = y(1) = 1 + c, \quad \Rightarrow \quad c = 1.$$

We conclude that $y(t) = t^2 + \frac{1}{t^2}$.

◁

Linear Variable coefficient equations (Sect. 1.2)

- ▶ Review: Linear constant coefficient equations.
- ▶ The Initial Value Problem.
- ▶ Linear variable coefficients equations.
- ▶ **The Bernoulli equation: A nonlinear equation.**

The Bernoulli equation

Remark: The Bernoulli equation is a **non-linear** differential equation that can be transformed into a **linear** differential equation.

Definition

Given functions $p, q : \mathbb{R} \rightarrow \mathbb{R}$ and a real number n , the differential equation in the unknown function $y : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$y' + p(t)y = q(t)y^n$$

is called the *Bernoulli equation*.

Theorem

The function $y : \mathbb{R} \rightarrow \mathbb{R}$ is a solution of the Bernoulli equation for

$$y' + p(t)y = q(t)y^n, \quad n \neq 1,$$

iff the function $v = 1/y^{(n-1)}$ is solution of the linear differential equation

$$v' - (n-1)p(t)v = -(n-1)q(t).$$

The Bernoulli equation

Example

Given arbitrary constants $a \neq 0$ and b , find every solution of the differential equation

$$y' = ay + by^3.$$

Solution: This is a Bernoulli equation. Divide the equation by y^3 ,

$$\frac{y'}{y^3} = \frac{a}{y^2} + b.$$

Introduce the function $v = 1/y^2$, with derivative $v' = -2\left(\frac{y'}{y^3}\right)$, into the differential equation above,

$$-\frac{v'}{2} = av + b \Rightarrow v' = -2av - 2b \Rightarrow v' + 2av = -2b.$$

The Bernoulli equation

Example

Given arbitrary constants $a \neq 0$ and b , find every solution of the differential equation

$$y' = ay + by^3.$$

Solution: Recall: $v' + 2av = -2b$.

The last equation is a linear differential equation for v . This equation can be solved using the integrating factor method.

Multiply the equation by $\mu(t) = e^{2at}$,

$$(e^{2at}v)' = -2be^{2at} \Rightarrow e^{2at}v = -\frac{b}{a}e^{2at} + c$$

We obtain that $v = ce^{-2at} - \frac{b}{a}$. Since $v = 1/y^2$,

$$\frac{1}{y^2} = ce^{-2at} - \frac{b}{a} \Rightarrow y(t) = \pm \frac{1}{(ce^{-2at} - \frac{b}{a})^{1/2}}. \quad \triangleleft$$