

On linear and non-linear equations. (Sect. 1.6).

- ▶ Review: Linear differential equations.
- ▶ Non-linear differential equations.
- ▶ Properties of solutions to non-linear ODE.
- ▶ Direction Fields.

Review: Linear differential equations.

Theorem (Variable coefficients)

Given continuous functions $a, b : (t_1, t_2) \rightarrow \mathbb{R}$, with $t_2 > t_1$, and given constants $t_0 \in (t_1, t_2)$, $y_0 \in \mathbb{R}$, the IVP

$$y' = -a(t)y + b(t), \quad y(t_0) = y_0,$$

has the unique solution $y : (t_1, t_2) \rightarrow \mathbb{R}$ given by

$$y(t) = \frac{1}{\mu(t)} \left[y_0 + \int_{t_0}^t \mu(s) b(s) ds \right], \quad (1)$$

where the integrating factor function is given by

$$\mu(t) = e^{A(t)}, \quad A(t) = \int_{t_0}^t a(s) ds.$$

Proof: Based on the integration factor method.

Review: Linear differential equations.

Remarks:

- ▶ The Theorem above assumes that the coefficients a, b , are continuous in $(t_1, t_2) \subset \mathbb{R}$.
- ▶ The Theorem above implies:
 - (a) There is an explicit expression for the solutions of a linear IVP, given in Eq. (1).
 - (b) For every initial condition $y_0 \in \mathbb{R}$ there exists a unique solution to a linear IVP.
 - (c) For every initial condition $y_0 \in \mathbb{R}$ the corresponding solution $y(t)$ of a linear IVP is defined for all $t \in (t_1, t_2)$.
- ▶ **None of these properties holds for solutions to non-linear differential equations.**

On linear and non-linear equations. (Sect. 1.6).

- ▶ Review: Linear differential equations.
- ▶ **Non-linear differential equations.**
- ▶ Properties of solutions to non-linear ODE.
- ▶ Direction Fields.

Non-linear differential equations.

Definition

An ordinary differential equation $y'(t) = f(t, y(t))$ is called *non-linear* iff the function $(t, u) \mapsto f(t, u)$ is non-linear in the second argument.

Example

- (a) The differential equation $y'(t) = \frac{t^2}{y^3(t)}$ is non-linear, since the function $f(t, u) = t^2/u^3$ is non-linear in the second argument.
- (b) The differential equation $y'(t) = 2ty(t) + \ln(y(t))$ is non-linear, since the function $f(t, u) = 2tu + \ln(u)$ is non-linear in the second argument, due to the term $\ln(u)$.
- (c) The differential equation $\frac{y'(t)}{y(t)} = 2t^2$ is linear, since the function $f(t, u) = 2t^2u$ is linear in the second argument.

On linear and non-linear equations. (Sect. 1.6).

- ▶ Review: Linear differential equations.
- ▶ Non-linear differential equations.
- ▶ **Properties of solutions to non-linear ODE.**
- ▶ Direction Fields.

Properties of solutions to non-linear ODE.

Theorem (Non-linear ODE)

Fix a non-empty rectangle $R = (t_1, t_2) \times (u_1, u_2) \subset \mathbb{R}^2$ and fix a function $f : R \rightarrow \mathbb{R}$ denoted as $(t, u) \mapsto f(t, u)$. If the functions f and $\partial_u f$ are continuous on R , and $(t_0, y_0) \in R$, then there exists a smaller open rectangle $\hat{R} \subset R$ with $(t_0, y_0) \in \hat{R}$ such that the IVP

$$y'(t) = f(t, y(t)), \quad y(t_0) = y_0$$

has a unique solution y on the set $\hat{R} \subset \mathbb{R}^2$.

Remarks:

- (i) There is no general explicit expression for the solution $y(t)$ to a non-linear ODE.
- (ii) Non-uniqueness of solution to the IVP above may happen at points $(t, u) \in \mathbb{R}^2$ where $\partial_u f$ is not continuous.
- (iii) Changing the initial data y_0 may change the domain on the variable t where the solution $y(t)$ is defined.

Properties of solutions to non-linear ODE.

Example

Given non-zero constants a_1, a_2, a_3, a_4 , find every solution y of

$$y' = \frac{t^2}{(y^4 + a_4 y^3 + a_3 y^2 + a_2 y + a_1)}.$$

Solution: The ODE is separable. So first, rewrite the equation as

$$(y^4 + a_4 y^3 + a_3 y^2 + a_2 y + a_1) y' = t^2,$$

then we integrate in t on both sides of the equation,

$$\int (y^4 + a_4 y^3 + a_3 y^2 + a_2 y + a_1) y' dt = \int t^2 dt + c.$$

Introduce the substitution $u = y(t)$, so $du = y'(t) dt$,

$$\int (u^4 + a_4 u^3 + a_3 u^2 + a_2 u + a_1) du = \int t^2 dt + c.$$

Properties of solutions to non-linear ODE.

Example

Given non-zero constants a_1, a_2, a_3, a_4 , find every solution y of

$$y' = \frac{t^2}{(y^4 + a_4 y^3 + a_3 y^2 + a_2 y + a_1)}.$$

Solution:

Recall: $\int (u^4 + a_4 u^3 + a_3 u^2 + a_2 u + a_1) du = \int t^2 dt + c.$

Integrate, and in the result substitute back the function y :

$$\frac{1}{5} y^5(t) + \frac{a_4}{4} y^4(t) + \frac{a_3}{3} y^3(t) + \frac{a_2}{2} y^2(t) + a_1 y(t) = \frac{t^3}{3} + c.$$

The solution is in implicit form. It is the root of a polynomial degree five. There is no formula for the roots of a general polynomial degree five or bigger.

There is no explicit expression for solutions y of the ODE. \triangleleft

Properties of solutions to non-linear ODE.

Example

Find every solution y of the initial value problem

$$y'(t) = y^{1/3}(t), \quad y(0) = 0.$$

Remark: The equation above is non-linear, separable, and the function $f(t, u) = u^{1/3}$ has derivative

$$\partial_u f = \frac{1}{3} \frac{1}{u^{2/3}},$$

so $\partial_u f$ is not continuous at $u = 0$.

The initial condition above is precisely where f is not continuous.

Solution: There are two solutions to the IVP above:

The first solution is

$$y_1(t) = 0.$$

Properties of solutions to non-linear ODE.

Example

Find every solution y of the initial value problem

$$y'(t) = y^{1/3}(t), \quad y(0) = 0.$$

Solution: The second solution is obtained as follows:

$$\int [y(t)]^{-1/3} y'(t) dt = \int dt + c.$$

Then, the substitution $u = y(t)$, with $du = y'(t) dt$, implies that

$$\int u^{-1/3} du = \int dt + c \Rightarrow \frac{3}{2} [y(t)]^{2/3} = t + c,$$

$$y(t) = \left[\frac{2}{3} (t + c) \right]^{3/2} \Rightarrow 0 = y(0) = \left(\frac{2}{3} c \right)^{3/2} \Rightarrow c = 0.$$

So, the second solution is: $y_2(t) = \left(\frac{2}{3} t \right)^{3/2}$. Recall $y_1(t) = 0$. \triangleleft

Properties of solutions to non-linear ODE.

Example

Find the solution y to the initial value problem

$$y'(t) = y^2(t), \quad y(0) = y_0.$$

Solution: This is a separable equation. So,

$$\int \frac{y' dt}{y^2} = \int dt + c \Rightarrow -\frac{1}{y} = t + c \Rightarrow y(t) = -\frac{1}{t + c}.$$

Using the initial condition in the expression above,

$$y_0 = y(0) = -\frac{1}{c} \Rightarrow c = -\frac{1}{y_0} \Rightarrow y(t) = \frac{1}{\left(\frac{1}{y_0} - t \right)}.$$

This solution diverges at $t = 1/y_0$, so its domain is $\mathbb{R} - \{y_0\}$.

The solution domain depends on the values of the initial data y_0 . \triangleleft

Properties of solutions to non-linear ODE.

Summary:

- ▶ Linear ODE:
 - (a) There is an explicit expression for the solution of a linear IVP.
 - (b) For every initial condition $y_0 \in \mathbb{R}$ there exists a unique solution to a linear IVP.
 - (c) The domain of the solution of a linear IVP is defined for every initial condition $y_0 \in \mathbb{R}$.

- ▶ Non-linear ODE:
 - (i) There is no general explicit expression for the solution $y(t)$ to a non-linear ODE.
 - (ii) Non-uniqueness of solution to a non-linear IVP may happen at points $(t, u) \in \mathbb{R}^2$ where $\partial_u f$ is not continuous.
 - (iii) Changing the initial data y_0 may change the domain on the variable t where the solution $y(t)$ is defined.

On linear and non-linear equations. (Sect. 1.6).

- ▶ Review: Linear differential equations.
- ▶ Non-linear differential equations.
- ▶ Properties of solutions to non-linear ODE.
- ▶ **Direction Fields.**

Direction Fields.

Remarks:

- ▶ One does not need to solve a differential equation $y'(t) = f(t, y(t))$ to have a qualitative idea of the solution.
- ▶ Recall that $y'(t)$ represents the slope of the tangent line to the graph of function y at the point $(t, y(t))$.
- ▶ A differential equation provides these slopes, $f(t, y(t))$, for every point $(t, y(t))$.
- ▶ **Key idea:** Graph the function $f(t, y)$ on the yt -plane, not as points, but as slopes of small segments.


Definition

A *Direction Field* for the differential equation $y'(t) = f(t, y(t))$ is the graph on the yt -plane of the values $f(t, y)$ as slopes of a small segments.

Direction Fields.

Example

We know that the solution of $y' = y$ are the exponentials $y(t) = y_0 e^t$. The graph of these solution is simple. So is the direction field:



Direction Fields.


Example

The solution of $y' = \sin(y)$ is simple to compute. The equation is separable. After some calculations the implicit solution are

$$\ln \left| \frac{\csc(y_0) + \cot(y)}{\csc(y) + \cot(y_0)} \right| = t.$$

for $y_0 \in \mathbb{R}$. The graph of these solution is not simple to do.

But the direction field is simple to plot:



Direction Fields.

Example

The solution of $y' = \frac{(1+y^3)}{(1+t^2)}$ could be hard to compute. But the direction field is simple to plot:

