


## §2.7. Rates of Change.

Key points: Functions of Motion.

We consider the following physical quantities describing motion as functions of time  $t$ .

- Position or displacement  $s(t)$ . (in feet). 
- Velocity at  $t$ :  $v(t) = s'(t)$ . (in ft/s)
- Speed: magnitude of velocity, i.e.,  $|v|$ .
- Acceleration:  $a(t) = v'(t)$ . (in ft/s<sup>2</sup>)
- Average =  $\frac{s(t_2) - s(t_1)}{t_2 - t_1}$  (sec 1.4)
- Distance

Rank:  $v > 0 \iff$  moving forward  $\iff s$  is increasing  
 $v < 0 \iff$  moving backward  $\iff s$  is decreasing  
 $|v|$  is increasing  $\iff$  speed up,  $|v|$  is decreasing  $\iff$  slow down.  
 $a > 0 \iff v$  is increasing  $\begin{cases} v > 0, |v| \text{ is increasing} \\ v < 0, |v| \text{ is decreasing} \end{cases}$

eg. 1. The position of a particle moving along the  $x$ -axis is  $x(t) = t^4 - 4t^3 + 1$ ,  $t > 0$  (5/6).

(a) when is the velocity negative? (b) when is the acceleration negative?

solution:  $v(t) = (t^4 - 4t^3 + 1)' = 4t^3 - 4 \cdot 3t^2 + 0 = \boxed{4t^3 - 12t^2}$

$$v(t) < 0 \iff 4t^3 - 12t^2 < 0 \iff 4t^2(t-3) < 0$$

The velocity is negative when  $t < 3$ .

$$a(t) = v'(t) = (4t^3 - 12t^2)' = 4 \cdot 3t^2 - 12 \cdot 2t = 12t^2 - 24t$$

$$a(t) < 0 \iff 12t^2 - 24t < 0$$

$$\iff 12t(t-2) < 0$$

The acceleration is negative when  $t < 2$ .

Rank: (a) is equivalent to ask "when is the particle moving in the negative direction"

eg. 2. A ball is thrown upward from the top of a building 50 feet tall.

The height of the ball is described by the function,  $h(t) = -t^2 + 5t + 50$ .

(a) When does the ball reach the maximum height?

(b) When does the ball reach the ground with what velocity?

Solution: (a) Maximum height  $\Leftrightarrow$  velocity zero.



$$v'(t) = (-t^2 + 5t + 50)' = -2t + 5 = 0$$

$$\Rightarrow t = 2.5 \text{ (s)}$$

$$(b). h(t) = -t^2 + 5t + 50 = 0 \Rightarrow t^2 - 5t - 50 = 0 \Rightarrow (t+5)(t-10) = 0$$


$$\text{and } v(10) = -2 \cdot 10 + 5 = -15 \text{ ft/s} \Rightarrow t = 10 \text{ (s)}$$

(c). Sketch the graph of  $h$  and  $v$ .



eg. 3. (WW \* 10)

Give the graph of  $v(t)$  in  $t \in [0, 10]$  as follow.



At  $t=9$ , the particle reaches its maximum speed.

The interval the particle moves forward:  $v > 0$   $t \in (4, 8)$

The interval the particle moves backward:  $v < 0$   $t \in (0, 2) \cup (8, 10)$

The interval the particle stops:  $v = 0$   $t \in [2, 4]$

The interval the particle speed up:  $|v|$  is increasing:  $(4, 5) \cup (8, 9)$

The interval the particle slow down:  $|v|$  is decreasing:  $(0, 2) \cup (7, 8) \cup (9, 10)$

The interval the acceleration is positive:  $v$  is increasing:  $(0, 2) \cup (4, 5) \cup (9, 10)$

The interval the acceleration is negative:  $v$  is decreasing:  $(7, 9)$

The interval the acceleration is zero:  $v$  is constant:  $(2, 4) \cup (5, 7)$

## 2.6 Implicit Differentiation.

Key points: • Explicit/Implicit functions


• Implicit Differentiation Rule.

Explicit function: From equation  $4x + 2y = 6$ , we can solve for  $y$  as.

$$y = -2x + 3 \text{ EXPLICITLY.}$$

Implicit function: While from  $x^2 + y^2 = 1$ , it is not so convenient to solve for  $y$ .

Instead of solving  $y$  (as two functions), we assume an implicit relation  $y = y(x)$  (unknown function) which satisfies the above equation. Such unknown functions are called **IMPLICIT** functions.



The goal is HOW TO TAKE THE DERIVATIVES of such unknown functions.

And the tangent line to the given curve (as an equation of  $x, y$ ).

eg. Suppose  $y$  and  $x$  satisfy the implicit equation  $\frac{x^2}{4} + \frac{y^2}{5} = 1$

$$\text{Find } y' = \frac{dy}{dx}.$$

Solution: Step 1: Take derivatives (with respect to  $x$ ) both sides of the equation:

$$\left(\frac{x^2}{4} + \frac{y^2}{5}\right)' = (1)' = 0$$

$$\Leftrightarrow \left(\frac{x^2}{4}\right)' + \left(\frac{y^2}{5}\right)' = 0 \Leftrightarrow \frac{1}{4} \cdot 2x + \frac{1}{5} \cdot (y^2)' = 0. \quad (*)$$

Caution:  $(y^2)' \neq 2 \cdot y$ .  $y = y(x)$  is a FUNCTION of  $x$ .

we need to apply CHAIN RULE, with outer function  $[\ ]^2$  and inner  $y(x)$

$$\left([y(x)]^2\right)' = [2y(x)] \cdot y'(x) = 2y \cdot y'$$

$$\text{i.e. } (*) \Leftrightarrow \frac{1}{4} \cdot 2x + \frac{1}{5} \cdot 2y \cdot y' = 0 \Leftrightarrow \left[\frac{2}{5}y\right] \cdot y' = -\frac{1}{2}x$$

Step 2: Solve for  $y'$  as a function of  $x, y$ :  $y' = -\frac{5}{4} \cdot \frac{x}{y}$

eg. 2. Suppose  $x, y$  satisfy the implicit equation  $y^2 + x \cdot y + x^3 = 3$ .

(F16). Find  $y' = \frac{dy}{dx}$  as a function of  $x, y$ .

Solution: Take derivatives both sides of the equation:  $(y^2 + x \cdot y + x^3)' = 3'$

Notice  $(y^2)' = 2y \cdot y'$ , (chain rule), and  $(x \cdot y)' = x' \cdot y + x \cdot y'$

Therefore,  $= 1 \cdot y + x \cdot y'$

Caution:  $y' \neq 1$  since  $y = y(x)$  is a function of  $x$

$$(*) \quad 2y \cdot y' + y + x \cdot y' + 3x^2 = 0.$$

Then fix  $x, y$  (treat them as some numbers) and solve for  $y'$ .

$$(2y + x) \cdot y' + y + 3x^2 = 0 \Leftrightarrow (2y + x) \cdot y' = -y - 3x^2$$

$$\Leftrightarrow \boxed{y' = \frac{-y - 3x^2}{2y + x}}$$

eg. 3. Consider the curve ~~the~~  $x^2 + y^3 + x \cdot y = 1$ .

(S16). (a). Find the slope of the tangent line of the curve at the point  $(2, -1)$ .

(b) Find the equation of the tangent line.

Hint: Recall slope of the tangent line = derivative of the "function" evaluated at "this point".

Here "the function" is the implicit function  $y = y(x)$  and the point (x-coordinate) is  $x = 2$ .

ie. (a) is equivalent to find  $\boxed{\left. \frac{dy}{dx} \right|_{x=2}}$ .

(a). Take derivative both sides:  $(x^2 + y^3 + x \cdot y)' = (1)'$   $\Leftrightarrow (x^2)' + (y^3)' + (x \cdot y)' = 0$ .

$(x^2)' = 2x$ .  $(x \cdot y)' = x' \cdot y + x \cdot y' = y + x \cdot y'$  (product rule).

$(y^3)'$ : outer function:  $\square^3$ ,  $((\square)^3)' = 3 \square^2$  Plug in inner  $y(x)$   
inner function:  $y(x)$ ,  $y'(x)$

Chain rule gives us:  $(y^3)' = 3y^2 \cdot y'$ . Therefore,  $2x + 3y^2 \cdot y' + y + x \cdot y' = 0$

Plug in  $(2, -1)$ , ie,  $x = 2, y = -1$ .  $2 \cdot 2 + 3(-1)^2 \cdot y' + (-1) + 2 \cdot y' = 0$

$$\Rightarrow 4 + 3 \cdot y' - 1 + 2 \cdot y' \Rightarrow 5y' = -3 \Rightarrow \boxed{y' = -\frac{3}{5}}$$

(b) Point slope formula:  $(2, -1)$ ;  $-\frac{3}{5}$ .  
point slope

$$\boxed{y = -\frac{3}{5}(x - 2) - 1}$$