Monomial Bases for NBC Complexes

Jason I. Brown
Department of Mathematics and Statistics
Dalhousie University
Halifax, NS B3H 3J5, CANADA

Bruce E. Sagan
Department of Mathematics
Michigan State University
East Lansing, MI 48824-1027, USA
sagan@math.msu.edu
www.math.msu.edu/~sagan

1. Complexes and chromatic polynomials

2. NBC complexes

3. Stanley-Reisner rings and hsop's

4. Monomial ideals

5. Comments

1. Complexes and chromatic polynomials

Let Δ be a *simplicial complex* on a finite set E, so Δ is a family of subsets of E satisfying

$$S \in \Delta \text{ and } T \subseteq S \text{ implies } T \in \Delta.$$

The $S \in \Delta$ are called *faces*. We assume Δ is *pure of rank* r meaning that $|S| = r$ for all maximal faces $S \in \Delta$. For $0 \leq i \leq r$, let

$$f_i = f_i(\Delta) = \# \text{ of faces } S \in \Delta \text{ with } |S| = i.$$

The *f-polynomial* of Δ is

$$f(x) = f_0 + f_1 x + f_2 x^2 + \cdots + f_r x^r.$$

The *h-polynomial* of Δ is

$$\begin{aligned} h(x) &= (1-x)^r f\left(\frac{x}{1-x}\right) \\ &= f_0(1-x)^r + f_1 x(1-x)^{r-1} + \cdots + f_r x^r. \end{aligned}$$

and let

$$h_i = h_i(\Delta) = \text{coefficient of } x^i \text{ in } h(x).$$

Let $G = (V, E)$ be a graph with $|V| = p$ and $|E| = q$. A *proper coloring* of G is $c : V \to \{1, 2, \ldots, \lambda\}$ such that

$$vw \in E \quad \text{implies} \quad c(v) \neq c(w).$$

The *chromatic polynomial* of G is

$$P(G) = P(G; \lambda) = \#\text{ of such proper colorings.}$$

Example. Let

$G =$ 

$$\begin{aligned} P(G) &= \#\text{ ways to color } t, \text{ then } u, \text{ then } v, \text{ then } w \\ &= \lambda(\lambda-1)(\lambda-2)(\lambda-2) \\ &= \lambda^4 - 5\lambda^3 + 8\lambda^2 - 4\lambda. \end{aligned}$$

Proposition 1 *Let $\overline{K}_p$ be the edgeless graph and let T be a tree on p vertices. Then*

$$P(\overline{K}_p; \lambda) = \lambda^p \quad \textit{and} \quad P(T; \lambda) = \lambda(\lambda-1)^{p-1}.$$

Let G be a graph and $e \in E$. Let

$$\begin{aligned} G \setminus e &= G \text{ with } e \text{ deleted,} \\ G/e &= G \text{ with } e \text{ contracted.} \end{aligned}$$

Example. Let $G =$



$G \setminus e =$



$G/e =$



Theorem 2 (Deletion-Contraction) *For $e \in E$*

$$P(G) = P(G \setminus e) - P(G/e)$$

Proof. If $e = vw$ then

$$\begin{aligned} P(G \setminus e) &= (\#\text{ proper } c \text{ for } G \setminus e \text{ s.t. } c(v) \neq c(w)) \\ &\quad + (\#\text{ proper } c \text{ for } G \setminus e \text{ s.t. } c(v) = c(w)) \\ &= P(G) + P(G/e). \qquad \blacksquare \end{aligned}$$

Corollary 3 *For any graph G:*

1. *$P(G;\lambda)$ is a monic polynomial in λ.*
2. *$\deg P(G;\lambda) = p = |V|$.*
3. *Coefficients of $P(G;\lambda)$ alternate in sign.*

2. NBC complexes

Define coefficients f_i by

$$P(G;\lambda) = f_0\lambda^p - f_1\lambda^{p-1} + \cdots$$

and coefficients h_i by

$$P(G;\lambda) = h_0\lambda(\lambda-1)^{p-1} - h_1\lambda(\lambda-1)^{p-2} + \cdots.$$

Let

$$\mathcal{C} = \mathcal{C}(G) = \text{set of cycles/circuits of } G.$$

Let G be *ordered* meaning that E has been given a total order $e_1 < e_2 < \ldots < e_q$. Then each $C \in \mathcal{C}$ has *broken circuit*

$$\overline{C} = C - \min C.$$

The *NBC complex* of G is

$$\Delta = \Delta(G) = \{S \subseteq E \ : \ S \text{ contains no } \overline{C}\}.$$

Then $\Delta(G)$ is a pure simplicial complex.

Theorem 4 *Let $P(G;\lambda)$ have coefficients f_i and h_i as defined above. Then for $0 \leq i \leq p$*

$$f_i = f_i(\Delta(G)) \quad \textit{and} \quad h_i = h_i(\Delta(G)).$$

$G =$ 

$\mathcal{C}(G) =$ 

$$\begin{aligned}
\bar{\mathcal{C}}(G) &= \{35, 34, 245\} \\
\Delta(G) &= \{\emptyset\} \cup \{1, 2, 3, 4, 5\} \\
&\quad \cup \{12, 13, 14, 15, 23, 24, 25, 45\} \\
&\quad \cup \{123, 124, 125, 145\} \\
(f_i(\Delta)) &= (1, 5, 8, 4, 0).
\end{aligned}$$

$$P(G; \lambda) = \lambda(\lambda - 1)(\lambda - 2)^2 = \lambda^4 - 5\lambda^3 + 8\lambda^2 - 4\lambda.$$

$P\left(\right.$  $; \lambda\left.\right) = P\left(\right.$  $; \lambda\left.\right) - P\left(\right.$  $; \lambda\left.\right)$

$= P\left(\right.$ $; \lambda\left.\right) - P\left(\right.$ $; \lambda\left.\right) - P\left(\right.$ $; \lambda\left.\right) + P\left(\right.$ $; \lambda\left.\right)$

$= \lambda(\lambda - 1)^3 - \lambda(\lambda - 1)^2 - \lambda(\lambda - 1)^2 + \lambda(\lambda - 1)$

$= \lambda(\lambda - 1)^3 - 2\lambda(\lambda - 1)^2 + \lambda(\lambda - 1)$

$(h_i(\Delta)) = (1, 2, 1, 0, 0)$

3. Stanley-Reisner rings and hsop's

Let $F[\mathbf{x}]$ be the polynomial ring over field F with variables $\mathbf{x} = \{x_1, \ldots, x_q\}$. If $E = \{e_1, \ldots, e_q\}$ then $S \subseteq E$ has *monomial*

$$\mathbf{x}^S = \prod_{e_i \in S} x_i.$$

Simplicial complex Δ has *Stanley-Reisner ring*

$$F(\Delta) = F[\mathbf{x}]/(\mathbf{x}^S \ : \ S \notin \Delta).$$

In particular, for an ordered graph G we let

$$F(G) = F(\Delta(G)) = F[\mathbf{x}]/(\mathbf{x}^{\overline{C}} \ : \ C \in \mathcal{C}(G)).$$

Now $F(G)$ has a *homogeneous system of parameters (hsop) of degree one* $\theta_1, \ldots, \theta_t$, i.e.,

1. θ_i is linear without constant term for all i,
2. $\theta_1, \ldots, \theta_t$ are algebraically independent,
3. $F(G)/(\theta_1, \ldots, \theta_t)$ is finite dim. over F.

Brown gave an explicit hsop for $F(G)$. WLOG G is connected and let T be a spanning tree of G. If $e \in E(T)$ then e has *fundamental disconnecting set*

$$D_e = D_e(G) = \{f \in E(G) \ : \ T - e + f \text{ connected}\}$$

and hsop element (when $F = \mathbb{Z}_2$)

$$\theta_e = \sum_{e_i \in D_e} x_i.$$

Example

$G =$ 

$T =$ 

$$\mathcal{C}(G) = \{13467, 2345, 12567\}$$

$$\mathbb{Z}_2(G) = \mathbb{Z}_2[x_1, \ldots, x_7]/(x_3x_4x_6x_7, x_3x_4x_5, x_2x_5x_6x_7)$$



$\theta_3 = x_3 + x_1 + x_2$



$\theta_4 = x_4 + x_1 + x_2$



$\theta_5 = x_5 + x_2$



$\theta_6 = x_6 + x_1$



$\theta_7 = x_7 + x_1$

4. Monomial ideals

If $F(\Delta)$ has an hsop $\theta_1, \ldots, \theta_t$ we let

$$R(\Delta) = F(\Delta)/(\theta_1, \ldots, \theta_t).$$

Consider

$$\text{Mon}(k) = \text{set of monomials in } F[x_1, \ldots, x_k].$$

A subset $L \subseteq \text{Mon}(k)$ is a *lower order ideal* if

$$m \in L \text{ and } n|m \text{ imples } n \in L.$$

The *lower order ideal generated* by $S \subseteq \text{Mon}(k)$ is

$$L(S) = \{m \in \text{Mon}(k) \ : \ m|n \text{ for some } n \in S\}.$$

Upper ideal and $U(S)$ are defined dually.

Theorem 5 (Macaulay, Stanley) *Suppose Δ is a simplicial complex and that the ring $F(\Delta)$ is Cohen-Macaulay. Then $R(\Delta)$ has a basis, L, which is a lower order ideal of monomials and*

$$h_i(\Delta) = \# \text{ of monomials of total degree } i \text{ in } L.$$

For a graph G, $F(G)$ is Cohen-Macaulay. We have a conjectured construction of a basis for $R(G)$.

An ordering $e_1 < \ldots < e_q$ is *standard* if the last $p-1$ edges form a tree. Let $k = |E(G) - E(T)|$. We can pick the monomial basis for $R(G)$ inside Mon(k) since Brown's θ_i can be used to eliminate the other variables, replacing each $\mathbf{x}^{\overline{C}}$ by a polynomial $p_{\overline{C}}$.

Example. In our running example, $k = 2$ and

$$\mathbb{Z}_2(G) = \mathbb{Z}_2[x_1, \ldots, x_7]/(x_3x_4x_6x_7, x_3x_4x_5, x_2x_5x_6x_7).$$
$$\theta_3 = x_3 + x_1 + x_2, \ \theta_4 = x_4 + x_1 + x_2, \ \theta_5 = x_5 + x_2.$$

So, picking one of the broken circuit monomials

$$\mathbf{x}^{\overline{C}} = x_3x_4x_5 \quad \text{becomes} \quad p_{\overline{C}} = (x_1 + x_2)^2 x_2.$$

For $1 \leq i \leq k$, the graph $T + e_i$ has a unique *fundamental circuit* C_i.

Conjecture 6 *Let G be connected. Then there is a standard ordering of E such that $R(G)$ has basis*

$$L(G) = \text{Mon}(k) - U(m_{\overline{C}} \ : \ C \in \mathcal{C}(G))$$

where

$$m_{\overline{C}} = \begin{cases} x_i^{\#\overline{C}_i} & \text{if } C = C_i \text{ fundamental,} \\ \min p_{\overline{C}} & \text{else.} \end{cases}$$

Here min p *picks out the lexicographically smallest monomial in p.*





Fundamental cycles:



$C_1 = \{1, 3, 4, 6, 7\}$ $\quad m_{\overline{C}_1} = x_1^4$



$C_2 = \{2, 3, 4, 5\}$ $\quad m_{\overline{C}_2} = x_2^3$

Nonfundamental cycle:



$C_3 = \{1, 2, 5, 6, 7\}$ $\quad m_{\overline{C}_3} = x_1^2 x_2^2$

$\mathbf{x}^{\overline{C}_3} = x_2 x_5 x_6 x_7$

$p_{\overline{C}_3} = x_2 x_2 x_1 x_1$

So $R(G)$ has basis

$$L(G) = \mathrm{Mon}(2) - U(x_1^4,\ x_2^3,\ x_1^2 x_2^2).$$

5. Comments

A graph with a standard ordering satisfying the conjecture is said to have a *broken circuit basis*.

a. (Generalized) theta graphs and phi graphs have broken circuit bases.

b. By only considering the fundamental circuits:

Proposition 7 *If G has a broken circuit basis and $c_i = |C_i|$ for $1 \leq i \leq k$, then $R(G)$ is spanned by*

$$L\left(\prod_{1\leq i\leq k} x_i^{c_i-2}\right).$$

Stanley showed that the number of acyclic orientations of G is given by $P(G;-1)$. So one can use this proposition to estimate their number.

c. The results we have about broken circuit bases are proved by deletion/contraction. It is hoped that together with the ear decomposition of a block we will be able to prove the full conjecture.

d. The conjecture may even be true for representable matroids.