Topological Properties of Activity Orders for Matroids

Rieuwert J. Blok
Dipartimento di Matematica
Università degli studi di Roma "Tor Vergata"
Via della ricerca scientifica
00133 Roma, ITALY
blokr@member.ams.org

and

Bruce E. Sagan
Department of Mathematics
Michigan State University
East Lansing, MI 48824-1027, USA
sagan@math.msu.edu
www.math.msu.edu/~sagan

0. Origin of the Problem
1. Matroids and the External Order
2. The Order Complex
3. The Main Theorem
4. Applications
5. An Open Problem

1. Matroids and the External Order

Let M be a matroid on a finite set E and let

$$\mathcal{B}(M) = \{B \subseteq E \mid B \text{ is a base of } M\}.$$

Ex. 1. The *vector matroid* $M = (V, +, \cdot,)$ has $E = V$, a finite dimensional vector space over a finite field, and

$$\mathcal{B}(V) = \{B| \ B \text{ is a basis of } V\}.$$

2. The *graph matroid* $M = G$ where G is a connected graph has $E = E(G)$ (edges of G) and

$$\mathcal{B}(G) = \{E(T) \mid T \text{ is a spanning tree of } G\}.$$

Here we will make no distinction between $F \subseteq E$ and the spanning graph of G with edge set F.

The *independent sets of* M are

$$\mathcal{I}(M) = \{I \mid I \subseteq B \text{ for some base } B\}.$$

The *circuits of* M are

$$\mathcal{C}(M) = \{C \mid C \text{ a minimal dependent set}\}.$$

If $M = G$ then a circuit of M is just a cycle in G.

From now on all matroids will be *ordered*, i.e., there will be a total order on E. Then $F \subseteq E$ has *external active set*

$$\mathrm{Act}(F) = \{e \in E \mid e = \min C \text{ where } C \in \mathcal{C}(F \cup e)\}.$$

Ex. Let $G =$ 

and $F =$ 

Then $\mathrm{Act}(F) = \{1\}$.

The *external lattice of the ordered matroid* M is

$$L(M) = \mathcal{B}(M) \uplus \hat{0}$$

where $\hat{0}$ is a unique minimal element and for bases $A, B \in \mathcal{B}(M)$

$$A \leq B \quad \text{iff} \quad A \subseteq B \cup \mathrm{Act}(B).$$

Theorem 1 (Las Vergnas) *$L(M)$ is a lattice with unique maximal element T which is the lexicographically largest base (writing sets in increasing order). $L(M)$ is also ranked with rank function*

$$\rho(B) = |\mathrm{Act}(B)| + 1.$$

$G =$



$L(G) =$



2. The Order Complex

An *abstract simplicial complex* is a finite family Δ of finite nonempty sets called *simplices* such that

$$S \in \Delta \text{ and } R \subseteq S \ (R \neq \emptyset) \text{ implies } R \in \Delta.$$

Let P be any finite poset with a unique minimal element $\hat{0}$ and unique maximal element $\hat{1}$. The *order complex of* P is

$$\Delta(P) = \{C \mid C \text{ is a chain in } P \setminus \hat{0}, \hat{1}\}.$$

Note that this is an abstract simplicial complex since a subset of a chain is again a chain.

We have the correspondences

P	$\Delta(P)$
1-chain	vertex
2-chain	edge
3-chain	triangle
k-chain	simplex of dimension $k-1$

If M is an ordered matroid then $\Delta(M) = \Delta(L(M))$.

$\Delta(G) =$



3. The Main Theorem

If M is a matroid on E then the *dual of* M is the matroid M^* defined by

$$\mathcal{B}(M^*) = \{B^* \mid B^* = E - B \text{ for some } B \in \mathcal{B}(M)\}.$$

Ex. For our example graph $E(G) = \{0, 1, 2, 3, 4\}$ so

$$\begin{aligned} \mathcal{B}(M) &= \{012, 013, 014, 024, 123, 124, 034, 234\} \\ \mathcal{B}(M^*) &= \{34, 24, 23, 13, 04, 03, 12, 01\} \end{aligned}$$

If M is a matroid on E and $F \subseteq E$ then the *restriction of* M *to* F is the matroid $M|F$ defined by

$$\mathcal{B}(M|F) = \{B \cap F \mid B \in \mathcal{B}(M) \text{ and } |B \cap F| \text{ maximum}\}.$$

Ex. (cont) Let $F = T = 234 = \hat{1}$ in $L(G)$

$$\mathcal{B}(M^*|T) = \{34, 24, 23\}.$$

If M is a matroid then the *matroid complex* $[M]$ is the abstract simplicial complex whose simplices are the nonempty independent sets of M.

Ex. (cont) We have a homotopy equivalence

$$[M^*|T] = \{34, 24, 23, 2, 3, 4\} \simeq S^1 \text{ (a 1-sphere)}$$

Theorem 2 (B & S) *If* $T = \hat{1}$ *in* $L(M)$ *then*

$$\Delta(M) \simeq [M^*|T].$$

4. Applications

Denote the Möbius function of $L(M)$ by

$$\mu(M) := \mu_{L(M)}(\hat{0}, T).$$

So $\mu(M)$ is the reduced Euler characteristic of $\Delta(M)$

$$\mu(M) = \sum_{i \geq -1} (-1)^i \dim \tilde{H}_i(\Delta(M)).$$

The *rank of* M is $\mathrm{rk}(M) := |B|$ for any $B \in \mathcal{B}(M)$.

Corollary 3 (B & S) *Let* $\mathrm{rk}(M) = r > 1$.

(i) If there is $B \in \mathcal{B}(M)$ *with* $B \subseteq E - T$, *then*

$$\tilde{H}_i(M) = \{0\} \text{ for all } i \geq 0 \text{ and } \mu(M) = 0.$$

(ii) If $E - S$ *contains a base of* M *for all* $S \subset T$ *but not for* $S = T$, *then*

$$\tilde{H}_i(M) = \begin{cases} \mathbb{Z} & \text{if } i = r-2, \\ \{0\} & \text{else,} \end{cases} \quad \text{and } \mu(M) = (-1)^{r-2}.$$

Proof (i) By definition $E - B \in \mathcal{B}(M^*)$ and we're given $E - B \supseteq T$. So $\Delta(M) \simeq [M^*|T] \simeq (r-1)$-ball and we are done.

(ii) Similarly $\Delta(M) \simeq [M^*|T] \simeq (r-2)$-sphere. ■

A *star* is a complete bipartite graph of the form $K_{1,n}$.

Corollary 4 (B & S) *Consider a matroid obtained by ordering the edges of the complete graph K_r, $r > 2$.*

(i) If T is not a star then then $\mu(K_r) = 0$.

(ii) If T is a star then $\mu(K_r) = (-1)^{r-3}$.

Proof (i) If T is not a star, then $K_r - E(T)$ is connected. So $K_r - E(T)$ contains a spanning tree and thus a base for K_r. Now use the previous corollary. (ii) is similar. ■

Note that $\Delta(M)$ is not necessarily shellable: Take an ordering of $E(K_r)$ so that the $r-1$ largest edges all meet at a vertex v. So every edge of $E - T$ is active and

$$\dim \Delta(K_r) = \rho(L(K_r)) - 2 = |\operatorname{Act}(T)| - 1 = \binom{r-1}{2} - 1.$$

But from Corollary 4, $\Delta(K_r)$ has non-zero homology in dimension $r-3$.

Theorem 5 (Björner) *The complex $[M]$ is shellable for any matroid M.*

5. An Open Problem

Las Vergnas also defined an *internal order* $\leq_I$ on M in a way equivalent to defining

$$A \leq_I B \text{ iff } E - B \leq E - A \text{ in } L(M^*).$$

If $L_I(M)$ is the resulting lattice, then it is isompor-phic to the poset dual of $L(M^*)$ and so has the same homology and Möbius function.

One can combine the external and internal orders as follows. Define the *internal-external order* $\leq_{IO}$ on $\mathcal{B}(M)$ by

$$A \leq_{IO} B \text{ iff } A \leq B \text{ or } A \leq_I B.$$

It follows from results of Las Vergnas that this is a well-defined partial order.

Question 6 *What can be said about the homology and Möbius function of the corresponding poset* $L_{IO}(M)$*?*