

Complex, distinct eigenvalues (Sect. 7.8)

- ▶ Review: Classification of 2×2 diagonalizable systems.
- ▶ Review: The case of diagonalizable matrices.
- ▶ The algebraic multiplicity of an eigenvalue.
- ▶ Non-diagonalizable matrices with a repeated eigenvalue.
- ▶ Phase portraits for 2×2 systems.

Review: Classification of 2×2 diagonalizable systems.

Remark:

Diagonalizable 2×2 matrices A with real coefficients are classified according to their eigenvalues.

- (a) $\lambda_1 \neq \lambda_2$, real-valued. Hence, A has two non-proportional eigenvectors $\mathbf{v}_1, \mathbf{v}_2$ (eigen-directions), (Section 7.5).
- (b) $\lambda_1 = \bar{\lambda}_2$, complex-valued. Hence, A has two non-proportional eigenvectors $\mathbf{v}_1 = \bar{\mathbf{v}}_2$, (Section 7.6).
- (c-1) $\lambda_1 = \lambda_2$ real-valued with two non-proportional eigenvectors $\mathbf{v}_1, \mathbf{v}_2$, (Section 7.8).

Remark:

- (c-2) $\lambda_1 = \lambda_2$ real-valued with only one eigen-direction. Hence, A is not diagonalizable, (Section 7.8).

Complex, distinct eigenvalues (Sect. 7.8)

- ▶ Review: Classification of 2×2 diagonalizable systems.
- ▶ **Review: The case of diagonalizable matrices.**
- ▶ The algebraic multiplicity of an eigenvalue.
- ▶ Non-diagonalizable matrices with a repeated eigenvalue.
- ▶ Phase portraits for 2×2 systems.

Review: The case of diagonalizable matrices.

Theorem (Diagonalizable matrix)

If $n \times n$ matrix A is diagonalizable, with a linearly independent eigenvectors set $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ and corresponding eigenvalues $\{\lambda_1, \dots, \lambda_n\}$, then the general solution \mathbf{x} to the homogeneous, constant coefficients, linear system

$$\mathbf{x}'(t) = A\mathbf{x}(t)$$

is given by the expression below, where $c_1, \dots, c_n \in \mathbb{R}$,

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + \dots + c_n \mathbf{v}_n e^{\lambda_n t}.$$

Complex, distinct eigenvalues (Sect. 7.8)

- ▶ Review: Classification of 2×2 diagonalizable systems.
- ▶ Review: The case of diagonalizable matrices.
- ▶ **The algebraic multiplicity of an eigenvalue.**
- ▶ Non-diagonalizable matrices with a repeated eigenvalue.
- ▶ Phase portraits for 2×2 systems.

The algebraic multiplicity of an eigenvalue.

Definition

Let $\{\lambda_1, \dots, \lambda_k\}$ be the set of eigenvalues of an $n \times n$ matrix, where $1 \leq k \leq n$, hence the characteristic polynomial is

$$p(\lambda) = (-1)^n (\lambda - \lambda_1)^{r_1} \cdots (\lambda - \lambda_k)^{r_k}.$$

The positive integer r_i , for $i = 1, \dots, k$, is called the *algebraic multiplicity* of the eigenvalue λ_i . The eigenvalue λ_i is called *repeated* iff $r_i > 1$.

Remark:

- ▶ A matrix with repeated eigenvalues may or may not be diagonalizable.
- ▶ Equivalently: An $n \times n$ matrix with repeated eigenvalues may or may not have a linearly independent set of n eigenvectors.

The algebraic multiplicity of an eigenvalue.

Example

Show that matrix A is diagonalizable but matrix B is not, where

$$A = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 3 & 1 & 1 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

Solution: The eigenvalues of A are the solutions of

$$\begin{vmatrix} (3 - \lambda) & 0 & 1 \\ 0 & (3 - \lambda) & 2 \\ 0 & 0 & (1 - \lambda) \end{vmatrix} = -(\lambda - 3)^2 (\lambda - 1) = 0,$$

We conclude: $\lambda_1 = 3$, $r_1 = 2$, and $\lambda_2 = 1$, $r_2 = 1$.

Verify that the eigenvalues are: $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ 2 \end{bmatrix} \right\}$.

We conclude: A is diagonalizable.

The algebraic multiplicity of an eigenvalue.

Example

Show that matrix A is diagonalizable but matrix B is not, where

$$A = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 3 & 1 & 1 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

Solution: The eigenvalues of B are the solutions of

$$\begin{vmatrix} (3 - \lambda) & 1 & 1 \\ 0 & (3 - \lambda) & 2 \\ 0 & 0 & (1 - \lambda) \end{vmatrix} = -(\lambda - 3)^2 (\lambda - 1) = 0,$$

We conclude: $\lambda_1 = 3$, $r_1 = 2$, and $\lambda_2 = 1$, $r_2 = 1$.

Verify that the eigenvalues are: $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$.

We conclude: B is not diagonalizable.

The algebraic multiplicity of an eigenvalue.

Example

Find a fundamental set of solutions to

$$\mathbf{x}'(t) = A\mathbf{x}(t), \quad A = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix},$$

Solution: Since matrix A is diagonalizable, with eigen-pairs,

$$\lambda_1 = 3, \quad \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad \lambda_2 = 1, \quad \left\{ \begin{bmatrix} -1 \\ -2 \\ 2 \end{bmatrix} \right\}.$$

We conclude that a set of fundamental solutions is

$$\left\{ \mathbf{x}_1(t) = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} e^{3t}, \mathbf{x}_2(t) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} e^{3t}, \mathbf{x}_3(t) = \begin{bmatrix} -1 \\ -2 \\ 2 \end{bmatrix} e^t \right\}. \quad \triangleleft$$

Complex, distinct eigenvalues (Sect. 7.8)

- ▶ Review: Classification of 2×2 diagonalizable systems.
- ▶ Review: The case of diagonalizable matrices.
- ▶ The algebraic multiplicity of an eigenvalue.
- ▶ **Non-diagonalizable matrices with a repeated eigenvalue.**
- ▶ Phase portraits for 2×2 systems.

Non-diagonalizable matrices with a repeated eigenvalue.

Theorem (Repeated eigenvalue)

If λ is an eigenvalue of an $n \times n$ matrix A having algebraic multiplicity $r = 2$ and only one associated eigen-direction, then the differential equation

$$\mathbf{x}'(t) = A\mathbf{x}(t),$$

has a linearly independent set of solutions given by

$$\{\mathbf{x}^{(1)}(t) = \mathbf{v} e^{\lambda t}, \quad \mathbf{x}^{(2)}(t) = (\mathbf{v} t + \mathbf{w}) e^{\lambda t}\}.$$

where the vector \mathbf{w} is solution of

$$(A - \lambda I)\mathbf{w} = \mathbf{v}$$

which always has a solution \mathbf{w} .

Non-diagonalizable matrices with a repeated eigenvalue.

Recall: The case of a single second order equation

$$y'' + a_1 y' + a_0 y = 0$$

with characteristic polynomial

$$p(r) = r^2 + a_1 r + a_0 = (r - r_1)^2.$$

In this case a fundamental set of solutions is

$$\{y_1(t) = e^{r_1 t}, \quad y_2(t) = t e^{r_1 t}\}.$$

This is not the case with systems of first order linear equations,

$$\{\mathbf{x}^{(1)}(t) = \mathbf{v} e^{\lambda t}, \quad \mathbf{x}^{(2)}(t) = (\mathbf{v} t + \mathbf{w}) e^{\lambda t}\}.$$

In general, $\mathbf{w} \neq \mathbf{0}$.

Non-diagonalizable matrices with a repeated eigenvalue.

Example

Find fundamental solutions of $\mathbf{x}' = A\mathbf{x}$, with $A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}$.

Solution: Find the eigenvalues of A . Its characteristic polynomial is

$$p(\lambda) = \begin{vmatrix} \left(-\frac{3}{2} - \lambda\right) & 1 \\ -\frac{1}{4} & \left(-\frac{1}{2} - \lambda\right) \end{vmatrix} = \left(\lambda + \frac{3}{2}\right)\left(\lambda + \frac{1}{2}\right) + \frac{1}{4}.$$

So $p(\lambda) = \lambda^2 + 2\lambda + 1 = (\lambda + 1)^2$. The roots and multiplicity are

$$\lambda = -1, \quad r = 2.$$

The corresponding eigenvectors are the solutions of $(A + I)\mathbf{v} = \mathbf{0}$,

$$\begin{bmatrix} \left(-\frac{3}{2} + 1\right) & 1 \\ -\frac{1}{4} & \left(-\frac{1}{2} + 1\right) \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} & 1 \\ -\frac{1}{4} & \frac{1}{2} \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 \\ 1 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix}$$

Non-diagonalizable matrices with a repeated eigenvalue.

Example

Find fundamental solutions of $\mathbf{x}' = A\mathbf{x}$, with $A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}$.

Solution: Recall: $\lambda = -1$, with $r = 2$, and $(A + I) \rightarrow \begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix}$.

The eigenvector components satisfy: $v_1 = 2v_2$. We obtain,

$$\lambda = -1, \quad \mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} v_2.$$

We conclude that this eigenvalue has only one eigen-direction.

Matrix A is not diagonalizable.

Theorem above says we need to find \mathbf{w} solution of $(A + I)\mathbf{w} = \mathbf{v}$.

$$\left[\begin{array}{cc|c} -\frac{1}{2} & 1 & 2 \\ -\frac{1}{4} & \frac{1}{2} & 1 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & -2 & -4 \\ 1 & -2 & -4 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & -2 & -4 \\ 0 & 0 & 0 \end{array} \right]$$

Non-diagonalizable matrices with a repeated eigenvalue.

Example

Find fundamental solutions of $\mathbf{x}' = A\mathbf{x}$, with $A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}$.

Solution: Recall that:

$$\lambda = -1, \quad \mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} v_2, \quad \text{and} \quad (A + I)\mathbf{w} = \mathbf{v} \Rightarrow \begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} -4 \\ 0 \end{bmatrix}.$$

We obtain $w_1 = 2w_2 - 4$. That is, $\mathbf{w} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} w_2 + \begin{bmatrix} -4 \\ 0 \end{bmatrix}$.

Given a solution \mathbf{w} , then $c\mathbf{v} + \mathbf{w}$ is also a solution, $c \in \mathbb{R}$.

We choose the simplest solution, $\mathbf{w} = \begin{bmatrix} -4 \\ 0 \end{bmatrix}$. We conclude,

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{-t}, \quad \mathbf{x}^{(2)}(t) = \left(\begin{bmatrix} 2 \\ 1 \end{bmatrix} t + \begin{bmatrix} -4 \\ 0 \end{bmatrix} \right) e^{-t}. \quad \triangleleft$$

Non-diagonalizable matrices with a repeated eigenvalue.

Example

Find the solution \mathbf{x} to the IVP

$$\mathbf{x}' = A\mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}.$$

Solution: The general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{-t} + c_2 \left(\begin{bmatrix} 2 \\ 1 \end{bmatrix} t + \begin{bmatrix} -4 \\ 0 \end{bmatrix} \right) e^{-t}.$$

The initial condition is $\mathbf{x}(0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = c_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} -4 \\ 0 \end{bmatrix}$.

$$\begin{bmatrix} 2 & -4 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 0 & 4 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1/4 \end{bmatrix}.$$

We conclude: $\mathbf{x}(t) = \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{-t} + \frac{1}{4} \left(\begin{bmatrix} 2 \\ 1 \end{bmatrix} t + \begin{bmatrix} -4 \\ 0 \end{bmatrix} \right) e^{-t}. \quad \triangleleft$

Complex, distinct eigenvalues (Sect. 7.8)

- ▶ Review: Classification of 2×2 diagonalizable systems.
- ▶ Review: The case of diagonalizable matrices.
- ▶ The algebraic multiplicity of an eigenvalue.
- ▶ Non-diagonalizable matrices with a repeated eigenvalue.
- ▶ **Phase portraits for 2×2 systems.**

Phase portraits for 2×2 systems.

Example

Sketch a phase portrait for solutions of


$$\mathbf{x}' = A\mathbf{x}, \quad A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}.$$

Solution:

We start plotting the vectors

$$\mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix},$$

$$\mathbf{w} = \begin{bmatrix} -4 \\ 0 \end{bmatrix}.$$



Phase portraits for 2×2 systems.

Example

Sketch a phase portrait for solutions of


$$\mathbf{x}' = A\mathbf{x}, \quad A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}.$$

Solution:

Now plot the solutions

$$\mathbf{x}^{(1)} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{-t}$$

$$\mathbf{x}^{(2)} = \left(\begin{bmatrix} 2 \\ 1 \end{bmatrix} t + \begin{bmatrix} -4 \\ 0 \end{bmatrix} \right) e^{-t}.$$



Phase portraits for 2×2 systems.

Example

Sketch a phase portrait for solutions of

$$\mathbf{x}' = A\mathbf{x}, \quad A = \frac{1}{4} \begin{bmatrix} -6 & 4 \\ -1 & -2 \end{bmatrix}.$$


Solution:

Now plot the solutions

$$\mathbf{x}^{(1)}, \quad -\mathbf{x}^{(1)},$$

$$\mathbf{x}^{(2)}, \quad -\mathbf{x}^{(2)},$$

This is the case $\lambda < 0$.



Phase portraits for 2×2 systems.

Example

Given any vectors \mathbf{v} and \mathbf{w} , and any constant λ , plot the phase portraits of the functions


$$\mathbf{x}^{(1)}(t) = \mathbf{v} e^{\lambda t}, \quad \mathbf{x}^{(2)}(t) = (\mathbf{v} t + \mathbf{w}) e^{\lambda t},$$

Solution:

The case $\lambda < 0$. We plot the functions

$$\mathbf{x}^{(1)}, \quad -\mathbf{x}^{(1)},$$

$$\mathbf{x}^{(2)}, \quad -\mathbf{x}^{(2)}.$$



Phase portraits for 2×2 systems.

Example

Given any vectors \mathbf{v} and \mathbf{w} , and any constant λ , plot the phase portraits of the functions

$$\mathbf{x}^{(1)}(t) = \mathbf{v} e^{\lambda t}, \quad \mathbf{x}^{(2)}(t) = (\mathbf{v} t + \mathbf{w}) e^{\lambda t},$$

Solution:

The case $\lambda > 0$. We plot the functions

$$\mathbf{x}^{(1)}, \quad -\mathbf{x}^{(1)},$$

$$\mathbf{x}^{(2)}, \quad -\mathbf{x}^{(2)}.$$

