

## Examples of the Fourier Theorem (Sect. 6.2).

- ▶ The Fourier Theorem: Continuous case.
- ▶ Example: Using the Fourier Theorem.
- ▶ The Fourier Theorem: Piecewise continuous case.
- ▶ Example: Using the Fourier Theorem.

## The Fourier Theorem: Continuous case.

### Theorem (Fourier Series)

If the function  $f : [-L, L] \subset \mathbb{R} \rightarrow \mathbb{R}$  is continuous, then  $f$  can be expressed as an infinite series

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[ a_n \cos\left(\frac{n\pi x}{L}\right) + b_n \sin\left(\frac{n\pi x}{L}\right) \right] \quad (1)$$

with the constants  $a_n$  and  $b_n$  given by

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx, \quad n \geq 0,$$
$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx, \quad n \geq 1.$$

Furthermore, the Fourier series in Eq. (1) provides a  $2L$ -periodic extension of function  $f$  from the domain  $[-L, L] \subset \mathbb{R}$  to  $\mathbb{R}$ .

## The Fourier Theorem: Continuous case.

Sketch of the Proof:

- ▶ Define the partial sum functions

$$f_N(x) = \frac{a_0}{2} + \sum_{n=1}^N \left[ a_n \cos\left(\frac{n\pi x}{L}\right) + b_n \sin\left(\frac{n\pi x}{L}\right) \right]$$

with  $a_n$  and  $b_n$  given by

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx, \quad n \geq 0,$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx, \quad n \geq 1.$$

- ▶ Express  $f_N$  as a convolution of Sine, Cosine, functions and the original function  $f$ .
- ▶ Use the convolution properties to show that

$$\lim_{N \rightarrow \infty} f_N(x) = f(x), \quad x \in [-L, L].$$

□

## Examples of the Fourier Theorem (Sect. 6.2).

- ▶ The Fourier Theorem: Continuous case.
- ▶ **Example: Using the Fourier Theorem.**
- ▶ The Fourier Theorem: Piecewise continuous case.
- ▶ Example: Using the Fourier Theorem.

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1 + x & x \in [-1, 0), \\ 1 - x & x \in [0, 1]. \end{cases}$$

**Solution:** In this case  $L = 1$ . The Fourier series expansion is

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos(n\pi x) + b_n \sin(n\pi x)],$$

where the  $a_n$ ,  $b_n$  are given in the Theorem. We start with  $a_0$ ,

$$a_0 = \int_{-1}^1 f(x) dx = \int_{-1}^0 (1 + x) dx + \int_0^1 (1 - x) dx.$$

$$a_0 = \left(x + \frac{x^2}{2}\right) \Big|_{-1}^0 + \left(x - \frac{x^2}{2}\right) \Big|_0^1 = \left(1 - \frac{1}{2}\right) + \left(1 - \frac{1}{2}\right)$$

We obtain:  $a_0 = 1$ .

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1 + x & x \in [-1, 0), \\ 1 - x & x \in [0, 1]. \end{cases}$$

**Solution:** Recall:  $a_0 = 1$ . Similarly, the rest of the  $a_n$  are given by,

$$a_n = \int_{-1}^1 f(x) \cos(n\pi x) dx$$
$$a_n = \int_{-1}^0 (1 + x) \cos(n\pi x) dx + \int_0^1 (1 - x) \cos(n\pi x) dx.$$

Recall the integrals  $\int \cos(n\pi x) dx = \frac{1}{n\pi} \sin(n\pi x)$ , and

$$\int x \cos(n\pi x) dx = \frac{x}{n\pi} \sin(n\pi x) + \frac{1}{n^2\pi^2} \cos(n\pi x).$$

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1+x & x \in [-1, 0), \\ 1-x & x \in [0, 1]. \end{cases}$$

**Solution:** It is not difficult to see that

$$a_n = \frac{1}{n\pi} \sin(n\pi x) \Big|_{-1}^0 + \left[ \frac{x}{n\pi} \sin(n\pi x) + \frac{1}{n^2\pi^2} \cos(n\pi x) \right] \Big|_{-1}^0 \\ + \frac{1}{n\pi} \sin(n\pi x) \Big|_0^1 - \left[ \frac{x}{n\pi} \sin(n\pi x) + \frac{1}{n^2\pi^2} \cos(n\pi x) \right] \Big|_0^1$$

$$a_n = \left[ \frac{1}{n^2\pi^2} - \frac{1}{n^2\pi^2} \cos(-n\pi) \right] - \left[ \frac{1}{n^2\pi^2} \cos(n\pi) - \frac{1}{n^2\pi^2} \right].$$

We then conclude that  $a_n = \frac{2}{n^2\pi^2} [1 - \cos(n\pi)]$ .

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1+x & x \in [-1, 0), \\ 1-x & x \in [0, 1]. \end{cases}$$

**Solution:** Recall:  $a_0 = 1$ , and  $a_n = \frac{2}{n^2\pi^2} [1 - \cos(n\pi)]$ .

Finally, we must find the coefficients  $b_n$ .

A similar calculation shows that  $b_n = 0$ .

Then, the Fourier series of  $f$  is given by

$$f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi^2} [1 - \cos(n\pi)] \cos(n\pi x).$$

◁

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1+x & x \in [-1, 0), \\ 1-x & x \in [0, 1]. \end{cases}$$

Solution: Recall:  $f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi^2} [1 - \cos(n\pi)] \cos(n\pi x).$

We can obtain a simpler expression for the Fourier coefficients  $a_n$ .

Recall the relations  $\cos(n\pi) = (-1)^n$ , then

$$f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi^2} [1 - (-1)^n] \cos(n\pi x).$$

$$f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi^2} [1 + (-1)^{n+1}] \cos(n\pi x).$$

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1+x & x \in [-1, 0), \\ 1-x & x \in [0, 1]. \end{cases}$$

Solution: Recall:  $f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi^2} [1 + (-1)^{n+1}] \cos(n\pi x).$

If  $n = 2k$ , so  $n$  is even, so  $n + 1 = 2k + 1$  is odd, then

$$a_{2k} = \frac{2}{(2k)^2\pi^2} (1 - 1) \Rightarrow a_{2k} = 0.$$

If  $n = 2k - 1$ , so  $n$  is odd, so  $n + 1 = 2k$  is even, then

$$a_{2k-1} = \frac{2}{(2k-1)^2\pi^2} (1 + 1) \Rightarrow a_{2k-1} = \frac{4}{(2k-1)^2\pi^2}.$$

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series expansion of the function

$$f(x) = \begin{cases} 1 + x & x \in [-1, 0), \\ 1 - x & x \in [0, 1]. \end{cases}$$

Solution:

Recall:  $f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi^2} [1 + (-1)^{n+1}] \cos(n\pi x)$ , and

$$a_{2k} = 0, \quad a_{2k-1} = \frac{4}{(2k-1)^2\pi^2}.$$

We conclude:  $f(x) = \frac{1}{2} + \sum_{k=1}^{\infty} \frac{4}{(2k-1)^2\pi^2} \cos((2k-1)\pi x)$ .  $\triangleleft$

## Examples of the Fourier Theorem (Sect. 6.2).

- ▶ The Fourier Theorem: Continuous case.
- ▶ Example: Using the Fourier Theorem.
- ▶ **The Fourier Theorem: Piecewise continuous case.**
- ▶ Example: Using the Fourier Theorem.

## The Fourier Theorem: Piecewise continuous case.

Recall:

### Definition

A function  $f : [a, b] \rightarrow \mathbb{R}$  is called *piecewise continuous* iff holds,

- (a)  $[a, b]$  can be partitioned in a finite number of sub-intervals such that  $f$  is continuous on the interior of these sub-intervals.
- (b)  $f$  has finite limits at the endpoints of all sub-intervals.

## The Fourier Theorem: Piecewise continuous case.

### Theorem (Fourier Series)

If  $f : [-L, L] \subset \mathbb{R} \rightarrow \mathbb{R}$  is *piecewise continuous*, then the function

$$f_F(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[ a_n \cos\left(\frac{n\pi x}{L}\right) + b_n \sin\left(\frac{n\pi x}{L}\right) \right]$$

where  $a_n$  and  $b_n$  given by

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx, \quad n \geq 0,$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx, \quad n \geq 1.$$

satisfies that:

(a)  $f_F(x) = f(x)$  for all  $x$  where  $f$  is continuous;

(b)  $f_F(x_0) = \frac{1}{2} \left[ \lim_{x \rightarrow x_0^+} f(x) + \lim_{x \rightarrow x_0^-} f(x) \right]$  for all  $x_0$  where  $f$  is discontinuous.

## Examples of the Fourier Theorem (Sect. 6.2).

- ▶ The Fourier Theorem: Continuous case.
- ▶ Example: Using the Fourier Theorem.
- ▶ The Fourier Theorem: Piecewise continuous case.
- ▶ **Example: Using the Fourier Theorem.**

### Example: Using the Fourier Theorem.

#### Example

Find the Fourier series of  $f(x) = \begin{cases} -1 & x \in [-1, 0), \\ 1 & x \in [0, 1). \end{cases}$   
and periodic with period  $T = 2$ .

**Solution:** We start computing the Fourier coefficients  $b_n$ ;

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx, \quad L = 1,$$

$$b_n = \int_{-1}^0 (-1) \sin(n\pi x) dx + \int_0^1 (1) \sin(n\pi x) dx,$$

$$b_n = \frac{(-1)}{n\pi} \left[ -\cos(n\pi x) \Big|_{-1}^0 \right] + \frac{1}{n\pi} \left[ -\cos(n\pi x) \Big|_0^1 \right],$$

$$b_n = \frac{(-1)}{n\pi} [-1 + \cos(-n\pi)] + \frac{1}{n\pi} [-\cos(n\pi) + 1].$$



## Example: Using the Fourier Theorem.

### Example

Find the Fourier series of  $f(x) = \begin{cases} -1 & x \in [-1, 0), \\ 1 & x \in [0, 1). \end{cases}$

and periodic with period  $T = 2$ .

Solution:  $b_n = \frac{(-1)}{n\pi} [-1 + \cos(-n\pi)] + \frac{1}{n\pi} [-\cos(n\pi) + 1]$ .

$$b_n = \frac{1}{n\pi} [1 - \cos(-n\pi) - \cos(n\pi) + 1] = \frac{2}{n\pi} [1 - \cos(n\pi)],$$

We obtain:  $b_n = \frac{2}{n\pi} [1 - (-1)^n]$ .

If  $n = 2k$ , then  $b_{2k} = \frac{2}{2k\pi} [1 - (-1)^{2k}]$ , hence  $b_{2k} = 0$ .

If  $n = 2k - 1$ , then  $b_{2k-1} = \frac{2}{(2k-1)\pi} [1 - (-1)^{2k-1}]$ ,

hence  $b_{2k} = \frac{4}{(2k-1)\pi}$ .

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series of  $f(x) = \begin{cases} -1 & x \in [-1, 0), \\ 1 & x \in [0, 1). \end{cases}$

and periodic with period  $T = 2$ .

Solution: Recall:  $b_{2k} = 0$ , and  $b_{2k} = \frac{4}{(2k-1)\pi}$ .

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx, \quad L = 1,$$

$$a_n = \int_{-1}^0 (-1) \cos(n\pi x) dx + \int_0^1 (1) \cos(n\pi x) dx,$$

$$a_n = \frac{(-1)}{n\pi} \left[ \sin(n\pi x) \Big|_{-1}^0 \right] + \frac{1}{n\pi} \left[ \sin(n\pi x) \Big|_0^1 \right],$$

$$a_n = \frac{(-1)}{n\pi} [0 - \sin(-n\pi)] + \frac{1}{n\pi} [\sin(n\pi) - 0] \Rightarrow a_n = 0.$$

## Example: Using the Fourier Theorem.

### Example

Find the Fourier series of  $f(x) = \begin{cases} -1 & x \in [-1, 0), \\ 1 & x \in [0, 1). \end{cases}$   
and periodic with period  $T = 2$ .

**Solution:** Recall:  $b_{2k} = 0$ ,  $b_{2k} = \frac{4}{(2k-1)\pi}$ , and  $a_n = 0$ .  
Therefore, we conclude that

$$f_F(x) = \frac{4}{\pi} \sum_{k=1}^{\infty} \frac{1}{(2k-1)} \sin((2k-1)\pi x).$$

◁