

## REVIEW MIDTERM EXAM 1

**Section Objective(s):**


- Exam Settings.
- A Sample of Review Exercises.

**Exam Main Settings:**

- Midterm Exam 1 Covers: **Chapter 1, 2, and 3.**
- Part A: Graded by Webwork. (100 points.)
  - 4 Problems,
  - 12 Grading Submissions.
  - 50 minutes.
  - In Recitation.
  - At Recitation Time.
- Part B: Graded by TA. (100 points.)
  - 4 Problems,
  - 90 minutes.
  - In N130 Business College Complex.
  - At 8:15-9:45 pm.
- Integration table provided
- There is not talking during the exam.
- No notes, no textbooks, no calculators, no phones, no electronic devices (other than the lab computer) are allowed during the exam.
- Phones and any other electronic devices must be in bags, not in pockets.

**Academic Dishonesty**

- Anyone who violates these instructions will have committed an act of academic dishonesty. Penalties for academic dishonesty can be very severe. All cases of academic dishonesty will be reported to the Dean of Undergraduate Studies and added to the student's academic record.

**Remark on Exam Starting and End Times:**

# REVIEW FOR MIDTERM EXAM 1

## Chapter 1

- (1) Consider the population of worms in a composting pile. Assume the worm population increases by 20% each week and that a farmer takes 10 worms from the pile each week.
- Write a differential equation of the form  $P' = F(P)$ , which models this situation, where  $P$  is the number of worms as a function of time.
  - Assume that the initial worm population is 40 worms. Solve the ordinary differential equation in part (a) above with this given initial condition.
  - Find the time  $t_1$  when the farmer runs out of worms.

- (2) A population of fish has a **growth rate proportional to the amount of fish present** at that time, with a proportionality factor of  $\frac{1}{5}$  per unit time.
- Write a differential equation of the form  $P' = F(P)$ , which models this situation, where  $P$  is the number of fish as a function of time.
  - Now, assume that we have the same fish population, reproducing as above, but we are harvesting fish at a **constant rate of 100** fish per unit time. Write the differential equation in this case.
  - Assume that the initial fish population is 600 fish. Solve the ordinary differential equation in part (b) above with this given initial condition.

- (3) Use the Picard iteration to find the first 4 of a sequence  $\{y_n\}$  of approximate solutions to the IVP

$$y'(t) = 8t^3y(t), \quad y(0) = 4.$$

- (4) Find the general solution to the following ODE

$$y' - 8y^2 \cos(t) - 7y^2 \sin(4t) = 0.$$

- (5) A radioactive material has a **decay rate** proportional to the amount of radioactive material present at that time, with a proportionality factor of 2 per unit time.
- Write a differential equation of the form  $P' = F(P)$ , which models this situation, where  $P$  is the amount of radioactive material (measured in micrograms) as a function of time.
  - Now, assume that we have the same radioactive material decaying as above, but we are **adding** additional material (of the same type) **at a constant rate of 6** micrograms per unit time. Write the differential equation in this case.
  - Solve the ordinary differential equation in part (b) above, assuming the initial amount of radioactive material is 70 micrograms.

- (6) Find the solution to the following initial value problem

$$y' + 8y^3 \cos(7t) = 0, \quad y(0) = 2.$$

(7) Find the solution to the following IVP

$$ty' = 2y - 3t^3 \cos(4t), \quad y(\pi/8) = 0.$$

(8) Consider the differential equation

$$\frac{dy}{dt} = y(y^2 - 4)(y^2 + 9).$$

- (a) Find the **equilibrium solutions** of the ODE.
- (b) Construct a phase diagram and determine the stability of the critical points.
- (c) Make rough sketches of typical solution curves.

(9) Find the solution to the following IVP

$$y' = \tan(t)y - 5t, \quad t \in [0, \frac{\pi}{2}), \quad y(0) = 3.$$

(10) Find an explicit expression for the solution  $y$  of the following initial value problem.

$$y' = \frac{3y^3 + t^3}{ty^2}, \quad y(1) = 2, \quad t \geq 1.$$

(11) A glass of cold soda is placed into a room held at 30 C.

- (a) If  $k$  is a (positive cooling constant), find the differential equation satisfied by the temperature,  $T(t)$  of the soda.
- (b) Find the soda temperature as a function of time (and  $k$ ), if the initial temperature of the soda was 2 C.
- (c) If after 40 minutes the soda temperature was 10 C, find the cooling constant  $k$ .

## Review for Chapter 2

- (1) An object of mass 2 gr is hanging at the bottom of a spring with a spring constant 3 gr/sq.sec. Let  $y(t)$  denote the vertical coordinate, positive downwards and  $y = 0$  be the resting position. Find the mechanical energy of the system. If the initial position of the object is  $y(0) = -3$  and its initial velocity is  $y'(0) = 3$ , find the maximum value of the position of the object, achieved during this motion.

- (2) Find the general solution of

$$y'' - 8y' + 16y = 0.$$

- (3) Solve the initial value problem

$$y'' - 5y' + 4y = 0, \quad y(0) = -5, \quad y'(0) = 3.$$

- (4) Solve the initial value problem

$$y'' - 8y' + 32y = 0, \quad y(0) = -2, \quad y'(0) = -4.$$

- (5) Solve the initial value problem

$$y'' - 8y' + 15y = 4e^t, \quad y(0) = 5, \quad y'(0) = 1.$$

- (6) Find the general solution of

$$y'' - 6y' + 8y = 3e^{2t}.$$

- (7) Find the general solution of

$$y'' - 6y' + 9y = 4e^{3t}.$$

- (8) Find the general solution of

$$y'' - 10y' + 24y = -3\sin(2t).$$

## Review for Chapter 3

(1) Use the Laplace transform to solve the initial value problem

$$y'' + 6y' + 10y = 0, \quad y(0) = -5, \quad y'(0) = -4.$$

(2) Solve the initial value problem

$$y'' - 8y' + 16y = 5\delta(t - 3), \quad y(0) = 0, \quad y'(0) = 0.$$

(3) Consider the following second order IVP with an arbitrary force term,  $g(t)$

$$y'' - 4y' + 20y = g(t), \quad y(0) = 0, \quad y'(0) = 0.$$

Let  $G(s) = \mathcal{L}[g]$  and  $Y(s) = \mathcal{L}[y]$ . Find  $H(s)$ , such that  $Y(s) = H(s)G(s)$  and  $h(t)$  such that  $y(t) = h \star g(t)$ .

(4) Find the Laplace transform of the function

$$f(t) = \begin{cases} 0, & t < 3 \\ t^2 - 6t + 7, & t \geq 3. \end{cases}$$

(5) Solve the initial value problem

$$y'' - 7y' + 12y = 5u(t - 3)e^{-3t}, \quad y(0) = 0, \quad y'(0) = 0.$$

(6) Solve the initial value problem

$$y'' - 5y' + 4y = -5u(t - 9), \quad y(0) = 0, \quad y'(0) = 0.$$

(7) Solve the initial value problem

$$y'' - 7y' + 6y = e^{5t}\delta(t - 5), \quad y(0) = 0, \quad y'(0) = 0.$$