

## The Lagrange Inversion Formula (LIF)

Following Wilf we consider the following functional equation

$$(1) \quad z = x\phi(z)$$

Can we solve for  $z$  as an explicit function of  $x$ ? Can we find a closed formula for the sequence of coefficients,  $[x^n]z(x)$ ? *Note:* The functional equation (1) implies  $z(0) = 0$ .

**Theorem 1. The Lagrange Inversion Formula** Suppose that  $W(z)$  and  $\phi(z)$  are formal power series in  $z$  with  $\phi(0) = 1$ . Then there is a unique formal power series  $z = z(x) = \sum_n z_n x^n$ , satisfying (1). In addition, the value of  $W(z(x))$  when expanded in a power series in  $x$  about  $x = 0$  satisfies

$$(2) \quad n[x^n]W(z(x)) = [z^{n-1}]\{W'(z)\phi^n(z)\}$$

The simplest version of the theorem occurs when we take  $W(z) = z$ . In that case, (2) reduces to

$$(3) \quad n[x^n]z(x) = [z^{n-1}]\phi^n(z)$$

At first glance, it may look as if we are trading one problem, coefficient extraction on  $W(z(x))$ , for another perhaps more difficult task, coefficient extraction on the more complicated expression  $W'(z)\phi^n(z)$ . However, in practice this is not the case. In fact, we will see that LIF can still be quite useful even when an explicit solution (1) is known.

Our first example is a familiar one. In Math 481 we saw that the Catalan numbers  $c_n := |C([n])|$  counted the number of legal strings of  $n$  pairs of matching parentheses. For example,  $c_3 = 5$  since

$$(4) \quad C([3]) = \{()()(), (())(), ()(()), ((())), (()())\}$$

are the only legal strings with 3 pairs of matching parentheses. If we define  $c_0 = 1$  then the first 10 Catalan numbers are

$$(5) \quad 1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, \dots$$

Now let  $C(x) = \sum_n c_n x^n$  be the ordinary power series generating function of the Catalan numbers. We showed that  $C(x)$  satisfied the functional equation

$$(6) \quad C(x) = 1 + xC^2(x)$$

This yielded explicit closed form

$$(7) \quad C(x) = \frac{1 - \sqrt{1 - 4x}}{2x}$$

It takes quite a bit more effort to conclude that

$$(8) \quad c_n = [x^n]C(x) = \frac{1}{n+1} \binom{2n}{n}$$

We now illustrate how to derive (8) using LIF.

**Example 2.** Let  $C(x)$  be as given above. Now let  $z = C(x) - 1$  and  $\phi(z) = (1 + z)^2$ . Then  $\phi(0) = 1$  and (6) becomes

$$z = C(x) - 1 = xC^2(x) = x\phi(z)$$

Now by (3), we have

$$\begin{aligned} [x^n]z(x) &= \frac{1}{n}[z^{n-1}](1+z)^{2n} \\ &= \frac{1}{n}[z^{n-1}] \sum_k \binom{2n}{k} z^k \\ &= \frac{1}{n} \binom{2n}{n-1} \\ &= \frac{1}{n+1} \binom{2n}{n} \end{aligned}$$

and we have recovered an explicit formula for the Catalan numbers with a lot less work. Notice also that in this example, we have an explicit solution to  $z = x\phi(z)$ , namely

$$z(x) = C(x) - 1 = \frac{1 - \sqrt{1 - 4x}}{2x} - 1$$

The next example illustrates how to use LIF to reprove Binomial Inversion. *Warning:* This is used to illustrate another aspect of LIF only. It is certainly not the preferred proof in this case.

**Example 3.** Suppose that

$$f_n = \sum_{k=0}^n \binom{n}{k} g_k$$

and let  $f(x) = \sum_n f_n x^n$  and  $g(x) = \sum_n g_n x^n$ . We leave it as an exercise to show that

$$(9) \quad f(x) = \frac{1}{1-x} g\left(\frac{x}{1-x}\right)$$

Now let  $y = x/(1-x)$ . Then  $x = y/(1+y)$  and it's easy to see that

$$(10) \quad g(y) = \frac{1}{1+y} f\left(\frac{y}{1+y}\right)$$

It is now a simple matter to mimic the proof of Theorem 2 (Stirling Inversion) to quickly conclude that

$$g_n = \sum_k f_k \binom{n}{k} (-1)^{n-k}$$

Instead we focus on (9) and rewrite the substitution that preceded it as  $x = y(1 - x)$ . In other words,

$$(11) \quad x(y) = y(1 - x) = y\phi(x)$$

where  $\phi(x) := 1 - x$ . Now (10) becomes

$$\begin{aligned} g(y) &= (1 - x(y))f(x(y)) \\ &= f(x(y)) - x(y)f(x(y)) \\ &= \sum_k f_k x^k(y) - x(y) \sum_k f_k x^k(y) \\ &= \sum_k f_k x^k(y) - \sum_k f_k x^{k+1}(y) \end{aligned}$$

so that

$$(12) \quad g_n = [y^n]g(y) = \sum_k f_k [y^n]x^k(y) - \sum_k f_k [y^n]x^{k+1}(y)$$

Evidently we need to compute  $[y^n]x^k(y)$  and  $[y^n]x^{k+1}(y)$ . So now we use (2) with  $W(z) = z^k$  and  $W(z) = z^{k+1}$ , respectively. Thus

$$\begin{aligned} [y^n]x^k(y) &= \frac{1}{n}[x^{n-1}]kx^{k-1}(1-x)^n \\ &= \frac{k}{n}[x^{k-n}] \sum_j \binom{n}{j} (-1)^j x^j \\ &= \frac{k}{n} \binom{n}{n-k} (-1)^{n-k} = \frac{k}{n} \binom{n}{k} (-1)^{n-k} \end{aligned}$$

We leave it as an exercise to show that

$$(13) \quad [y^n]x^{k+1}(y) = \frac{k+1}{n} \binom{n}{k+1} (-1)^{n-k-1}$$

Returning to (12), we have

$$\begin{aligned} g_n &= [y^n]g(y) = \sum_k f_k [y^n]x^k(y) - \sum_k f_k [y^n]x^{k+1}(y) \\ &= \sum_k f_k \frac{k}{n} \binom{n}{k} (-1)^{n-k} - \sum_k f_k \frac{k+1}{n} \binom{n}{k+1} (-1)^{n-k-1} \\ &= \sum_k f_k \left( \frac{k}{n} \binom{n}{k} + \frac{k+1}{n} \binom{n}{k+1} \right) (-1)^{n-k} \\ &= \sum_k f_k \binom{n}{k} (-1)^{n-k} \end{aligned}$$

as expected. Here the last line follows from [Exercise 4](#) from 02/28.