

Worksheet (5)-Solutions

(4)

$$f(x, y, z) = xyz$$


$$C: \vec{r}(t) = \cos(t)\hat{i} + \sin(t)\hat{j} + 3t\hat{k}, \quad 0 \leq t \leq 4\pi.$$

$$\vec{v}(t) = \langle -\sin t, \cos t, 3 \rangle; \quad |\vec{v}(t)| = \sqrt{\sin^2 t + \cos^2 t + 9} = \sqrt{10}$$

$$\begin{aligned} \int_C f ds &= \int_0^{4\pi} \cos(t) \sin(t) \cdot 3t \cdot \sqrt{10} dt \\ &= 3\sqrt{10} \int_0^{4\pi} t \sin(t) \cos(t) dt \\ &= 3\sqrt{10} \int_0^{4\pi} t \cdot \frac{1}{2} \sin(2t) dt \\ &= \frac{3\sqrt{10}}{2} \int_0^{4\pi} t \cdot (\cos(2t))' dt \quad \left(\frac{-1}{2}\right) \\ &= \frac{-3\sqrt{10}}{4} \left(t \cos(2t) \Big|_0^{4\pi} - \int_0^{4\pi} \cos(2t) dt \right) \\ &= \frac{-3\sqrt{10}}{4} \left(4\pi - 0 - \frac{1}{2} \sin(2t) \Big|_0^{4\pi} \right) = \boxed{-3\sqrt{10}\pi} \end{aligned}$$

(1)

$$\int_C xy^4 ds, \quad C: \text{right half of circle } x^2 + y^2 = 16$$




$$\begin{aligned} x &= 4\cos t \\ y &= 4\sin t \\ -\frac{\pi}{2} &\leq t \leq \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} \vec{r}(t) &= \langle 4\cos t, 4\sin t \rangle \\ \vec{r}'(t) &= \langle -4\sin t, 4\cos t \rangle \\ |\vec{r}'(t)| &= \sqrt{16\sin^2 t + 16\cos^2 t} = \sqrt{16} = 4. \end{aligned}$$

$$\int_C xy^4 ds = \int_{-\pi/2}^{\pi/2} 4\cos t (4\sin t)^4 \cdot 4 dt$$

$$\begin{aligned} &= 4^6 \left(\frac{\sin^5 t}{5} \right) \Big|_{-\pi/2}^{\pi/2} = 4^6 \cdot \left(\frac{1}{5} - \left(-\frac{1}{5} \right) \right) = \frac{2 \cdot 4^6}{5} \\ &= \boxed{\frac{8192}{5}} \end{aligned}$$

(2) $\int_C (2+x^2y) ds$, C: upper half of unit circle $x^2+y^2=1$.



$$\vec{r}(t) = \langle \cos t, \sin t \rangle \quad 0 \leq t \leq \pi$$

$$|\vec{v}(t)| = 1$$

$$\begin{aligned} \int_C (2+x^2y) ds &= \int_0^\pi (2 + \cos^2 t \sin t) dt \\ &= 2t \Big|_0^\pi - \frac{\cos^3 t}{3} \Big|_0^\pi \end{aligned}$$

$$= 2\pi - \left(\frac{(-1)^3}{3} - \frac{1}{3} \right) = \boxed{2\pi + \frac{2}{3}}$$

(3) $\int_C \frac{x^2}{y^{4/3}} ds$; C: $\vec{r}(t) = \langle t^2, t^3 \rangle$, $-3 \leq t \leq 1$.

$$\vec{v}(t) = \langle 2t, 3t^2 \rangle; \quad |\vec{v}(t)| = \sqrt{4t^2 + 9t^4} = \sqrt{t^2(4+9t^2)} = \underline{\underline{|t| \sqrt{4+9t^2}}}$$

$$\int_C \frac{x^2}{y^{4/3}} ds = \int_{-3}^1 \left(\frac{(t^2)^2}{(t^3)^{4/3}} \right) |t| \sqrt{4+9t^2} dt$$

$$= \int_{-3}^0 (-t) \sqrt{4+9t^2} dt + \int_0^1 t \sqrt{4+9t^2} dt$$

$$= -\frac{1}{9} \cdot \frac{1}{2} \cdot \frac{2}{3} (4+9t^2)^{3/2} \Big|_{-3}^0 + \frac{1}{9} \cdot \frac{1}{2} \cdot \frac{2}{3} (4+9t^2)^{3/2} \Big|_0^1$$

$$= -\frac{1}{27} (4^{3/2} - 85^{3/2}) + \frac{1}{27} (13^{3/2} - 4^{3/2})$$

$$= \frac{1}{27} (85\sqrt{85} - 8 + 13\sqrt{13} - 8)$$

$$= \boxed{\frac{1}{27} (85\sqrt{85} + 13\sqrt{13} - 16)}$$