

The Schur Complement and Symmetric Positive Semidefinite (and Definite) Matrices

Jean Gallier

December 10, 2010

1 Schur Complements

In this note, we provide some details and proofs of some results from Appendix A.5 (especially Section A.5.5) of *Convex Optimization* by Boyd and Vandenberghe [1].

Let M be an $n \times n$ matrix written as a 2×2 block matrix

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

where A is a $p \times p$ matrix and D is a $q \times q$ matrix, with $n = p + q$ (so, B is a $p \times q$ matrix and C is a $q \times p$ matrix). We can try to solve the linear system

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix},$$

that is

$$\begin{aligned} Ax + By &= c \\ Cx + Dy &= d, \end{aligned}$$

by mimicking Gaussian elimination, that is, assuming that D is invertible, we first solve for y getting

$$y = D^{-1}(d - Cx)$$

and after substituting this expression for y in the first equation, we get

$$Ax + B(D^{-1}(d - Cx)) = c,$$

that is,

$$(A - BD^{-1}C)x = c - BD^{-1}d.$$

If the matrix $A - BD^{-1}C$ is invertible, then we obtain the solution to our system

$$\begin{aligned}x &= (A - BD^{-1}C)^{-1}(c - BD^{-1}d) \\y &= D^{-1}(d - C(A - BD^{-1}C)^{-1}(c - BD^{-1}d)).\end{aligned}$$

The matrix, $A - BD^{-1}C$, is called the *Schur Complement* of D in M . If A is invertible, then by eliminating x first using the first equation we find that the Schur complement of A in M is $D - CA^{-1}B$ (this corresponds to the Schur complement defined in Boyd and Vandenberghe [1] when $C = B^\top$).

The above equations written as

$$\begin{aligned}x &= (A - BD^{-1}C)^{-1}c - (A - BD^{-1}C)^{-1}BD^{-1}d \\y &= -D^{-1}C(A - BD^{-1}C)^{-1}c + (D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1})d\end{aligned}$$

yield a formula for the inverse of M in terms of the Schur complement of D in M , namely

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1} \end{pmatrix}.$$

A moment of reflexion reveals that

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} (A - BD^{-1}C)^{-1} & 0 \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} \end{pmatrix} \begin{pmatrix} I & -BD^{-1} \\ 0 & I \end{pmatrix},$$

and then

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} I & 0 \\ -D^{-1}C & I \end{pmatrix} \begin{pmatrix} (A - BD^{-1}C)^{-1} & 0 \\ 0 & D^{-1} \end{pmatrix} \begin{pmatrix} I & -BD^{-1} \\ 0 & I \end{pmatrix}.$$

It follows immediately that

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} I & BD^{-1} \\ 0 & I \end{pmatrix} \begin{pmatrix} A - BD^{-1}C & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} I & 0 \\ D^{-1}C & I \end{pmatrix}.$$

The above expression can be checked directly and has the advantage of only requiring the invertibility of D .

Remark: If A is invertible, then we can use the Schur complement, $D - CA^{-1}B$, of A to obtain the following factorization of M :

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} I & 0 \\ CA^{-1} & I \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & D - CA^{-1}B \end{pmatrix} \begin{pmatrix} I & A^{-1}B \\ 0 & I \end{pmatrix}.$$

If $D - CA^{-1}B$ is invertible, we can invert all three matrices above and we get another formula for the inverse of M in terms of $(D - CA^{-1}B)$, namely,

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1}CA^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{pmatrix}.$$

If A, D and both Schur complements $A - BD^{-1}C$ and $D - CA^{-1}B$ are all invertible, by comparing the two expressions for M^{-1} , we get the (non-obvious) formula

$$(A - BD^{-1}C)^{-1} = A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1}CA^{-1}.$$

Using this formula, we obtain another expression for the inverse of M involving the Schur complements of A and D (see Horn and Johnson [5]):

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} (A - BD^{-1}C)^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{pmatrix}.$$

If we set $D = I$ and change B to $-B$ we get

$$(A + BC)^{-1} = A^{-1} - A^{-1}B(I - CA^{-1}B)^{-1}CA^{-1},$$

a formula known as the *matrix inversion lemma* (see Boyd and Vandenberghe [1], Appendix C.4, especially C.4.3).

2 A Characterization of Symmetric Positive Definite Matrices Using Schur Complements

Now, if we assume that M is symmetric, so that A, D are symmetric and $C = B^\top$, then we see that M is expressed as

$$M = \begin{pmatrix} A & B \\ B^\top & D \end{pmatrix} = \begin{pmatrix} I & BD^{-1} \\ 0 & I \end{pmatrix} \begin{pmatrix} A - BD^{-1}B^\top & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} I & BD^{-1} \\ 0 & I \end{pmatrix}^\top,$$

which shows that M is similar to a block-diagonal matrix (obviously, the Schur complement, $A - BD^{-1}B^\top$, is symmetric). As a consequence, we have the following version of ‘‘Schur’s trick’’ to check whether $M \succ 0$ for a symmetric matrix, M , where we use the usual notation, $M \succ 0$ to say that M is positive definite and the notation $M \succeq 0$ to say that M is positive semidefinite.

Proposition 2.1 *For any symmetric matrix, M , of the form*

$$M = \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix},$$

if C is invertible then the following properties hold:

- (1) $M \succ 0$ iff $C \succ 0$ and $A - BC^{-1}B^\top \succ 0$.
- (2) If $C \succ 0$, then $M \succeq 0$ iff $A - BC^{-1}B^\top \succeq 0$.

Proof. (1) Observe that

$$\begin{pmatrix} I & BD^{-1} \\ 0 & I \end{pmatrix}^{-1} = \begin{pmatrix} I & -BD^{-1} \\ 0 & I \end{pmatrix}$$

and we know that for any symmetric matrix, T , and any invertible matrix, N , the matrix T is positive definite ($T \succ 0$) iff NTN^\top (which is obviously symmetric) is positive definite ($NTN^\top \succ 0$). But, a block diagonal matrix is positive definite iff each diagonal block is positive definite, which concludes the proof.

(2) This is because for any symmetric matrix, T , and any invertible matrix, N , we have $T \succeq 0$ iff $NTN^\top \succeq 0$. \square

Another version of Proposition 2.1 using the Schur complement of A instead of the Schur complement of C also holds. The proof uses the factorization of M using the Schur complement of A (see Section 1).

Proposition 2.2 *For any symmetric matrix, M , of the form*

$$M = \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix},$$

if A is invertible then the following properties hold:

(1) $M \succ 0$ iff $A \succ 0$ and $C - B^\top A^{-1}B \succ 0$.

(2) If $A \succ 0$, then $M \succeq 0$ iff $C - B^\top A^{-1}B \succeq 0$.

When C is singular (or A is singular), it is still possible to characterize when a symmetric matrix, M , as above is positive semidefinite but this requires using a version of the Schur complement involving the pseudo-inverse of C , namely $A - BC^\dagger B^\top$ (or the Schur complement, $C - B^\top A^\dagger B$, of A). But first, we need to figure out when a quadratic function of the form $\frac{1}{2}x^\top Px + x^\top b$ has a minimum and what this optimum value is, where P is a symmetric matrix. This corresponds to the (generally nonconvex) quadratic optimization problem

$$\text{minimize } f(x) = \frac{1}{2}x^\top Px + x^\top b,$$

which has no solution unless P and b satisfy certain conditions.

3 Pseudo-Inverses

We will need pseudo-inverses so let's review this notion quickly as well as the notion of SVD which provides a convenient way to compute pseudo-inverses. We only consider the case of square matrices since this is all we need. For comprehensive treatments of SVD and pseudo-inverses see Gallier [3] (Chapters 12, 13), Strang [7], Demmel [2], Trefethen and Bau [8], Golub and Van Loan [4] and Horn and Johnson [5, 6].

Recall that every square $n \times n$ matrix, M , has a *singular value decomposition*, for short, *SVD*, namely, we can write

$$M = U\Sigma V^\top,$$

where U and V are orthogonal matrices and Σ is a diagonal matrix of the form

$$\Sigma = \text{diag}(\sigma_1, \dots, \sigma_r, 0, \dots, 0),$$

where $\sigma_1 \geq \dots \geq \sigma_r > 0$ and r is the rank of M . The σ_i 's are called the *singular values* of M and they are the positive square roots of the nonzero eigenvalues of MM^\top and $M^\top M$. Furthermore, the columns of V are eigenvectors of $M^\top M$ and the columns of U are eigenvectors of MM^\top . Observe that U and V are not unique.

If $M = U\Sigma V^\top$ is some SVD of M , we define the *pseudo-inverse*, M^\dagger , of M by

$$M^\dagger = V\Sigma^\dagger U^\top,$$

where

$$\Sigma^\dagger = \text{diag}(\sigma_1^{-1}, \dots, \sigma_r^{-1}, 0, \dots, 0).$$

Clearly, when M has rank $r = n$, that is, when M is invertible, $M^\dagger = M^{-1}$, so M^\dagger is a “generalized inverse” of M . Even though the definition of M^\dagger seems to depend on U and V , actually, M^\dagger is uniquely defined in terms of M (the same M^\dagger is obtained for all possible SVD decompositions of M). It is easy to check that

$$\begin{aligned} MM^\dagger M &= M \\ M^\dagger MM^\dagger &= M^\dagger \end{aligned}$$

and both MM^\dagger and $M^\dagger M$ are symmetric matrices. In fact,

$$MM^\dagger = U\Sigma V^\top V\Sigma^\dagger U^\top = U\Sigma\Sigma^\dagger U^\top = U \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} U^\top$$

and

$$M^\dagger M = V\Sigma^\dagger U^\top U\Sigma V^\top = V\Sigma^\dagger \Sigma V^\top = V \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} V^\top.$$

We immediately get

$$\begin{aligned} (MM^\dagger)^2 &= MM^\dagger \\ (M^\dagger M)^2 &= M^\dagger M, \end{aligned}$$

so both MM^\dagger and $M^\dagger M$ are orthogonal projections (since they are both symmetric). We claim that MM^\dagger is the orthogonal projection onto the range of M and $M^\dagger M$ is the orthogonal projection onto $\text{Ker}(M)^\perp$, the orthogonal complement of $\text{Ker}(M)$.

Obviously, $\text{range}(MM^\dagger) \subseteq \text{range}(M)$ and for any $y = Mx \in \text{range}(M)$, as $MM^\dagger M = M$, we have

$$MM^\dagger y = MM^\dagger Mx = Mx = y,$$

so the image of MM^\dagger is indeed the range of M . It is also clear that $\text{Ker}(M) \subseteq \text{Ker}(M^\dagger M)$ and since $MM^\dagger M = M$, we also have $\text{Ker}(M^\dagger M) \subseteq \text{Ker}(M)$ and so,

$$\text{Ker}(M^\dagger M) = \text{Ker}(M).$$

Since $M^\dagger M$ is Hermitian, $\text{range}(M^\dagger M) = \text{Ker}(M^\dagger M)^\perp = \text{Ker}(M)^\perp$, as claimed.

It will also be useful to see that $\text{range}(M) = \text{range}(MM^\dagger)$ consists of all vector $y \in \mathbb{R}^n$ such that

$$U^\top y = \begin{pmatrix} z \\ 0 \end{pmatrix},$$

with $z \in \mathbb{R}^r$.

Indeed, if $y = Mx$, then

$$U^\top y = U^\top Mx = U^\top U \Sigma V^\top x = \Sigma V^\top x = \begin{pmatrix} \Sigma_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} V^\top x = \begin{pmatrix} z \\ 0 \end{pmatrix},$$

where Σ_r is the $r \times r$ diagonal matrix $\text{diag}(\sigma_1, \dots, \sigma_r)$. Conversely, if $U^\top y = \begin{pmatrix} z \\ 0 \end{pmatrix}$, then $y = U \begin{pmatrix} z \\ 0 \end{pmatrix}$ and

$$\begin{aligned} MM^\dagger y &= U \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} U^\top y \\ &= U \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} U^\top U \begin{pmatrix} z \\ 0 \end{pmatrix} \\ &= U \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} \begin{pmatrix} z \\ 0 \end{pmatrix} \\ &= U \begin{pmatrix} z \\ 0 \end{pmatrix} = y, \end{aligned}$$

which shows that y belongs to the range of M .

Similarly, we claim that $\text{range}(M^\dagger M) = \text{Ker}(M)^\perp$ consists of all vector $y \in \mathbb{R}^n$ such that

$$V^\top y = \begin{pmatrix} z \\ 0 \end{pmatrix},$$

with $z \in \mathbb{R}^r$.

If $y = M^\dagger M u$, then

$$y = M^\dagger M u = V \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} V^\top u = V \begin{pmatrix} z \\ 0 \end{pmatrix},$$

for some $z \in \mathbb{R}^r$. Conversely, if $V^\top y = \begin{pmatrix} z \\ 0 \end{pmatrix}$, then $y = V \begin{pmatrix} z \\ 0 \end{pmatrix}$ and so,

$$\begin{aligned} M^\dagger M V \begin{pmatrix} z \\ 0 \end{pmatrix} &= V \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} V^\top V \begin{pmatrix} z \\ 0 \end{pmatrix} \\ &= V \begin{pmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{pmatrix} \begin{pmatrix} z \\ 0 \end{pmatrix} \\ &= V \begin{pmatrix} z \\ 0 \end{pmatrix} = y, \end{aligned}$$

which shows that $y \in \text{range}(M^\dagger M)$.

If M is a symmetric matrix, then in general, there is no SVD, $U\Sigma V^\top$, of M with $U = V$. However, if $M \succeq 0$, then the eigenvalues of M are nonnegative and so the nonzero eigenvalues of M are equal to the singular values of M and SVD's of M are of the form

$$M = U\Sigma U^\top.$$

Analogous results hold for complex matrices but in this case, U and V are unitary matrices and MM^\dagger and $M^\dagger M$ are Hermitian orthogonal projections.

If M is a normal matrix which, means that $MM^\top = M^\top M$, then there is an intimate relationship between SVD's of M and block diagonalizations of M . As a consequence, the pseudo-inverse of a normal matrix, M , can be obtained directly from a block diagonalization of M .

If M is a (real) normal matrix, then it can be block diagonalized with respect to an orthogonal matrix, U , as

$$M = U\Lambda U^\top,$$

where Λ is the (real) block diagonal matrix,

$$\Lambda = \text{diag}(B_1, \dots, B_n),$$

consisting either of 2×2 blocks of the form

$$B_j = \begin{pmatrix} \lambda_j & -\mu_j \\ \mu_j & \lambda_j \end{pmatrix}$$

with $\mu_j \neq 0$, or of one-dimensional blocks, $B_k = (\lambda_k)$. Assume that B_1, \dots, B_p are 2×2 blocks and that $\lambda_{2p+1}, \dots, \lambda_n$ are the scalar entries. We know that the numbers $\lambda_j \pm i\mu_j$, and the λ_{2p+k} are the eigenvalues of A . Let $\rho_{2j-1} = \rho_{2j} = \sqrt{\lambda_j^2 + \mu_j^2}$ for $j = 1, \dots, p$, $\rho_{2p+j} = \lambda_j$ for $j = 1, \dots, n - 2p$, and assume that the blocks are ordered so that $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$. Then, it is easy to see that

$$UU^\top = U^\top U = U\Lambda U^\top U\Lambda^\top U^\top = U\Lambda\Lambda^\top U^\top,$$

with

$$\Lambda\Lambda^\top = \text{diag}(\rho_1^2, \dots, \rho_n^2)$$

so, the singular values, $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n$, of A , which are the nonnegative square roots of the eigenvalues of AA^\top , are such that

$$\sigma_j = \rho_j, \quad 1 \leq j \leq n.$$

We can define the diagonal matrices

$$\Sigma = \text{diag}(\sigma_1, \dots, \sigma_r, 0, \dots, 0)$$

where $r = \text{rank}(A)$, $\sigma_1 \geq \dots \geq \sigma_r > 0$, and

$$\Theta = \text{diag}(\sigma_1^{-1}B_1, \dots, \sigma_{2p}^{-1}B_p, 1, \dots, 1),$$

so that Θ is an orthogonal matrix and

$$\Lambda = \Theta\Sigma = (B_1, \dots, B_p, \lambda_{2p+1}, \dots, \lambda_r, 0, \dots, 0).$$

But then, we can write

$$A = U\Lambda U^\top = U\Theta\Sigma U^\top$$

and we if let $V = U\Theta$, as U is orthogonal and Θ is also orthogonal, V is also orthogonal and $A = V\Sigma U^\top$ is an SVD for A . Now, we get

$$A^+ = U\Sigma^+V^\top = U\Sigma^+\Theta^\top U^\top.$$

However, since Θ is an orthogonal matrix, $\Theta^\top = \Theta^{-1}$ and a simple calculation shows that

$$\Sigma^+\Theta^\top = \Sigma^+\Theta^{-1} = \Lambda^+,$$

which yields the formula

$$A^+ = U\Lambda^+U^\top.$$

Also observe that if we write

$$\Lambda_r = (B_1, \dots, B_p, \lambda_{2p+1}, \dots, \lambda_r),$$

then Λ_r is invertible and

$$\Lambda^+ = \begin{pmatrix} \Lambda_r^{-1} & 0 \\ 0 & 0 \end{pmatrix}.$$

Therefore, the pseudo-inverse of a normal matrix can be computed directly from any block diagonalization of A , as claimed.

Next, we will use pseudo-inverses to generalize the result of Section 2 to symmetric matrices $M = \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix}$ where C (or A) is singular.

4 A Characterization of Symmetric Positive Semidefinite Matrices Using Schur Complements

We begin with the following simple fact:

Proposition 4.1 *If P is an invertible symmetric matrix, then the function*

$$f(x) = \frac{1}{2}x^\top Px + x^\top b$$

has a minimum value iff $P \succeq 0$, in which case this optimal value is obtained for a unique value of x , namely $x^ = -P^{-1}b$, and with*

$$f(P^{-1}b) = -\frac{1}{2}b^\top P^{-1}b.$$

Proof. Observe that

$$\frac{1}{2}(x + P^{-1}b)^\top P(x + P^{-1}b) = \frac{1}{2}x^\top Px + x^\top b + \frac{1}{2}b^\top P^{-1}b.$$

Thus,

$$f(x) = \frac{1}{2}x^\top Px + x^\top b = \frac{1}{2}(x + P^{-1}b)^\top P(x + P^{-1}b) - \frac{1}{2}b^\top P^{-1}b.$$

If P has some negative eigenvalue, say $-\lambda$ (with $\lambda > 0$), if we pick any eigenvector, u , of P associated with λ , then for any $\alpha \in \mathbb{R}$ with $\alpha \neq 0$, if we let $x = \alpha u - P^{-1}b$, then as $Pu = -\lambda u$ we get

$$\begin{aligned} f(x) &= \frac{1}{2}(x + P^{-1}b)^\top P(x + P^{-1}b) - \frac{1}{2}b^\top P^{-1}b \\ &= \frac{1}{2}\alpha u^\top P\alpha u - \frac{1}{2}b^\top P^{-1}b \\ &= -\frac{1}{2}\alpha^2\lambda \|u\|_2^2 - \frac{1}{2}b^\top P^{-1}b, \end{aligned}$$

and as α can be made as large as we want and $\lambda > 0$, we see that f has no minimum. Consequently, in order for f to have a minimum, we must have $P \succeq 0$. In this case, as $(x + P^{-1}b)^\top P(x + P^{-1}b) \geq 0$, it is clear that the minimum value of f is achieved when $x + P^{-1}b = 0$, that is, $x = -P^{-1}b$. \square

Let us now consider the case of an arbitrary symmetric matrix, P .

Proposition 4.2 *If P is a symmetric matrix, then the function*

$$f(x) = \frac{1}{2}x^\top Px + x^\top b$$

has a minimum value iff $P \succeq 0$ and $(I - PP^\dagger)b = 0$, in which case this minimum value is

$$p^* = -\frac{1}{2}b^\top P^\dagger b.$$

Furthermore, if $P = U^\top \Sigma U$ is an SVD of P , then the optimal value is achieved by all $x \in \mathbb{R}^n$ of the form

$$x = -P^\dagger b + U^\top \begin{pmatrix} 0 \\ z \end{pmatrix},$$

for any $z \in \mathbb{R}^{n-r}$, where r is the rank of P .

Proof. The case where P is invertible is taken care of by Proposition 4.1 so, we may assume that P is singular. If P has rank $r < n$, then we can diagonalize P as

$$P = U^\top \begin{pmatrix} \Sigma_r & 0 \\ 0 & 0 \end{pmatrix} U,$$

where U is an orthogonal matrix and where Σ_r is an $r \times r$ diagonal invertible matrix. Then, we have

$$\begin{aligned} f(x) &= \frac{1}{2}x^\top U^\top \begin{pmatrix} \Sigma_r & 0 \\ 0 & 0 \end{pmatrix} Ux + x^\top U^\top Ub \\ &= \frac{1}{2}(Ux)^\top \begin{pmatrix} \Sigma_r & 0 \\ 0 & 0 \end{pmatrix} Ux + (Ux)^\top Ub. \end{aligned}$$

If we write $Ux = \begin{pmatrix} y \\ z \end{pmatrix}$ and $Ub = \begin{pmatrix} c \\ d \end{pmatrix}$, with $y, c \in \mathbb{R}^r$ and $z, d \in \mathbb{R}^{n-r}$, we get

$$\begin{aligned} f(x) &= \frac{1}{2}(Ux)^\top \begin{pmatrix} \Sigma_r & 0 \\ 0 & 0 \end{pmatrix} Ux + (Ux)^\top Ub \\ &= \frac{1}{2}(y^\top, z^\top) \begin{pmatrix} \Sigma_r & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} y \\ z \end{pmatrix} + (y^\top, z^\top) \begin{pmatrix} c \\ d \end{pmatrix} \\ &= \frac{1}{2}y^\top \Sigma_r y + y^\top c + z^\top d. \end{aligned}$$

For $y = 0$, we get

$$f(x) = z^\top d,$$

so if $d \neq 0$, the function f has no minimum. Therefore, if f has a minimum, then $d = 0$. However, $d = 0$ means that $Ub = \begin{pmatrix} c \\ 0 \end{pmatrix}$ and we know from Section 3 that b is in the range of P (here, U is U^\top) which is equivalent to $(I - PP^\dagger)b = 0$. If $d = 0$, then

$$f(x) = \frac{1}{2}y^\top \Sigma_r y + y^\top c$$

and as Σ_r is invertible, by Proposition 4.1, the function f has a minimum iff $\Sigma_r \succeq 0$, which is equivalent to $P \succeq 0$.

Therefore, we proved that if f has a minimum, then $(I - PP^\dagger)b = 0$ and $P \succeq 0$. Conversely, if $(I - PP^\dagger)b = 0$ and $P \succeq 0$, what we just did proves that f does have a minimum.

When the above conditions hold, the minimum is achieved if $y = -\Sigma_r^{-1}c$, $z = 0$ and $d = 0$, that is for x^* given by $Ux^* = \begin{pmatrix} -\Sigma_r^{-1}c \\ 0 \end{pmatrix}$ and $Ub = \begin{pmatrix} c \\ 0 \end{pmatrix}$, from which we deduce that

$$x^* = -U^\top \begin{pmatrix} \Sigma_r^{-1}c \\ 0 \end{pmatrix} = -U^\top \begin{pmatrix} \Sigma_r^{-1}c & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} c \\ 0 \end{pmatrix} = -U^\top \begin{pmatrix} \Sigma_r^{-1}c & 0 \\ 0 & 0 \end{pmatrix} Ub = -P^\dagger b$$

and the minimum value of f is

$$f(x^*) = -\frac{1}{2}b^\top P^\dagger b.$$

For any $x \in \mathbb{R}^n$ of the form

$$x = -P^\dagger b + U^\top \begin{pmatrix} 0 \\ z \end{pmatrix}$$

for any $z \in \mathbb{R}^{n-r}$, our previous calculations show that $f(x) = -\frac{1}{2}b^\top P^\dagger b$. \square

We now return to our original problem, characterizing when a symmetric matrix, $M = \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix}$, is positive semidefinite. Thus, we want to know when the function

$$f(x, y) = (x^\top, y^\top) \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = x^\top Ax + 2x^\top By + y^\top Cy$$

has a minimum with respect to both x and y . Holding y constant, Proposition 4.2 implies that $f(x, y)$ has a minimum iff $A \succeq 0$ and $(I - AA^\dagger)By = 0$ and then, the minimum value is

$$f(x^*, y) = -y^\top B^\top A^\dagger By + y^\top Cy = y^\top (C - B^\top A^\dagger B)y.$$

Since we want $f(x, y)$ to be uniformly bounded from below for all x, y , we must have $(I - AA^\dagger)B = 0$. Now, $f(x^*, y)$ has a minimum iff $C - B^\top A^\dagger B \succeq 0$. Therefore, we established that $f(x, y)$ has a minimum over all x, y iff

$$A \succeq 0, \quad (I - AA^\dagger)B = 0, \quad C - B^\top A^\dagger B \succeq 0.$$

A similar reasoning applies if we first minimize with respect to y and then with respect to x but this time, the Schur complement, $A - BC^\dagger B^\top$, of C is involved. Putting all these facts together we get our main result:

Theorem 4.3 *Given any symmetric matrix, $M = \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix}$, the following conditions are equivalent:*

- (1) $M \succeq 0$ (M is positive semidefinite).

$$(2) A \succeq 0, \quad (I - AA^\dagger)B = 0, \quad C - B^\top A^\dagger B \succeq 0.$$

$$(2) C \succeq 0, \quad (I - CC^\dagger)B^\top = 0, \quad A - BC^\dagger B^\top \succeq 0.$$

If $M \succeq 0$ as in Theorem 4.3, then it is easy to check that we have the following factorizations (using the fact that $A^\dagger AA^\dagger = A^\dagger$ and $C^\dagger CC^\dagger = C^\dagger$):

$$\begin{pmatrix} A & B \\ B^\top & C \end{pmatrix} = \begin{pmatrix} I & BC^\dagger \\ 0 & I \end{pmatrix} \begin{pmatrix} A - BC^\dagger B^\top & 0 \\ 0 & C \end{pmatrix} \begin{pmatrix} I & 0 \\ C^\dagger B^\top & I \end{pmatrix}$$

and

$$\begin{pmatrix} A & B \\ B^\top & C \end{pmatrix} = \begin{pmatrix} I & 0 \\ B^\top A^\dagger & I \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & C - B^\top A^\dagger B \end{pmatrix} \begin{pmatrix} I & A^\dagger B \\ 0 & I \end{pmatrix}.$$

References

- [1] Stephen Boyd and Lieven Vandenbergh. *Convex Optimization*. Cambridge University Press, first edition, 2004.
- [2] James W. Demmel. *Applied Numerical Linear Algebra*. SIAM Publications, first edition, 1997.
- [3] Jean H. Gallier. *Geometric Methods and Applications, For Computer Science and Engineering*. TAM, Vol. 38. Springer, first edition, 2000.
- [4] H. Golub, Gene and F. Van Loan, Charles. *Matrix Computations*. The Johns Hopkins University Press, third edition, 1996.
- [5] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, first edition, 1990.
- [6] Roger A. Horn and Charles R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, first edition, 1994.
- [7] Gilbert Strang. *Linear Algebra and its Applications*. Saunders HBJ, third edition, 1988.
- [8] L.N. Trefethen and D. Bau III. *Numerical Linear Algebra*. SIAM Publications, first edition, 1997.