

Topology and geometry of quantum invariants (open problems)

Effie Kalfagianni

Michigan State University

MSRI, Berkeley, January 2020

Settings and talk theme

3-manifolds: M =compact, orientable, with empty or tori boundary.

Links: Smooth embedding $K : \coprod S^1 \rightarrow M$.

Link complements: $\overline{M \setminus n(K)}$; toroidal boundary




Figure credit: J. Cantarella, UGA

Talk: Relations between perspectives with focus on open problems.

3-manifold topology/geometry:

- Invariants arising from geometric structures:(e.g. hyperbolic volume)
- π_1 -injective embedded surfaces


Quantum topology invariants:

- Jones type knot polynomials
- Witten -Reshetikhin-Turaev invariants of 3-manifolds and knots
- TQFTs



Warm up: Geometrization in 2-d:

Every (closed, orientable) surface can be written as $S = X/G$, where X is a model geometry and G is a discrete group of isometries.



$$X = \mathbf{S}^2$$



$$X = \mathbb{E}^2$$



$$X = \mathbb{H}^2$$



- *Curvature:* $k = 1, 0, -1$

Geometrization in 3-d:

In dimension 3, there are eight model geometries:

$$X = \mathbf{S}^3, \mathbb{E}^3, \mathbb{H}^3, \mathbf{S}^2 \times \mathbb{R}, \mathbb{H}^2 \times \mathbb{R}, \text{Sol}, \text{Nil}, \widetilde{SL_2(\mathbb{R})}$$

Recall M = compact, oriented, ∂M = empty or tori

Theorem (Thurston 1980 + Perelman 2003)

*For every 3-manifold M , there is a **canonical** way to cut M along spheres and tori into pieces M_1, \dots, M_n , such that each piece is $M_i = X_i / G_i$, where G_i is a discrete group of isometries of the model geometry X_i .*

- **Canonical**: “Unique” collection of spheres and tori.
- Poincare conjecture: \mathbf{S}^3 is the only compact mode.
- **Hyperbolic** 3-manifolds form a rich and very interesting class.
- Cutting along tori, manifolds with toroidal boundary will naturally arise. Knot complements fit in this class.

Geometric decomposition picture for this talk:

Theorem (Kneser, Milnor 60's, Jaco-Shalen, Johanson 1970, Thurston 1980 + Perelman 2003)

M=oriented, compact, with empty or toroidal boundary.

- 1 *There is a unique collection of 2-spheres that decompose M*

$$M = M_1 \# M_2 \# \dots \# M_p \# (\# S^2 \times S^1)^k,$$

*where M_1, \dots, M_p are compact orientable **irreducible** 3-manifolds.*

- 2 *For $M=$ irreducible, there is a unique collection of disjointly embedded **essential** tori \mathcal{T} such that all the connected components of the manifold obtained by cutting M along \mathcal{T} , are either **Seifert fibered manifolds** or **hyperbolic**.*

- **Seifert fibered manifolds:** For this talk, think of it as


$S^1 \times$ surface with boundary + union of solid tori.

Complete topological classification [Seifert, 60']


- **Hyperbolic:** Interior admits complete, hyperbolic metric of finite volume.

Three types of knots:

Satellite Knots: Complement contains embedded “essential” tori; There is a *canonical* (finite) collection of such tori.



Torus knots: Knot embeds on standard torus in T in S^3 and is determined by its class in $H_1(T)$. Complement is SFM.



Hyperbolic knots: Rest of them.

Rigidity for hyperbolic 3-manifolds:

Theorem (Mostow, Prasad 1973)

Suppose M is compact, oriented, and ∂M is a possibly empty union of tori. If M is hyperbolic (that is: $M \setminus \partial M = \mathbb{H}^3/G$), then G is unique up to conjugation by hyperbolic isometries. In other words, a hyperbolic metric on M is essentially unique.

- By rigidity, every geometric measurement (e.g. **volume**) of a hyperbolic M is a **topological invariant**
- **Gromov norm of M** : $v_{\text{tet}} ||M|| = \text{Vol}(H)$, where
- $\text{Vol}(H)$ = sum of the hyperbolic volumes of hyperbolic parts in geometric decomposition
- v_{tet} = volume of the regular hyperbolic tetrahedron.
- If M Seifert fibered then $||M|| = 0$
- **Cutting along tori**: If M' is obtained from M by cutting along an embedded torus T then

$$||M|| \leq ||M'||,$$

with equality if T is incompressible.

Jones Polynomials—Quantum invariants

1980's: Jones polynomial (subfactor theory)— Ideas originated in physics and in representation theory led to vast families invariants of knots and 3-manifolds. (*Quantum invariants*)

For this talk we discuss:


- The *Colored Jones Polynomials*: Infinite sequence of Laurent polynomials $\{J_{K,n}(t)\}_n$ encoding the *Jones polynomial* of K and these of the links K^s that are the **parallels** of K .
- Formulae for $J_{K,n}(t)$ come from representation theory of $SU(2)$ (decomposition of tensor products of representations).

They look like

$$J_K^1(t) = 1, \quad J_K^2(t) = J_K(t) - \text{Original JP,}$$


$$J_K^3(t) = J_{K^2}(t) - 1, \quad J_K^4(t) = J_{K^3}(t) - 2J_K(t), \dots$$

- $J_{K,n}(t)$ can be calculated from any knot diagram via processes such as *Skein Theory*, *State sums*, *R-matrices*, *Fusion rules*....



The skein theory approach

- A or B resolutions, D_A, D_B , of a crossing of $D = D(K)$.



- Kauffman bracket: polynomial $\langle D \rangle \in \mathbb{Z}[t^{\pm 1/4}]$, regular isotopy invariant:

- $\langle L \amalg \bigcirc \rangle = -(t^{1/2} + t^{-1/2})\langle L \rangle := \delta \langle L \rangle$
- $\langle L \rangle = t^{-1/4} \langle D_A \rangle + t^{1/4} \langle D_B \rangle$
- $\langle \bigcirc \rangle = -t^{1/2} - t^{-1/2}$

- Chebyshev polynomials:

$$S_{n+2}(x) = xS_{n+1}(x) - S_n(x), \quad S_1(x) = x, \quad S_0(x) = 1.$$

- D^m diagram obtained from D by taking m parallel copies.
- For $n > 0$, we define (where $w = w(D) = \text{writhe}$):

$$J_K^n(t) := ((-1)^{n-1} t^{(n^2-1)/4})^w (-1)^{n-1} \langle S_{n-1}(D) \rangle$$

- $\langle S_{n-1}(D) \rangle$ is linear extension on combinations of diagrams.

The CJP predicts Volume?

- **Question:** How do the *CJP* relate to geometry/topology of knot complements?
- *Renormalized CJP*.

$$J'_{K,n}(t) := \frac{J_K^n(t)}{J_\circ^n(t)}.$$

Volume conjecture. [Kashaev+ H. Murakami - J. Murakami] Suppose K is a knot in S^3 . Then

$$2\pi \cdot \lim_{n \rightarrow \infty} \frac{\log |J'_{K,n}(e^{2\pi i/n})|}{n} = \text{Vol}(S^3 \setminus K)$$

- The conjecture is wide open:
- 4_1 (by Ekholm), knots up to 7 crossings (by Ohtsuki)
- torus knots (Kashaev-Tirkkonen); satellites of torus knots (Zheng).

• Some difficulties:

- For families of **links** we have $J_K^n(e^{2\pi i/n}) = 0$, for all n .
- “State sum” for $J_K^n(e^{2\pi i/n})$ has oscillation/cancelation.
- No good behavior of $J_K^n(e^{2\pi i/n})$ with respect to geometric decompositions.

Colored Jones polynomials are q-homonomic

- (Garoufalidis-Le, 04): The sequence $\{J_{K,n}(t)\}_n$ has a *recursive relation*. For a fixed knot K the colored Jones function $J_K(n)$ satisfies a non-trivial linear recurrence relation of the form

$$a_d(t^{2n}, t)J_K(n+d) + \cdots + a_0(t^{2n}, t)J_K(n) = 0$$

for all n , where $a_j(u, v) \in \mathbb{C}[u, v]$.

- **Example:** For K =the trefoil knot

$$J_K^n = t^{-6(n^2-1)} \sum_{j=-\frac{n-1}{2}}^{\frac{n-1}{2}} t^{24j^2+12j} \frac{t^{8j+2} - t^{-(8j+2)}}{t^2 - t^{-2}}.$$

- The relation is

$$(t^{8n+12} - 1)J_K^{n+2} + (t^{-4n-6} - t^{-12n-10} - t^{8n+10} + t^{-2})J_K^{n+1} - (t^{-4n+4} - t^{-12n-8})J_K^n = 0.$$

Coarse relations: Colored Jones polynomial

For a knot K , and $n = 1, 2, \dots$, we write its *n -colored Jones polynomial*:

$$J_K^n(t) := \alpha_n t^{m_n} + \beta_n t^{m_n-1} + \dots + \beta'_n t^{k_n+1} + \alpha'_n t^{l_n} \in \mathbb{Z}[t, t^{-1}]$$

- (Garoufalidis-Le, 04): Each of $\alpha'_n, \beta'_n \dots$ satisfies a *linear recursive relation* in n , with integer coefficients .

$$(\text{e. g. } \alpha'_{n+1} + (-1)^n \alpha'_n = 0).$$

- Given a knot K any diagram $D(K)$, there exist **explicitly given** functions $M(n, D)$ and $L(n, D)$ $m_n \leq M(n, D)$ and $l_n \geq L(n, D)$. For “nice” knots, where we have $m_n = M(n, D)$ and $l_n = L(n, D)$, we have *stable coefficients*
- (Dasbach-Lin, Armond) If $m_n = M(n, D)$, then

$$\beta'_K := |\beta'_n| = |\beta'_2|, \quad \text{and} \quad \beta_K := |\beta_n| = |\beta_2|,$$

for every $n > 1$.

- **Remark.** “nice” turns out to be what Lickorish-Thistlethwaite call *adequate knots*.

A Coarse Volume Conjecture?

Theorem (Lackenby, Dasbach-Lin, Futer-K.-Purcell, Giambone, 05-'15')

There universal constants $A, B > 0$ such that for any hyperbolic link that is “nice” we have

$$A(\beta'_K + \beta_K) \leq \text{Vol}(S^3 \setminus K) < B(\beta'_K + \beta_K).$$

Question. Does there exist function $B(K)$ of the coefficients of the colored Jones polynomials of a knot K , that is easy to calculate from a “nice” knot diagram such that for hyperbolic knots, $B(K)$ is coarsely related to hyperbolic volume $\text{Vol}(S^3 \setminus K)$?

Are there constants $C_1 \geq 1$ and $C_2 \geq 0$ such that

$$C_1^{-1}B(K) - C_2 \leq \text{Vol}(S^3 \setminus K) \leq C_1B(K) + C_2,$$

for all hyperbolic K ?

- C. Lee, Proved CVC for classes of links that don't satisfy the standard “nice” hypothesis (2017)

How Strong is the Degree of CJP?

- $d_+[J_K^n]$ = maximum degree of CJP and $d_+[J_K^n]$ = minimum degree
- **Problem.** The degree $d_+[J_K^n]$ detects the unknot: If $d_+[J_K^n] = d_+[J_\circ^n] = 0.5n$, for all n , then $K = \circ$.
- **Problem.** The degrees $d_+[J_K^n]$, $d_-[J_K^n]$ detect all torus knots $T_{p,q}$: If

$$d_+[J_K^n] = d_+[J_{T_{p,q}}^n] \text{ and } d_-[J_K^n] = d_-[J_{T_{p,q}}^n],$$

then, $K = T_{p,q}$ (or $K = T_{q,p}$)

- **Problem.** Does the span $d_+[J_K^n] - d_-[J_K^n]$ “characterize” alternating knots? Is the knot K is alternating if and only if there are $a, b \in \mathbb{Z}$ (depending only on K) such that

$$a + b = 1 \text{ and } d_+[J_K^n] - d_-[J_K^n] = an^2 + bn - (b + a)?$$

- **Problem.** Do $d_+[J_K^n]$, $d_-[J_K^n]$ detect the figure-8 knot? If $d_\pm[J_K^n] = d_\pm[J_{4_1}^n]$, then is $K = 4_1$?
- All are implications of variations of *Slopes Conjectures*

The topology of the degree of CJP

- $d_+[J_K^n]$ = maximum degree of CJP
- The q -holonomicity property of CJP implies:
- Given K there is $N_K > 0$, such that, for $n \geq N_K$,

$$d_+[J_K^n] = a_K(n) n^2 + b_K(n)n + c_K(n),$$

- where $a_K(n), b_K(n), c_K(n) : \mathbf{N} \rightarrow \mathbb{Q}$ are **periodic** functions.
- Similarly, $d_-[J_K^n]$ = maximum degree of CJP:

$$d_-[J_K^n] = a_K^*(n) n^2 + b_K^*(n)n + c_K^*(n),$$

- where $a_K^*(n), b_K^*(n), c_K^*(n) : \mathbf{N} \rightarrow \mathbb{Q}$ are **periodic** functions.
- We have finitely many cluster points


$$js_K = \{4a_K(n)\}' \quad \text{and} \quad js_K^* = \{4a_K^*(n)\}',$$

- (called **Jones Slopes**) and finitely many cluster points

$$xs_K = \{2b_K(n)\}', \quad xs_K^* = \{4b_K^*(n)\}'.$$

Surfaces in knot complements

- There are several properly embedded surfaces in knot complements—some non-orientable.



- Definition.** A surface S , properly embedded in $\mathbf{S}^3 \setminus K$ is called *essential* if inclusion induces injection

$$\pi_1(S, \partial S) \longrightarrow \pi_1(\mathbf{S}^3 \setminus K, \partial(\mathbf{S}^3 \setminus K)).$$

- Definition.** A (primitive) class in $H_1(\partial(\mathbf{S}^3 \setminus K)) \cong \mathbb{Z} \times \mathbb{Z}$, determined by an element in $s \in \mathbf{Q} \cup \{\infty\}$, is called *a boundary slope of K* if there is an *essential* surface S such that each component of ∂S represents s .

Slopes Conjectures

- **Definition.** A **Jones surface** of K is an essential surface $S \subset M_K = S^3 \setminus K$
- ∂S represents a Jones slope $4a(n) = a/b \in js_K$, with $b > 0$, $\gcd(a, b) = 1$, and

$$\frac{\chi(S)}{|\partial S|b} = 2b_K(n).$$

- Similarly, for a Jones slope $4a(n) = a^*/b^* \in js_K$, with $b^* > 0$, $\gcd(a^*, b^*) = 1$, and

$$\frac{\chi(S)}{|\partial S|b^*} = -2b_K^*(n).$$

- **Strong Slope Conjecture.**
- (Garoufalidis) All Jones slopes are boundary slopes.
- ($K+Tran$) *All Jones slopes are realized by Jones surfaces.*
- **Remark/Problem.** No knots with more than two Jones slopes are known.

Figure-8/alternating

- **Question.** Are there knots with total number of Jones slopes > 2 ?
- **Definition.** A Jones surface S of a knot K is called *characteristic* if the number of sheets $b|\partial S|$ divides the *Jones period* of K .
- For all the knots the SSC is known, the Jones surfaces can be taken to be characteristic!
- **Question.** Is every Jones slope realized by a characteristic Jones surface?
- Howie and Greene gave a characterization of alternating knots in terms of their (spanning) surfaces. Their result, together with positive answer to last question, imply that if K satisfies the SSC and $d_{\pm}[J_K^n] = d_{\pm}[J_{4_1, n}]$, then K is isotopic to 4_1 .
- A positive answer to last question, together with Howie's theorem, imply the CJP characterization of alternating knots:

$$K \text{ is alternating iff } d_+[J_K^n] - d_-[J_K^n] = an^2 + (1 - a)n - 1?$$

- **SSC known for:**
- Alternating knots (Garoufalidis)
- Adequate knots (Futer-K-Purcell)
- Knots up to nine crossings (Garoufalidis, Tran-K., Howie)
- Montesinos knots (Lee-van der Veen, Garoufalidis-Lee-van der Veen, Leng-Yang-Liu)
- Iterated torus knots
- Graph knots (Motegi-Takata, Baker-Motegi-Takata)
- families of non-Montesinos knots, non-adequate knots (Howie-Do, Lee)
- **SSC is closed under:**
- Connect sums (Motegi-Takata)
- Cabling operations (Tran-K.)
- Whiterhead doubling (Baker-Motegi-Takata)

A Volume Conjecture for all 3-manifolds

- (Turaev-Viro, 1990): For odd integer r and $q = e^{\frac{2\pi i}{r}}$

$$TV_r(M) := TV_r(M, q),$$

a real valued invariant of compact oriented 3-manifolds M

- $TV_r(M, q)$ are combinatorially defined invariants and can be computed from triangulations of M by a *state sum* formula. Sums involve *quantum 6j-symbols*. Terms are highly “oscillating” and there is term cancellation. **Combinatorics have roots in representation theory of quantum groups.**
- We work with the $SO(3)$ quantum group.
- (Q. Chen- T. Yang, 2015): compelling experimental evidence supporting
- **Volume Conjecture** : For M compact, orientable

$$2\pi \cdot \lim_{r \rightarrow \infty} \frac{\log(TV_r(M, e^{\frac{2\pi i}{r}}))}{r} = v_{\text{tet}} \|M\|,$$

where r runs over odd integers.

What we know:

The Conjecture is verified for the following.

- (*Detcherry-K.-Yang, 2016*) (First examples) of **hyperbolic** links in S^3 : The complement of 4_1 knot and of the Borromean rings.
- (*Ohtsuki, 2017*) Infinite family of closed **hyperbolic** 3-manifolds: Manifolds obtained by *Dehn filling* along the 4_1 knot complement.
- (*Belletti-Detcherry-K- Yang, 2018*) Infinite family of cusped **hyperbolic** 3-manifolds that are **universal**: They produce all M by Dehn filling!
- (*Kumar 2019*) Infinite families of octahedral **hyperbolic** links in S^3 .
- (*Detcherry-K, 2017*) All links **zero Gromov norm** links in S^3 and in connected sums of copies of $S^1 \times S^2$.
- (*Detcherry, Detcherry-K, 2017*) Several families of 3-manifolds with **non-zero Gromov**, with or with or without boundary.
- For links in S^3 Turaev-Viro invariants relate to colored Jones polynomials (**Next**)

Link complements in S^3 :

$TV_r(S^3 \setminus K, e^{\frac{2\pi i}{r}})$ is expressed via (multi)-colored JP. For knots we have:

Theorem (Detcherry-K.-Yang, 2016)

For $K \subset S^3$ and $r = 2m + 1$ there is a constant η_r independent of K so that

$$TV_r(S^3 \setminus K, e^{\frac{2\pi i}{r}}) = \eta_r^2 \sum_{n=1}^m |J_K^n(e^{\frac{4\pi i}{r}})|^2.$$


- Theorem implies $TV_r(S^3 \setminus K, e^{\frac{2\pi i}{r}}) \neq 0$ for **any** link in S^3 ! The quantity $\log(TV_r(S^3 \setminus K))$ is well defined.
- The values of CJP in Theorem are different that these in “original” volume conjecture. Calculations of Detcherry-K.-Yang and K. H. Wong (*Whitehead chains*) prompt the following.
- **Question.** Is it true that for any hyperbolic link L in S^3 ,

$$2\pi \cdot \lim_{m \rightarrow \infty} \frac{\log |J_L^m(e^{\frac{4\pi i}{2m+1}})|}{m} = \text{Vol}(S^3 \setminus L)?$$

- where J_L^m denotes the unicolored CJP of L .

Building blocks of TV invariants relate to volumes

- Color the edges of a triangulation with certain “quantum ” data



- Colored tetrahedra get “6j-symbol” $\mathbf{Q} := Q(a_1, a_2, a_3, a_4, a_5, a_6)$ = function of the a_i and r . $TV_r(M)$ is a weighted sum over all tetrahedra of triangulation (*State sum*).
- (*BDKY*) Asymptotics of \mathbf{Q} relate to volumes of geometric polyhedra:

$$\frac{2\pi}{r} \log (|\mathbf{Q}|) \leq v_{\text{oct}} + O\left(\frac{\log r}{r}\right).$$

- Proved VC for “octahedral” 3-manifolds, where TV_r have “nice” forms. **In general, hard to control term cancellation in state sum.**

A more Robust statement?:

$$LTV(M) = \limsup_{r \rightarrow \infty} \frac{2\pi}{r} \log(TV_r(M)), \quad \text{and} \quad ITV(M) = \liminf_{r \rightarrow \infty} \frac{2\pi}{r} \log(TV_r(M))$$

Conjecture: There exists universal constants $B, C > 0$ and $E \geq 0$ such that for any compact orientable 3-manifold M with empty or toroidal boundary we have

$$B \|M\| - E \leq ITV(M) \leq LTV(M) \leq C \|M\|.$$

- Half is done:

Theorem (Detcherry-K., 2017)

There exists a universal constant $C > 0$ such that for any compact orientable 3-manifold M with empty or toroidal boundary we have

$$ITV(M) \leq LTV(M) \leq C \|M\|,$$

In particular, if $ITV(M) > 0$ then $\|M\| > 0$.

Outline of last theorem:

- 1 Use asymptotics of the quantum $6j$ -symbols to show

$$ITV(M) \leq LTV(M) < v_8(\# \text{ of tetrahedra needed to triangulate } M).$$

- 2 View TV invariants as part of the $SO(3)$ -Witten- Reshetikhin-Turaev and TQFT (*Roberts*). Use approach of Blanchet-Habegger-Masbaum-Vogel.
- 3 Use TQFT properties to show that if M is a Seifert fibered manifold, then

$$LTV(M) = \|M\| = 0.$$

- 4 Show that If M contains an embedded torus T and M' is obtained from M by cutting along T then

$$LTV(M) \leq LTV(M').$$

- 5 Use a theorem of Thurston to show that there is $C > 0$ such that for any hyperbolic 3-manifold M

$$LTV(M) \leq C\|M\|.$$

- 6 Use parallel behavior of $LTV(M)$ and $\|M\|$ under geometric decomposition.

Exponential growth and AMU conjecture:

- Σ =compact, oriented surface and $\text{Mod}(\Sigma)$ its mapping class group.
- For $f \in \text{Mod}(\Sigma)$ let M_f =mapping torus of f . The Invariants $TV_r(M)$ grow exponentially in r , iff

$$ITV(M_f) := \liminf_{r \rightarrow \infty} \frac{2\pi}{r} \log(TV_r(M)) > 0.$$

- **Exponential growth conjecture.** For $f \in \text{Mod}(\Sigma)$, we have $ITV(M_f) > 0$ if and only if $\|M_f\| > 0$.
- For odd integer r , and a primitive $2r$ -th root of unity there is quantum representation

$$\rho_r : \text{Mod}(\Sigma) \rightarrow \text{PGL}_{d_r}(\mathbb{C}).$$

- **AMU conjecture** (*Aderesen-Masbaum-Ueno*) A mapping class $\phi \in \text{Mod}(\Sigma)$ has pseudo-Anosov parts if and only if, for any big enough r , we have $\rho_{r,c}(\phi)$ has infinite order.
- Exponential growth conjecture implies AMU conjecture. Detcherry-K. gave many constructions of manifolds with $ITV(M) > 0$ and used these constructions to build substantial evidence for AMU conjecture. 

Some References

- S. Garoufalidis and T. Le, *A survey of q-holonomic functions*, L'Enseignement Mathématique, 62 (2016) 501-525.
- D. Futer, E. Kalfagianni, and J. Purcell, "*Jones polynomials, volume, and essential knot surfaces: a survey.*" Proceedings of Knots in Poland III. Vol 100, 51-77 (2014)..
- D. Futer, E. Kalfagianni, and J. Purcell, "*Guts of surfaces and the colored Jones polynomial.*" Research Monograph, Lecture Notes in Mathematics (Springer), vol. 2069, 2013, x+170 pp.
- "*Interactions between hyperbolic geometry, Quantum topology and number theory*, , Contemp. Math., 541, AMS., Providence, RI, 2011.
- Baker-Montegi-Takata, *The Strong Slope Conjecture for twisted generalized Whitehead doubles*, Quantum Topology, to appear.
- G. Belletti, R. Detcherry, E. Kalfagianni, T. Yang "*Growth of quantum $6j$ -symbols and applications to the Volume Conjecture.*", Journal of Differential Geometry, to appear
- R. Detcherry, E. Kalfagianni "*Quantum representations and monodromies of fibered links.*" Advances in Math., 351, 676-701 (2019).
- R. Detcherry, E. Kalfagianni "*Gromov norm and Turaev-Viro invariants of 3-manifolds.*", Ann. Sci. de l'Ecole Normale Sup., to appear.