

# A NOTE ON QUANTUM 3-MANIFOLD INVARIANTS AND HYPERBOLIC VOLUME

EFSTRATIA KALFAGIANNI

ABSTRACT. For a closed, oriented 3-manifold  $M$  and an integer  $r > 0$ , let  $\tau_r(M)$  denote the  $SU(2)$  Reshetikhin-Turaev-Witten invariant of  $M$ , at level  $r$ . We show that for every  $n > 0$ , and for  $r_1, \dots, r_n > 0$  sufficiently large integers, there exist infinitely many non-homeomorphic hyperbolic 3-manifolds  $M$ , all of which have different hyperbolic volume, and such that  $\tau_{r_i}(M) = 1$ , for  $i = 1, \dots, n$ .

*Key words:* Brunnian link, colored Jones polynomial, Reshetikhin-Turaev-Witten invariants, hyperbolic volume.

*Mathematics Subject Classification 2000:* 57M25, 57N10

## 1. INTRODUCTION

Throughout the paper,  $M$  will denote a closed orientable 3-manifold. The Reshetikhin-Turaev-Witten  $SU(2)$ -invariants of  $M$  are complex valued numbers ([RT], [KiM]) parametrized by positive integers (levels). Given a positive integer  $r \in \mathbb{N}$ , let  $\tau_r(M)$  denote the  $SU(2)$  invariant of  $M$  at level  $r$ . In the case that  $M$  is hyperbolic, let  $\text{vol}(M)$  denote the hyperbolic volume of  $M$ . It has been speculated that  $\text{vol}(M)$  is determined by the entire collection of the invariants  $T_M := \{\tau_r(M) \mid r = 1, 2, \dots\}$  (see [Mu]). However, at the moment, it is not clear what the precise statement of a conjecture in this direction should be. In this paper we are concerned with the question of the extent to which the finite sub sequences of  $T_M$  determine  $\text{vol}(M)$ . The main result is the following theorem that shows that for most finite sub sequences the answer to this question is an emphatic *no*.

**Theorem 1.1.** *Fix  $n > 0$ . There is a constant  $C_n > 0$  such that for every  $n$ -tuple of integers  $r_1, \dots, r_n > C_n$ , there exist and infinite sequence of hyperbolic 3-manifolds  $\{M_k\}_{k \in \mathbb{N}}$  such that*

- (a) *For  $i = 1, \dots, n$ , and every  $k \in \mathbb{N}$  we have  $\tau_{r_i}(M_k) = 1$ ; and*
- (b)  *$\dots > \text{vol}(M_{k+1}) > \text{vol}(M_k) > \dots > \text{vol}(M_0) > \frac{n}{2}$ .*

---

Supported in part by NSF grant DMS-0306995 and NSF-FRG grant DMS-0456155.

Given a framed link  $L$  in  $S^3$ , there is a sequence of Laurent polynomials  $\{J_N(L, t)\}_{N \in \mathbb{N}}$ ; the colored Jones polynomials ([KiM]). For the trivial knot  $U$ , if equipped with the 0-framing, we have

$$J_N(U, t) = [N] := \frac{t^N - t^{-N}}{t - t^{-1}}.$$

To prove Theorem 1.1 we need the following:

**Theorem 1.2.** *Given integers  $n > 0$  and  $r_1, \dots, r_n > 2$ , there is a knot  $K \subset S^3$  such that for any common framing on  $K$  and  $U$  we have*

$$J_N(K, e_{r_i}) = J_N(U, e_{r_i}) \quad \text{for all } N \in \mathbb{N}.$$

Here, for  $i = 1, \dots, n$ ,  $e_{r_i} := e^{\frac{2\pi\sqrt{-1}}{r_i}}$  is a primitive  $r_i$ -th root of unity. For fixed  $n$ , if  $r_1, \dots, r_n$  are sufficiently large, then  $K$  can be chosen hyperbolic with  $\text{vol}(S^3 \setminus K) > n$ . Furthermore, if  $M$  is a hyperbolic 3-manifold obtained by  $\frac{p}{q}$ -surgery on  $K$ , for some  $|q| > 12$ , then we have

$$\text{vol}(M) \geq \left(1 - \frac{127}{q^2}\right)^{\frac{3}{2}} n. \quad (1)$$

The proof of the first part of Theorem 1.1 uses a result of Lackenby ([L]) and a construction of [K]. For the remaining claims, we need Thurston's hyperbolic Dehn surgery theorem ([Th]) and a result proved jointly with Futer and Purcell ([FKP]).

**Corollary 1.3.** *Given an integer  $n > 0$ , there is a sequence of hyperbolic 3-manifolds  $\{M_i\}_{i \in \mathbb{N}}$  and an increasing sequence of positive integers  $\{m_i\}_{i \in \mathbb{N}}$  such that*

$$\text{vol}(M_i) > \frac{n}{2} \quad \text{and} \quad \tau_{m_i}(M_i) = 1,$$

for all  $i \in \mathbb{N}$ .

**Acknowledgment.** I thank Oliver Dasbach for helpful discussions.

## 2. THE PROOFS

**2.1. Some properties of the colored Jones polynomials.** A crossing disc of a knot  $J$  is an embedded disc  $D \subset S^3$  that intersects  $J$  only in its interior exactly twice geometrically and with zero algebraic intersection number. The curve  $\partial D$  is a crossing circle for  $J$ . A knot  $K$  is said to be obtained from  $J$  by a generalized crossing of order  $r \in \mathbb{Z}$  iff  $K$  is the result of  $J$  under surgery of  $S^3$  along  $\partial D$  with surgery slope  $\frac{1}{r}$ .

**Definition 2.1.** (Definition 1.1, [L]) *Let  $r \in \mathbb{N}$  and let  $J$  and  $K$  be two 0-framed knots in  $S^3$ . We say  $K$  and  $J$  are congruent modulo  $(r, 2)$ , iff  $K$  is obtained from  $J$  by a collection of generalized crossing changes of order  $r$  supported on disjoint crossing discs. In this case we will write  $J \equiv K \pmod{(r, 2)}$ .*

For  $j \in \mathbb{N}$ , let  $K^j$  denote the  $j$ -th parallel cable of  $K$  formed with 0-framing and let  $J(K^j, t)$  denote the Jones polynomial of  $K^j$ . We need the following result of Lackenby.

**Lemma 2.2.** (Corollary 2.8, [L]) *Let  $r > 2$  be an integer and let  $e_r := e^{\frac{2\pi\sqrt{-1}}{r}}$  denote a primitive  $r$ -th root of unity. Suppose that  $J$  and  $K$  are 0-framed knots in  $S^3$ . If  $J \equiv K \pmod{(r, 2)}$ , then,*

$$J(K^j, e_r) = J(J^j, e_r), \quad \text{for all } j \in \mathbb{N}.$$

We recall that for a 0-framed link  $L$  the value  $J_N(L, e_r)$  is a linear combination of  $J(L^j, e_r)$ , with the coefficients of the combination being constants independent of  $L$  (Theorem 4.15, [KiM]). Using this fact and Lemma 2.2 we have:

**Corollary 2.3.** *Let  $r > 2$  be an integer and let  $e_r := e^{\frac{2\pi\sqrt{-1}}{r}}$  denote a primitive  $r$ -th root of unity. Suppose that  $J$  and  $K$  are 0-framed knots in  $S^3$ . If  $J \equiv K \pmod{(r, 2)}$ , then, for every integer  $q > 0$ , we have  $J_N(K^q, e_r) = J_N(J^q, e_r)$ , for all  $N \in \mathbb{N}$ . In particular, we have  $J_N(K, e_r) = J_N(J, e_r)$ , for all  $N \in \mathbb{N}$ .*

**2.2. A construction of hyperbolic Brunnian links.** To prove Theorem 1.2 we will need the following lemma which summarizes results proved in [K] and uses results proved jointly with Askitas in [AK]. Below we will sketch the proof referring the reader to the original references for details.

**Lemma 2.4.** *For every  $n > 0$ , there is an  $(n + 1)$ -component link  $L_n := U \cup K_1 \cup \dots \cup K_n$  with the following properties:*

- (a)  $L_n$  is Brunnian; that is every proper sublink of  $L_n$  is a trivial link.
- (b) For  $i = 1, \dots, n$ ,  $K_i$  bounds a crossing disc  $D_i \subset S^3$  of  $U$ .
- (c)  $L_n$  is hyperbolic; that is the interior of the 3-manifold  $\overline{M_n} := S^3 \setminus \eta(L_n)$  admits a complete hyperbolic metric of finite volume. Here,  $\eta(L_n)$  denotes a tubular neighborhood of  $L_n$ .

(d) For  $n > 1$ , any collection of generalized crossing changes along any collection of discs formed by a proper subset of  $\{D_1, \dots, D_n\}$ , leaves  $U$  unknotted.

(e) Every knot obtained by a collection of  $n$  generalized crossing changes of order  $r_1, \dots, r_n > 0$  along  $D_1, \dots, D_n$ , respectively, is non-trivial.

**Proof:** For  $n = 1$ , we take  $L_1 := U \cup K_1$  to be the Whitehead link and for  $n = 2$ , we take  $L_2 := U \cup K_1 \cup K_2$  to be the 3-component Borromean link (see Figure 1). It is well known that they are both hyperbolic ([Th]). For  $n = 2$ , condition (d) is clearly satisfied and for  $n = 1$  and  $n = 2$ , condition (e) is true since the resulting knot will be a non-trivial twist knot.




FIGURE 1. The link  $L_n$  for  $n = 1$  and  $n = 2$ .

For  $n > 2$  the construction of a link  $L_n$  as claimed above, is given in the proof of Theorem 3.1 of [K]. We now outline the construction using the terminology of [AK] and [K]. Recall that a knot  $\hat{K}$  is  $n$ -adjacent to the unknot, for some  $n \in \mathbf{N}$ , if  $\hat{K}$  admits an embedding containing  $n$  generalized crossings such that changing any  $0 < m \leq n$  of them yields an embedding of the unknot. A collection of crossing circles corresponding to these crossings is called an  $n$ -trivializer. Theorem 3.5 of [K] proves the following: For every  $n > 2$ , there exists a knot  $\hat{K}$  that is  $n$ -adjacent to the unknot, and it admits an  $n$ -trivializer  $K_1 \cup \dots \cup K_n$  such that the link  $L_n^* = \hat{K} \cup K_1 \cup \dots \cup K_n$  is hyperbolic. Let  $D_1, \dots, D_n$  denote crossing discs corresponding to  $K_1, \dots, K_n$ , respectively. Now let  $L_n$  denote the  $(n + 1)$ -component link obtained from  $L_n^*$  by performing the  $n$  generalized crossing changes (along  $D_1, \dots, D_n$ ) that exhibit  $\hat{K}$  as  $n$ -adjacent to the unknot, simultaneously. We can write  $L_n := U \cup K_1 \cup \dots \cup K_n$ , where  $U$  is the unknot resulting from  $\hat{K}$  after these crossing changes. Since the links  $L_n^*$  and  $L_n$  differ by twists along a collection of discs, they have homeomorphic complements.

We conclude that  $L_n$  is hyperbolic. Thus  $L_n$  satisfies conditions (b) and (c) of the statement of the lemma. Next, let us focus on a proper sub collection of crossing discs; say, without loss of generality,  $D_1, \dots, D_{n-1}$ . Since  $K_1 \cup \dots \cup K_n$  is an  $n$ -trivializer for  $\hat{K}$ , by the definition of adjacency to the unknot,  $K_1 \cup \dots \cup K_{n-1}$  is an  $(n-1)$ -trivializer. By Theorem 2.2 of [AK], and its proof,  $U$  bounds an embedded disc, say  $\Delta$ , in the complement of  $K_1 \cup \dots \cup K_{n-1}$  such that  $D_i \cap \Delta$  is a single arc properly embedded on  $\Delta$ . It follows that  $U \cup K_1 \cup \dots \cup K_{n-1}$  is the trivial link and that every collection of generalized crossing changes supported along  $D_1, \dots, D_{n-1}$  leaves  $U$  unknotted. This proves (b) and (d). To see (e) suppose, on the contrary, that there is a collection of  $n$  generalized crossing changes of order  $r_1, \dots, r_n > 0$  along  $D_1, \dots, D_n$ , respectively, that leaves  $U$  unknotted. Let  $U'$  denote the result of  $U$  after performing the crossings changes along  $D_1, \dots, D_{n-1}$  only, leaving  $D_n$  intact. By (b) and (d),  $U' \cup K_1 \cup \dots \cup K_{n-1}$  is the trivial link. Then, a crossing change of order  $r_n > 0$  along  $D_n$  leaves  $U'$  unknotted. By the argument in the proof of Theorem 2.2 in [AK], we conclude that  $L_n$  is the trivial link. This is a contradiction since  $L_n$  is hyperbolic.  $\square$

**2.3. Proof of Theorem 1.2.** Fix  $n \in \mathbb{N}$  and let  $L_n$  be a link as in Lemma 2.4. We will consider the component  $U$  of  $L_n$  as a 0-framed unknot in  $S^3$ . Given an  $n$ -tuple of integers  $\mathbf{r} := (r_1, \dots, r_n)$ , with  $r_i > 2$ , let  $M_n(\mathbf{r})$  denote the 3-manifold obtained from  $M_n$  as follows: For  $1 \leq i \leq n$ , perform Dehn filling with slope  $\frac{1}{r_i}$  along  $\partial\eta(K_i)$ . Let  $K := U(\mathbf{r})$  denote the image of  $U$  in  $M_n(\mathbf{r})$ . Clearly,  $M_n(\mathbf{r}) := \overline{S^3 \setminus \eta(U(\mathbf{r}))}$ . Note, that since the linking number of  $K_i$  and  $U$  is zero the framing on  $K$  induced by that of  $U$  is zero. Thus  $K$  is a 0-framed knot in  $S^3$  that is obtained from  $U$  by a generalized crossing of order  $r_1, \dots, r_n$  along  $D_1, \dots, D_n$ , respectively. By Lemma 2.4 (e),  $K$  is non-trivial.

*Claim:* We have  $U \equiv K(\text{mod}(r_i, 2))$ , for every  $1 \leq i \leq n$ .

*Proof of Claim:* Fix  $1 \leq i \leq n$ . Consider  $U(i)$  the knot obtained from  $U$  as follows: For every  $1 \leq j \neq i \leq n$ , perform a generalized crossing change of order  $r_j$  along  $D_j$ . By Lemma 2.4 (d),  $U(i)$  is a (0-framed) trivial knot. Since by construction,  $K$  is obtained from  $U := U(i)$  by a generalized crossing change of order  $r_i$  along  $D_i$ , we have  $U \equiv K(\text{mod}(r_i, 2))$ . This proves the claim.

Now by Corollary 2.3 we have that, if both  $U$  and  $K$  are given the 0-framing, then  $J_N(K, e_{r_i}) = J_N(U, e_{r_i})$ , for  $i = 1, \dots, n$ . Suppose now that  $U$  and  $K$  are given any framing  $f \in \mathbb{Z}$ . By Lemma 3.27 of [KiM], under the

frame change from 0 to  $f$  both of  $J_N(K, t)$  and  $J_N(U, t)$  are changed by the factor  $t^{f(N^2-1)}$ . Thus, the equation  $J_N(K, e_{r_i}) = J_N(U, e_{r_i})$  remains true for every framing. This proves the first part of the theorem.

Now we prove the remaining claims made in the statement of the theorem: By Thurston's hyperbolic Dehn filling theorem ([Th]), there is a constant  $C_n := C(L_n) > 0$ , such that if  $r_1, \dots, r_n > C_n$  then  $M_n(\mathbf{r})$  admits a complete hyperbolic structure of finite volume. Thus  $K := U(\mathbf{r})$  is a hyperbolic knot. By the proof of Thurston's theorem, the hyperbolic metric on  $M_n(\mathbf{r})$  can be chosen so that it is arbitrarily close to the metric of  $M_n$ , provided that the numbers  $r_i \gg 0$  are all sufficiently large. Thus by choosing the  $r_i$ 's large we may ensure that the volume of  $M_n(\mathbf{r})$  is arbitrarily close to that of  $M_n$ . Since  $\partial M_n$  has  $n + 1$  components, the interior of  $M_n$  has  $n + 1$  cusps. By [Ad], we have  $\text{vol}(M_n) \geq (n + 1)v_3$ , where  $v_3 (\approx 1.01494)$  is the volume of regular hyperbolic ideal tetrahedron. Thus for  $r_1, \dots, r_n \gg C_n$  we have

$$\text{vol}(S^3 \setminus K) = \text{vol}(M_n) > nv_3 > n. \quad (2)$$

Now we turn our attention to closed 3-manifolds obtained by surgery of  $S^3$  along a knot  $K$  as above: Suppose that  $M$  is a hyperbolic 3-manifold obtained by  $\frac{p}{q}$ -surgery on  $K$ . In Theorem 3.4 of [FKP] it is shown that if  $|q| > 12$ , then  $\text{vol}(M) \geq (1 - \frac{127}{q^2})^{\frac{3}{2}} \text{vol}(S^3 \setminus K)$ . Combining this with (2) above we immediately obtain (1). This finishes the proof of the theorem.  $\square$

**2.4. Proof of Theorem 1.1.** Fix  $n > 0$  and  $r_1, \dots, r_n > 2$  and let  $K$  be a knot as in Theorem 1.2. For a positive integer  $q$ , let  $M := M_q(K)$  denote the 3-manifold obtained by surgery of  $S^3$  along  $K$  with surgery slope  $\frac{1}{q}$ . Let  $L := K^q$  denote the  $q$ -th cable of  $K$  formed with the 0-framing as before. Similarly let  $U^q$  denote the  $q$ -th cable of the unknot  $U$ . By Corollary 2.3, if both  $L$  and  $U^q$  are considered with 0-framing, we have  $J_N(L, e_{r_i}) = J_N(U^q, e_{r_i})$ , for all  $N \in \mathbb{N}$ . Now by Kirby calculus  $M$  can be obtained by surgery on the link  $L := K^q$  and where the framing on each of the  $q$  components is 1 (see, for example, Figure 8 of [G] for details.) By formula (1.9) on page 479 of [KiM],  $\tau_{r_i}(M)$  is a linear combination of the values  $\{J_N(L, e_{r_i}) \mid N < r_i\}$  with the linear coefficients depending only on  $r_i$  and the linking matrix of  $L$ . From this and our earlier observations, the value of  $\tau_{r_i}(M)$  remains the same if we replace  $L$  with  $U^q$  and keep the same framings. But then the 3-manifold obtained by surgery on this later framed link, which is the same as this obtained by  $\frac{1}{q}$ -surgery on the unknot  $U$ , is

clearly  $S^3$ . Since, by [KiM],  $\tau_r(S^3) = 1$ , for every  $r > 0$ , we have

$$\tau_{r_i}(M) = \tau_{r_i}(S^3) = 1, \quad \text{for } i = 1, \dots, m.$$

Next suppose that we have chosen the values  $r_1, \dots, r_n > 2$  large enough so that  $K$  is hyperbolic. By Thurston's hyperbolic Dehn filling theorem, if  $q \gg 0$  then the 3-manifold  $M$  is also hyperbolic. We may, without loss of generality, assume that  $|q| > 12$ . Now Theorem 1.2 implies that  $\text{vol}(M) \geq (1 - \frac{127}{q^2})^{\frac{3}{2}}n$ , and for  $q \gg 12$ , we can assure that

$$\text{vol}(M) \geq (1 - \frac{127}{q^2})^{\frac{3}{2}}n > \frac{n}{2}.$$

Next we show that the set  $A_K := \{M_q(K) \mid q \in \mathbb{Z}\}$ , contains infinitely many non-homeomorphic 3-manifolds. By [Th], for  $q \gg 0$ , we have  $\text{vol}(M_q(K)) < \text{vol}(S^3 \setminus K)$  and  $\lim_{q \rightarrow \infty} \text{vol}(M_q(K)) = \text{vol}(S^3 \setminus K)$ . Thus we can find a sequence  $\{M_k\}_{k \in \mathbb{N}}$  as claimed in the statement of Theorem 1.1.  $\square$

**2.5. Proof of Corollary 1.3.** We will use the notation and the setting established in the proofs Theorems 1.2 and 1.1: Fix  $n > 1$  and choose  $r'_1, \dots, r'_n \gg 0$  as in the proof of Theorem 1.2, so that for  $K := U(\mathbf{r})$  we have  $\text{vol}(S^3 \setminus K) > n$ , for every  $\mathbf{r} := (r_1, \dots, r_n)$  with  $r_j \geq r'_j$ . For  $i \in \mathbb{N}$  set

$$\mathbf{r}_i := (i + r_1, \dots, r_n) \quad \text{and} \quad K_i := U(\mathbf{r}_i).$$

By the arguments in the proof of Theorem 1.1, we have

$$\text{vol}(S^3 \setminus K_i) > n \quad \text{and} \quad K \equiv U(\text{mod}(m_i, 2)),$$

where  $m_i := i + r_1$ . Thus Corollary 2.3 and the argument in the proof of 1.1, imply that there is a 3-manifold  $M_i$  obtained by surgery along  $K_i$  such that: i)  $\tau_{m_i}(M_i) = 1$ ; and ii)  $M_i$  is hyperbolic with  $\text{vol}(M_i) > \frac{n}{2}$ .  $\square$

#### REFERENCES

- [Ad] C. C. Adams, *Volumes of  $N$ -cusped hyperbolic 3-manifolds*, J. London Math. Soc. (2) 38 (1988), no. 3, 555–565.
- [AK] N. Askitas and E. Kalfagianni, *On knot adjacency*, Topology and Its Applications, Vol.126 (2002), 63-81.
- [FKP] D. Futer, E. Kalfagianni and J. S. Purcell, *Dehn filling, volume, and the Jones polynomial*, Journal of Differential Geometry, in press. ArXiv:math.GT/0612138.
- [G] S. Garoufalidis, *On finite type invariants of 3-manifolds I*, Journal of Knot Theory and Its Ramifications, 4 (1996) 441-462
- [K] E. Kalfagianni, *Alexander polynomial, finite type invariants and volume of hyperbolic knots*, Algebraic and Geometric Topology, Vol. 4 (2004), 1111-1123.
- [KiM] R. Kirby and P. Melvin, *The 3-manifold invariants of Witten and Reshetikhin-Turaev for  $sl(2, C)$* , Inventiones Mathematicae, vol. 105 (1991), no. 3, 473–545.

- [L] M. Lackenby, *Fox's congruence classes and the quantum-SU(2) invariants of links in 3-manifolds*, Comment. Math. Helv. 71 (1996) 664-677.
- [Mu] H. Murakami, *Optimistic calculations about the Witten-Reshetikhin-Turaev invariants of closed three-manifolds obtained from the figure-eight knot by integral Dehn surgeries*, in *Recent Progress towards the volume conjecture* Surikaiseikikenkyusho Kokyuroku 1172, 2000, pp. 70-79.
- [RT] N. Reshetikhin and V. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. 103 (1991), no. 3, 547-597.
- [Th] W. Thurston, *The Geometry and Topology of 3-manifolds*, Available at <http://www.msri.org/publications/books/gt3m>.

MICHIGAN STATE UNIVERSITY, E. LANSING, MI, 48823

*E-mail address:* `kalfagia@math.msu.edu`