

We count the number of possible functions f with input set $[k] = \{1, 2, \dots, k\}$ and output set $[n] = \{1, 2, \dots, n\}$, subject to restrictions (injective or surjective). We may picture f as a way of distributing k balls (marked $1, \dots, k$) into n baskets (marked $1, \dots, n$). A map is injective if each basket contains at most one ball, or surjective if no basket is empty.

Indistinguishable $[k]$ means we consider two functions the same whenever they differ by a permutation of the inputs $[k]$; so we picture the k balls as identical, unmarked. Similarly, *indistinguishable* $[n]$ means we consider classes of functions up to permutation of the outputs $[n]$, so we picture the n baskets as identical and movable, and we cannot distinguish a first basket, second basket, etc.

$f : [k] \rightarrow [n]$	ALL FUNCTIONS	INJECTIONS ($k \leq n$)	SURJECTIONS ($k \geq n$)
DIST DIST	① n^k	② $n^{\underline{k}}$	③ $\text{surj}(k, n) = \sum_{i=0}^n (-1)^i \binom{n}{i} (n-i)^k$
IND DIST	④ $\binom{\binom{n}{k}}{k} = \frac{n^{\bar{k}}}{k!}$ $\binom{\binom{n}{k}}{k} = \binom{\binom{n}{k-1}}{k} + \binom{\binom{n-1}{k}}{k}$	⑤ $\binom{n}{k} = \frac{n^{\underline{k}}}{k!}$ $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$	⑥ $a(k, n) = \binom{\binom{n}{k-n}}{k} = \binom{k-1}{n-1}$
DIST IND	⑦ $\left\{ \begin{matrix} k \\ \leq n \end{matrix} \right\} = \left\{ \begin{matrix} k \\ 1 \end{matrix} \right\} + \left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} + \dots + \left\{ \begin{matrix} k \\ n \end{matrix} \right\}$	⑧ 1	⑨ $\left\{ \begin{matrix} k \\ n \end{matrix} \right\} = \frac{\text{surj}(k, n)}{n!}$
IND IND	⑩ $p_{\leq n}(k) = p_1(k) + p_2(k) + \dots + p_n(k)$	⑪ 1	⑫ $p_n(k) = p_{\leq n}(k-n)$

- Binomial coefficient, choose number $\binom{n}{k}$: $(1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k$. Multiset, multi-choose number $\binom{n}{k}$: $\frac{1}{(1-x)^n} = \sum_{k \geq 0} \binom{n}{k} x^k$, $\binom{n}{k} = \binom{n+k-1}{k}$.
- Stirling partition number (second kind) $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$: $x^n = \sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k$, $x^n = \sum_{k=1}^n (-1)^{n-k} \left[\begin{matrix} n \\ k \end{matrix} \right] x^k$, $\sum_{n \geq 0} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \frac{x^n}{n!} = \frac{1}{k!} (e^x - 1)^k$.
- Stirling cycle number (first kind) $\left[\begin{matrix} n \\ k \end{matrix} \right]$: counts permutations $w \in S_n$ with k cycles, $\left[\begin{matrix} n \\ k \end{matrix} \right] = \left[\begin{matrix} n-1 \\ k-1 \end{matrix} \right] + (n-1) \left[\begin{matrix} n-1 \\ k \end{matrix} \right]$, $\sum_{n \geq 0} \left[\begin{matrix} n \\ k \end{matrix} \right] \frac{x^n}{n!} = \log^k \left(\frac{1}{1-x} \right)$.
- Bell number $B_k = \left\{ \begin{matrix} k \\ \leq k \end{matrix} \right\} = \left\{ \begin{matrix} k \\ 1 \end{matrix} \right\} + \left\{ \begin{matrix} k \\ 2 \end{matrix} \right\} + \dots + \left\{ \begin{matrix} k \\ k \end{matrix} \right\}$. Dobinski: $B_k = \frac{1}{e} \sum_{i \geq 0} \frac{i^k}{i!}$. $\sum_{k \geq 0} B_k \frac{x^k}{k!} = e^{(e^x - 1)}$. Recurrence: $B_k = \sum_{i=0}^{k-1} \binom{k-1}{i} B_i$.
- Partition number $p(k) = p_{\leq k}(k) = p_1(k) + p_2(k) + \dots + p_k(k)$: $\sum_{k \geq 0} p(k) x^k = \prod_{i \geq 1} \frac{1}{1-x^i}$, Hardy-Ramanujan: $p(k) \sim \frac{1}{4n\sqrt{3}} \exp(\pi \sqrt{\frac{2n}{3}})$.
- Fibonacci number $F_k = F_{k-1} + F_{k-2}$ from $F_0 = 0$, $F_1 = 1$. Binet: $F_k = \frac{1}{\sqrt{5}} (\phi^k - (-\psi)^k) = \text{round}(\frac{\phi^k}{\sqrt{5}})$, where $\phi = \frac{\sqrt{5}+1}{2}$, $\psi = \frac{\sqrt{5}-1}{2}$.
- Catalan number $C_k = \sum_{i=0}^{k-1} C_i C_{k-i}$ from $C_0 = 1$; $C_k = \frac{1}{k+1} \binom{2k}{k}$. Counts: binary ordered trees ($k+1$ leaves); ordered trees (k nodes).
- Derangement number (perms with no fixed points) $D_n = n! (1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots + \frac{(-1)^n}{n!})$. $\sum_{k \geq 0} D_k \frac{x^k}{k!} = \frac{e^{-x}}{1-x}$. $D_n = (n-1)(D_{n-1} + D_{n-2})$.
- Cayley: labeled, unordered trees $T_n = n^{n-2}$. Unlabeled, unordered rooted trees r_n : $\sum_{n \geq 1} r_n x^n = x \prod_{i \geq 1} \frac{1}{(1-x^i)^{r_i}}$; $r_{n+1} = \frac{1}{n} \sum_{k=1}^n \sum_{j|k} j r_j r_{n-k+1}$.