

**COMMUTATIVE ALGEBRA**  
**HOMEWORK\*6, DUE 2-27-08**

- (1) Let  $k$  be a field,  $R = k[x, y, z]$  the polynomial ring in 3 variables over  $k$ . Show:
- (a)  $x(x-1), xy-1, xz$  is a regular sequence in  $R$ .
  - (b)  $x(x-1), xz, xy-1$  is *not* a regular sequence in  $R$ .

- (2) Let  $k$  be a field,  $R = k[x, y]_{(x, y)}$  the localized polynomial ring, and  $S = R[z]/(xz, yz, z^2)$ , where  $z$  is a variable over  $R$ . Determine  $\dim(S)$  and  $\text{depth}(S)$ .

- (3) Let  $k$  be a field and  $x_1, x_2, x_3, x_4$  variables over  $k$ . Show that the ring  $A = (k[x_1, x_2, x_3, x_4]/(x_1x_3, x_1x_4, x_2x_3, x_2x_4))_{(x_1, x_2, x_3, x_4)}$  is not a CM-ring.

**Definition.** Let  $A$  be a ring and  $M$  an  $A$ -module. A sequence  $a_\bullet = a_1, \dots, a_n \in A$  is called *weakly  $M$ -regular* or a *weak  $M$ -sequence*, if for all  $1 \leq i \leq n$  the element  $a_i$  is a nonzero divisor in  $M/(a_1, \dots, a_{i-1})M$ .

- (4) Let  $A$  be a ring,  $M$  an  $A$ -module, and  $a_\bullet$  a weakly  $M$ -regular sequence. Let

$$N_2 \longrightarrow N_1 \longrightarrow N_0 \longrightarrow M \longrightarrow 0$$

is an exact sequence of  $A$ -modules. Show that the induced sequence

$$N_2/(a_\bullet)N_2 \longrightarrow N_1/(a_\bullet)N_1 \longrightarrow N_0/(a_\bullet)N_0 \longrightarrow M/(a_\bullet)M \longrightarrow 0$$

is exact.

- (5) Let  $A$  be a ring and

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

an exact sequence of  $A$ -modules. Suppose that the sequence  $a_\bullet$  is a weakly regular for  $M'$  and  $M''$ . Show that  $a_\bullet$  is weakly  $M$ -regular.

(6) Extend the result of problem (4) to show: Let  $A$  be a ring, and  $a_\bullet = a_1, \dots, a_n \in A$  a sequence of elements in  $A$ . If the complex

$$N_\bullet : \dots \rightarrow N_m \xrightarrow{\varphi_m} N_{m-1} \rightarrow \dots \rightarrow N_0 \rightarrow N_{-1} \rightarrow 0$$

is exact and  $a_\bullet$  is weakly  $N_i$ -regular for all  $i$ , then the complex  $N_\bullet \otimes_A A/(a_\bullet)$  is exact.

(7) Let  $A$  be a ring,  $M$  an  $A$ -module.

- (a) Prove that if  $a_\bullet$  is weakly  $M$ -regular then  $\text{Tor}_1^A(M, A/(a_\bullet)) = 0$ .
- (b) If, in addition,  $a_\bullet$  is a weak  $A$ -sequence, prove that  $\text{Tor}_i^A(M, A/(a_\bullet)) = 0$  for all  $i \geq 1$ .

(8) Let  $A$  be a ring,  $M$  an  $A$ -module,  $a_1, a_2, \dots, a_n \in A$ . Set  $I = (a_1, \dots, a_n) \subseteq A$  and assume that  $IM \neq M$ . Prove that the sequence  $a_1, \dots, a_n$  is  $M$ -quasi regular if and only if  $a_1 + I^2, \dots, a_n + I^2 \in I/I^2$  is  $\text{gr}_I(M)$ -regular.