

SOME FREE BOUNDARY PROBLEMS RECAST AS NONLOCAL PARABOLIC EQUATIONS

HÉCTOR A. CHANG-LARA, NESTOR GUILLEN, AND RUSSELL W. SCHWAB

ABSTRACT. In this work we demonstrate that a class of some one and two phase free boundary problems can be recast as nonlocal parabolic equations on a submanifold. The canonical examples would be one-phase Hele Shaw flow, as well as its two-phase analog. We also treat nonlinear versions of both one and two phase problems. In the special class of free boundaries that are graphs over \mathbb{R}^d , we give a precise characterization that shows their motion is equivalent to that of a solution of a nonlocal (fractional), nonlinear parabolic equation for functions on \mathbb{R}^d . Our main observation is that the free boundary condition defines a nonlocal operator having what we call the Global Comparison Property. A consequence of the connection with nonlocal parabolic equations is that for free boundary problems arising from translation invariant elliptic operators in the positive and negative phases, one obtains, in a uniform treatment for all of the problems (one and two phase), a propagation of modulus of continuity for viscosity solutions of the free boundary flow.

1. INTRODUCTION

In this work we demonstrate that some free boundary problems can be formulated as nonlocal parabolic equations. For both simplicity and technical restrictions, we focus on those problems that describe the motion of the graph of a function over \mathbb{R}^d . Further assumptions appear in Section 2.2. We elaborate on these assumptions and the possibility to reduce and/or modify them in Section 12. Specifically, we investigate the class of free boundary problems that have the following form:

$$U : \mathbb{R}^{d+1} \times [0, T] \rightarrow \mathbb{R},$$

and U solves (in a way to be made precise in Section 9),

$$\begin{cases} F_1(D^2U, \nabla U) = 0 & \text{in } \{U > 0\} \\ F_2(D^2U, \nabla U) = 0 & \text{in } (\{U > 0\}^c)^\circ \\ V = G(\partial_n^+ U, \partial_n^- U) & \text{on } \Gamma = \partial\{U > 0\}, \end{cases} \quad (1.1)$$

where n is the inward normal direction to the positivity region, $\{U > 0\}$. The derivatives, $\partial_n^\pm U$, are computed from respectively the positive and the negative phases, normalized so that both $\partial_n^+ U$ and $\partial_n^- U$ are positive quantities. This means that the free boundary, denoted as $\Gamma(t)$, evolves by the normal velocity field given by V , which of course depends on U (so, when U is nice enough, these dynamics track the evolution of the graph of $\partial\{U(\cdot, t) > 0\}$). Or, if one prefers to think in a level set formulation of the flow, this velocity condition would mean that

$$\text{on } \partial\{U(\cdot, t) > 0\}, \text{ the equation is } \partial_t U = G(\partial_n^+ U, \partial_n^- U) |\nabla U|.$$

As one can see, ∇U^\pm play the roles of the gradient of the pressures in their respective phases; these quantities drive the flow, and the velocity is determined by the balance law, G . Some common examples are expanded upon in Section 3.

Date: Sunday 21st July, 2019, arXiv Ver. 2.

2010 Mathematics Subject Classification. 35B51, 35R09, 35R35, 45K05, 47G20, 49L25, 60J75, 76D27, 76S05 .

Key words and phrases. Global Comparison Principle, Integro-differential operators, Dirichlet-to-Neumann, Free Boundaries, Hele-Shaw, Fully Nonlinear Equations.

Two of the authors acknowledge partial support from the NSF leading to the completion of this work: N. Guillen DMS-1700307; R. Schwab DMS-1665285.

Here, the operators, F_1 and F_2 , are translation invariant and either both uniformly elliptic or one is uniformly elliptic and the other is zero only when U is identically zero (for one-phase problems). More detailed assumptions appear in Section 2.2. The canonical example would be $F_1(D^2U, \nabla U) = \Delta U$ and the operator $F_2(D^2U, \nabla U)$ is only zero when U^- is identically zero, which would give what many authors call the one-phase Hele-Shaw flow. The two-phase analog would be when, e.g. $F_1(D^2U, \nabla U) = F_2(D^2U, \nabla U) = \Delta U$. The nonlocal parabolic equations that arise take place on \mathbb{R}^d , and they have the form,

$$\begin{cases} f : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R} \\ \partial_t f = G(I^+(f)) \cdot \sqrt{1 + |\nabla f|} \text{ in } \mathbb{R}^d \times [0, T] \text{ (for the one-phase problem),} \\ \text{or } \partial_t f = G(I^+(f), I^-(f)) \cdot \sqrt{1 + |\nabla f|} \text{ in } \mathbb{R}^d \times [0, T] \text{ (for the two-phase problem),} \end{cases} \quad (1.2)$$

where G is the prescribed velocity function as above, and I^\pm are (nonlinear) fractional order nonlocal operators acting on functions $f \in C^{1,\gamma}(\mathbb{R}^d)$, and they enjoy the global comparison principle. Roughly speaking, I^\pm linearize to integro-differential operators in the class that given by

$$L(f, x) = b(x) \cdot \nabla f(x) + \int_{\mathbb{R}^d} f(x+h) - f(x) - (\nabla f(x) \cdot h) \mathbb{1}_{B_1}(h) \mu(x, dh), \quad (1.3)$$

where for each x , $\mu(x, dh)$ is a Lévy measure on \mathbb{R}^d .

We informally state our main results here, and then they will be expanded upon and split into separate, more precise results in later sections. They state, under a graph assumption for the free boundary, an equivalence between viscosity solutions of the free boundary and fractional parabolic equations, and also a preservation of modulus of continuity of the initial free boundary. This first theorem will be addressed more precisely in Sections 8, 9, and 10. We also tie together all of the arguments into one place in Section 11.

Theorem 1.1. *If F_1 and F_2 are uniformly elliptic and rotationally invariant in the Hessian variable, G is Lipschitz and monotone (see the assumptions of Section 2.2), and $f : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$ and $U_f : \mathbb{R}^{d+1} \times [0, T] \rightarrow \mathbb{R}$ are such that*

$$\forall t \in [0, T], \quad \Gamma(t) = \partial\{U_f(\cdot, t) > 0\} = \text{graph}(f(\cdot, t)),$$

then,

- (i) U_f is a viscosity solution of the free boundary evolution, (1.1), with appropriate boundary conditions, if and only if f is a viscosity solution of the fractional parabolic equation, (1.2);
- (ii) if additionally, $\text{graph}(f(\cdot, 0)) = \partial\{U_f(\cdot, 0)\}$ enjoys a modulus of continuity, $|f(x, 0) - f(y, 0)| \leq \omega(|x - y|)$, then for all $t \in [0, T]$ this modulus is preserved for $f(\cdot, t)$ (and hence $\partial\{U_f(\cdot, t) > 0\}$);
- (iii) if $\text{graph}(f(\cdot, 0)) = \partial\{U_f(\cdot, 0)\}$ enjoys a modulus of continuity, $|f(x, 0) - f(y, 0)| \leq \omega(|x - y|)$, then there exists a unique viscosity solution to both the respective parabolic equation and the free boundary evolution, assuming the graph assumption for $\partial\{U > 0\}$ listed above.

Remark 1.2. *The precise definitions of viscosity solutions are given in Section 8 for the parabolic nonlocal equation and Section 9 for the free boundary problem.*

Remark 1.3. *As mentioned above, the one-phase problem can be interpreted as taking F_2 to only be zero when U is identically zero. Another way to capture the one-phase problem from Theorem 1.1 is to take $G(\partial_n^+ U, \partial_n^- U) = g(\partial_n^+ U)$, which ignores whatever is the prescribed equation in the set $\{U > 0\}^C$.*

The main feature behind the results in Theorem 1.1 is the analysis of what could be called a non-linear version of the Dirichlet-to-Neumann operator, but modified to suit the free boundary problems like (1.1). This is the operator we called I , above, and we define it here. Assume that F satisfies assumptions listed in Section 2.2, and that $f \in C^{1,\gamma}(\mathbb{R}^d)$ with the property that $f \geq \delta > 0$. Then, for the domain given by f , which we define as

$$D_f = \{(x, x_{d+1}) \in \mathbb{R}^d \times \mathbb{R}_+ : 0 < x_{d+1} < f(x)\}, \quad (1.4)$$

there is a unique solution, U_f , to the equation

$$\begin{cases} F(D^2U_f, \nabla U_f) = 0 & \text{in } D_f \\ U_f = 1 & \text{on } \mathbb{R}^d \times \{0\} \\ U_f = 0 & \text{on } \text{graph}(f). \end{cases} \quad (1.5)$$

This allows us to define an operator, I , via the (inward) normal derivative of U_f , which is given by

$$I(f, x) := \partial_n U_f(x, f(x)). \quad (1.6)$$

One of the main results in our work that leads to Theorem 1.1 is the following property of I . Again, as with Theorem 1.1, we state it informally here, and revisit it with more precise assumptions later, in Sections 5, 7, and 11.

Theorem 1.4. *If F is uniformly elliptic and rotationally invariant in the Hessian variable (see the assumptions of Section 2.2), I is defined via (1.5) and (1.6), and $\gamma \in (0, 1)$ is fixed, then there exists γ' with $0 < \gamma' < \gamma$ so that*

$$I : \left(\bigcup_{\delta > 0} C^{1,\gamma}(\mathbb{R}^d) \cap \{f : f \geq \delta > 0\} \right) \rightarrow C^{\gamma'}(\mathbb{R}^d), \quad \text{and } I \text{ is locally Lipschitz.}$$

Additionally, I enjoys the following representation for a^{ij} , c^{ij} , b^{ij} , and $\mu^{ij}(dh)$ that are independent of x , and that depend on the bounded set $\{f : f \geq \delta, \|f\|_{C^{1,\gamma}} \leq m\}$:

$$I(f, x) = \min_i \max_j \left\{ a^{ij} + c^{ij} f(x) + b^{ij} \cdot \nabla f(x) + \int_{\mathbb{R}^d} (f(x+h) - f(x) - \mathbb{1}_{B_1}(h) \nabla f(x) \cdot h) \mu^{ij}(dh) \right\}.$$

Furthermore, there exists a C such that $|a^{ij}| \leq C$, $-C \leq c^{ij} \leq 0$, $|b^{ij}| \leq C$, and

$$\int_{\mathbb{R}^d} \min(|h|^{1+\gamma}, 1) \mu^{ij}(dh) \leq C \quad \text{and} \quad \int_{\mathbb{R}^d \setminus B_R} \mu^{ij}(dh) \leq C\omega(R),$$

for some modulus, ω , with $\omega(R) \rightarrow 0$ as $R \rightarrow \infty$. The constant, C , depends on the Lipschitz bound for I over the corresponding subset of $(\bigcup_{\delta > 0} C^{1,\gamma}(\mathbb{R}^d) \cap \{f : f \geq \delta > 0\})$.

Remark 1.5. *We want to emphasize that in most of our results, above and below, there is a technical assumption that F_i must be rotationally invariant in the Hessian variable. The curious reader can see its use in the proof of Theorem 5.13, but it is not clear if this was an artifact of our chosen method or if it is a true obstacle to such representations of the free boundary problem.*

Remark 1.6. *We note that to properly study the parabolic equation (1.2), the most important part is to understand the Lipschitz, comparison, and integro-differential properties of the map, I , defined in (1.6). As the reader will see at the end of Section 7, all the desired properties then also carry over to the operator*

$$G(I(f)) \cdot \sqrt{1 + |\nabla f|},$$

which is the one that appears in (1.2) and corresponds to the free boundary evolution.

Remark 1.7. *In the interest of transparency, we would like to mention what is new and what is closer to what is already known about the results that appear above.*

- *In principle, the free boundary existence and uniqueness results mentioned in Theorem 1.1 are not surprising, and most experts would probably say they closely mirror the results on existence and uniqueness found in [46]. However, to the best of our knowledge, even with such special assumptions as the graph condition, we are not aware that the fully nonlinear nor the two-phase cases have been written or appeared anywhere. Thus, this step is technically new, but maybe not substantially new.*

- *The preservation of a modulus of continuity for the free boundary again, is to the best of our knowledge, new, but given the fact that it is really a consequence of the definition of viscosity solutions, combined with the underlying translation invariance of the problem, one could also suggest that this may follow in the Hele-Shaw case from some arguments similar to [46]. However, this result seems to only appear when the modulus is a Lipschitz norm and the flow is Hele-Shaw, which was proved in [27] (under a star-shaped assumption instead of a graph assumption). Thus for the two-phase and the fully nonlinear one-phase cases, we believe this part of the results are technically new.*
- *The Lipschitz property of the mapping, I , to the best of our knowledge is completely new. The integro-differential formula in the min-max of Theorem 1.4 is new, even though previous works had noted connections with operators similar to the 1/2-Laplacian (more discussion of this appears in Section 4).*
- *The correspondence of the free boundary evolution with the viscosity solution of the parabolic equation (1.2) is relatively new, and the closest result is that of two of the authors, where in [25] they showed that certain blow-up limits of the free boundary will be viscosity solutions of a parabolic equation. The novelty in our current work is to avoid the blow-up argument, and work directly with the original free boundary as a fractional parabolic equation.*

1.1. A note on the level-set formulation of (1.1). One way to see the connection between the original free boundary evolution given by (1.1) and the nonlinear fractional parabolic equation in (1.2) is by recognizing the flow as a level-set equation. Indeed, the graph assumption means that we naturally have two different choices of defining functions for the boundary, $\partial\{U(\cdot, t) > 0\} = \partial\Omega(t)$. Of course, this is the zero level set of U , which is the obvious defining function. However, if we assume that we have parametrized $\Omega(t)$ by *some* defining function, $\Phi(X, t)$, we have

$$\partial\Omega(t) = \{\Phi(\cdot, t) = 0\}.$$

Regardless of from where the prescribed evolution of $\Omega(t)$ originates, if one insists that the normal velocity is V and the (outer!) normal vector is n , then it is not hard to check that the level set equation becomes

$$0 = \nabla\Phi \cdot (nV) + \partial_t\Phi.$$

Of course, as a defining function of a level set, we have, depending upon whether or not $\Phi > 0$ in $\Omega(t)$ or $\Phi < 0$ in $\Omega(t)$,

$$n = \pm \frac{\nabla\Phi}{|\nabla\Phi|},$$

and hence the flow reduces to

$$\partial_t\Phi = \pm V |\nabla\Phi|.$$

In our case, to be consistent with the natural parametrization, we choose $\Phi > 0$ in $\Omega(t)$, and hence a second natural choice then becomes (where $X = (x, x_{d+1})$)

$$\Phi(X, t) = f(x, t) - x_{d+1}.$$

Furthermore, we have chosen $V = G(\partial_n^+ U_f, \partial_n^- U_f)$, and so given the definition of the operator I (and correspondingly I^+ and I^-), we finally obtain

$$\partial_t f = G(I^+(f), I^-(f)) \cdot \sqrt{1 + |\nabla f|^2}.$$

2. SOME DEFINITIONS AND ASSUMPTIONS

2.1. Definitions.

Definition 2.1. *The translation operator, τ_x , is defined, for a fixed x , acting on functions on \mathbb{R}^d , as*

$$\tau_x u := u(\cdot + x).$$

Definition 2.2 (GCP). (Part 1) The global comparison property (GCP) for $I : C^{1,\gamma}(\mathbb{R}^d) \rightarrow C^0(\mathbb{R}^d)$ requires that for all $u, v \in C^{1,\gamma}(\mathbb{R}^d)$ such that $u(x) \leq v(x)$ for all $x \in \mathbb{R}^d$ and such that for some x_0 , $u(x_0) = v(x_0)$, then the operator I satisfies $I(u, x_0) \leq I(v, x_0)$. That is to say that I preserves ordering of functions on \mathbb{R}^d at any points where their graphs touch.

(Part 2) Analogously, we say that I has the GCP at x_0 if the above property is only required to hold for one fixed x_0 , instead of all possible x_0 .

Definition 2.3 (Extremal Operators). The second order (λ, Λ) -Pucci extremal operators are defined as \mathcal{M}^- and \mathcal{M}^+ , for a function, U that is second differentiable at X , via

$$\mathcal{M}^-(D^2U, X) = \min_{\lambda \text{Id} \leq B \leq \Lambda \text{Id}} (\text{tr}(BD^2U(X))) \quad \text{and} \quad \mathcal{M}^+(D^2U, X) = \max_{\lambda \text{Id} \leq B \leq \Lambda \text{Id}} (\text{tr}(BD^2U(X))).$$

When $\{e_i\}_{i=1,\dots,d+1}$ are the eigenvalues of $D^2U(X)$, an equivalent representation is

$$\mathcal{M}^-(D^2U, X) = \Lambda \sum_{e_i \leq 0} e_i + \lambda \sum_{e_i > 0} e_i \quad \text{and} \quad \mathcal{M}^+(D^2U, X) = \lambda \sum_{e_i \leq 0} e_i + \Lambda \sum_{e_i > 0} e_i.$$

Definition 2.4 (Uniformly Elliptic). When F is linear, i.e. $F(D^2U, \nabla U) = \text{tr}(A(X)D^2U(X)) + B(X) \cdot \nabla U(X)$ we say that F is (λ, Λ) -uniformly elliptic if $\|B\|_{L^\infty} \leq \Lambda$ and

$$\lambda \text{Id} \leq A(X) \leq \Lambda \text{Id} \quad \text{for all } X,$$

and when F is nonlinear, if for all $U, V \in C^2$,

$$\begin{aligned} \mathcal{M}^-(D^2U - D^2V) - \Lambda|\nabla U - \nabla V| &\leq F(D^2U, \nabla U) - F(D^2V, \nabla V) \\ &\leq \mathcal{M}^+(D^2U - D^2V) + \Lambda|\nabla U - \nabla V|. \end{aligned}$$

Next, we recall some definitions from Clarke's book on nonsmooth analysis, [29]. They are appropriately modified for our context.

Definition 2.5 (Upper Gradient). Assume that \mathcal{K} is an open convex subset of $C^{1,\gamma}(\mathbb{R}^d)$ and that $\phi : \mathcal{K} \subset C^{1,\gamma}(\mathbb{R}^d) \rightarrow \mathbb{R}$ is Lipschitz. The upper gradient of ϕ at $f \in \mathcal{K}$ in the direction of $g \in C^{1,\gamma}$, is defined as

$$\phi^0(f; g) := \limsup_{t \searrow 0} \frac{\phi(f + tg) - \phi(f)}{t}.$$

This can be seen as a function $\phi^0 : \mathcal{K} \times C^{1,\gamma} \rightarrow \mathbb{R}$.

Definition 2.6 (Subdifferential). Let ϕ be as in Definition 2.5. The generalized gradient, or as we will say, Clarke differential of ϕ at $f \in \mathcal{K}$ is the subset of $(C^{1,\gamma}(\mathbb{R}^d))^*$ given by

$$\partial\phi(f) := \{\ell \in (C^{1,\gamma}(\mathbb{R}^d))^* \mid \forall \psi \in C^{1,\gamma}(\mathbb{R}^d), \phi^0(f; \psi) \geq \langle \ell, \psi \rangle\}.$$

We will denote simply by $[\partial\phi]_{\mathcal{K}}$ the convex hull of the union of $\partial\phi(f)$,

$$[\partial\phi]_{\mathcal{K}} := \text{hull} \left(\bigcup_{f \in \mathcal{K}} \partial\phi(f) \right).$$

Remark 2.7. We want to stress that $[\partial\phi]_{\mathcal{K}}$ depends on the original set, \mathcal{K} . This plays an essential role in Sections 7 and 8.

2.2. Assumptions. We keep the following standing assumptions on F_i for $i = 1, 2$:

- (a) there exists $\lambda \leq \Lambda$ such that F_i is uniformly elliptic in the sense of Definition 2.4;
- (b) $F(0, 0) = 0$;
- (c) for all p , $F(A, p)$ is rotationally invariant in the A variable (the Hessian variable);
- (d) $G : (0, \infty)^2 \rightarrow \mathbb{R}$, and for a.e. (a, b) , $\lambda_0 \leq \frac{\partial}{\partial a} G(a, b) \leq \Lambda_0$, $\lambda_0 \leq -\frac{\partial}{\partial b} G(a, b) \leq \Lambda_0$.

2.3. Notation. Here we will collect some notation that is used in this work.

- F is an elliptic second order operator, described in Definition 2.4
- Upper case U will be functions $\mathbb{R}^{d+1} \rightarrow \mathbb{R}$, and lower case u will be functions $\mathbb{R}^d \rightarrow \mathbb{R}$.
- similarly, we try to keep the convention that upper case letters, $X \in \mathbb{R}^{d+1}$, and lower case letters, $x \in \mathbb{R}^d$.
- $D_f = \{(x, x_{d+1}) \in \mathbb{R}^d \times \mathbb{R}_+ : 0 < x_{d+1} < f(x)\}$
- $\Gamma_f = \text{graph}(f)$, and in most cases, $\Gamma_f = \partial\{U_f > 0\}$.
- $\Gamma_0 = \mathbb{R}^d \times \{0\}$
- $\Gamma_L = \mathbb{R}^d \times \{L\}$, where $L > 0$.
- n is the inward normal vector to D_f , along Γ_f , which in most cases is the inward normal vector to $\{U > 0\}$.
- For $X_0 \in \Gamma_f$,

$$\partial_n^+ U(X_0) = \lim_{t \rightarrow 0^+} \frac{U(X_0 + tn(X_0)) - U(X_0)}{t} \text{ and } \partial_n^- U(X_0) = - \lim_{t \rightarrow 0^+} \frac{U(X_0 - tn(X_0)) - U(X_0)}{t}$$

(note that in most cases, $-tn(X_0) \in \{U < 0\}$ and $U(X_0) = 0$; $\partial_n^- U$ is normalized to be non-negative).

- The translation operator, τ_h , is given in Definition 2.1.
- $C^{1,\gamma}(\mathbb{R}^d) = \{f : \mathbb{R}^d \rightarrow \mathbb{R} : \|f\|_{L^\infty} + [f]_{C^\gamma} + \|\nabla f\|_{L^\infty} + [\nabla f]_{C^\gamma} < \infty\}$
- (equivalently)

$$C^{1,\gamma}(\mathbb{R}^d) = \{f \in L^\infty(\mathbb{R}^d) : \sup_{z \in \mathbb{R}^d} \sup_{r > 0} r^{-1-\gamma} \inf_{P(x)=c+px : c \in \mathbb{R}, p \in \mathbb{R}^d} \|f - P\|_{L^\infty(B_r(z))} < \infty\}$$

- Punctually $C^{1,\gamma}(x)$ is defined in Definition 5.2.
- For $\gamma \in (0, 1)$, $\delta > 0$, $m > 0$, the convex set, $\mathcal{K}(\gamma, \delta, m)$ is defined as

$$\mathcal{K}(\gamma, \delta, m) := \{f \in C^{1,\gamma}(\mathbb{R}^d) \mid f(x) \geq \delta \ \forall x \in \mathbb{R}^d, \|f\|_{C^{1,\gamma}(\mathbb{R}^d)} \leq m\}.$$

- $\mathcal{K}(\gamma, \delta) = \bigcup_{m > 0} \mathcal{K}(\gamma, \delta, m)$.
- $\mathcal{K}(\delta) = \bigcup_{\gamma \in (0,1)} \bigcup_{m > 0} \mathcal{K}(\gamma, \delta, m)$.
- $\mathcal{K}^*(\gamma, \delta, m)$ and $\mathcal{K}_*(\gamma, \delta, m)$ appear after Definition 5.2, in equations (5.4), (5.5).

3. THREE EXAMPLES

Here, we mention three natural examples to which Theorems 1.1 and 1.4 will apply. The first two examples are treated explicitly in this paper, and the third is closely related, but not explicitly treated. We want to point out that although the equations that govern the pressure and free boundary may be linear, the resulting nonlocal parabolic flow is fully nonlinear.

3.1. One phase Hele-Shaw on a half space. To every $f : \mathbb{R}^d \rightarrow \mathbb{R}$ which is continuous, non-negative, bounded, and bounded away from zero we associate the domain

$$D_f := \{X = (x, x_{d+1}) \in \mathbb{R}^{d+1} \mid 0 < x_{d+1} < f(x)\},$$

as well as the hypersurface


$$\Gamma_f := \{X = (x, x_{d+1}) \in \mathbb{R}^{d+1} \mid x_{d+1} = f(x)\}.$$

Let $U_f : D_f \rightarrow \mathbb{R}$ be the unique bounded solution to the Dirichlet problem

$$\begin{cases} \Delta U = 0 & \text{in } D_f, \\ U = 1 & \text{on } \{x_{d+1} = 0\}, \\ U = 0 & \text{on } \Gamma_f. \end{cases}$$

Then, with n denoting the unit normal to Γ_f pointing towards D_f , define

$$I(f, x) := (\partial_n^+ U_f)(x, f(x)).$$



For sufficiently smooth f , the property to be a solution of

$$\partial_t f = I(f, x) \cdot \sqrt{1 + |\nabla f|^2}, \quad \text{on } \mathbb{R}^d \times [0, T],$$

is equivalent to $U = U_f$ being a solution of the free boundary problem

$$\begin{cases} \Delta U = 0 & \text{in } \{U > 0\}, \\ U = 1 & \text{along } \{x_{d+1} = 0\}, \\ V = \partial_n^+ U & \text{along } \partial\{U > 0\}. \end{cases}$$

This is the one-phase Hele-Shaw problem on the upper half space, and it is not hard to check that the map, I , enjoys the GCP. The operator I is easily seen to be translation invariant, and we emphasize that it is also nonlinear and nonlocal. The existence/comparison results in Sections 8-9 and regularity result in Section 10 apply to this problem.

3.2. Two-phase problems along an infinite strip. In our framework, dealing with the two-phase problem is nearly at the same level of complexity and difficulty as the one-phase problem. We fix an upper boundary, $L > 0$, and a globally Lipschitz function

$$G : (0, \infty)^2 \rightarrow \mathbb{R},$$

which satisfies a monotonicity assumption,

$$\lambda_0 \leq \frac{\partial}{\partial a} G(a, b) \leq \Lambda_0, \quad \lambda_0 \leq -\frac{\partial}{\partial b} G(a, b) \leq \Lambda_0,$$

for example, $G(a, b) = a^2 - b^2$. This function, G , will give the normal velocity of the flow, depending upon $\partial_n^\pm U$ along $\partial\{U > 0\}$.

Now, to every $f : \mathbb{R}^d \rightarrow \mathbb{R}$ which is continuous, $0 \leq f \leq L$, and bounded away from zero and L , we associate the two domains (for $\{U > 0\}$ and $\{U < 0\}$)

$$\begin{aligned} D_f^+ &= \{(x, x_{d+1}) \mid 0 < x_{d+1} < f(x)\}, \\ D_f^- &= \{(x, x_{d+1}) \mid f(x) < x_{d+1} < L\}, \end{aligned}$$


and we also associate to such f the same hypersurface, Γ_f , that is defined in the previous example. Let $U_f^+ : D_f^+ \rightarrow \mathbb{R}$ and $U_f^- : D_f^- \rightarrow \mathbb{R}$ be respectively the unique bounded solution to the Dirichlet

problem

$$\begin{cases} \Delta U^\pm = 0 & \text{in } D_f^\pm \\ U^\pm = \pm 1 & \text{on } \{x_{d+1} = 0\} \text{ (resp. } \{x_{d+1} = L\}), \\ U^\pm = 0 & \text{on } \Gamma_f. \end{cases}$$

Then, we define $H(f, x)$ by

$$H(f, x) := G(\partial_n^+ U(x, f(x)), \partial_n^- U(x, f(x))).$$



As in the previous example, as long as f is sufficiently smooth, solving

$$\partial_t f = H(f, x) \cdot \sqrt{1 + |\nabla f|^2}, \quad \text{on } \mathbb{R}^d \times [0, T],$$

is equivalent to $U = U_f$ solving the two-phase free boundary problem

$$\begin{cases} \Delta U = 0 & \text{in } \{U \neq 0\}, \\ U = 1 & \text{on } \{x_{d+1} = 0\}, \\ U = -1 & \text{on } \{x_{d+1} = L\}, \\ V = G(\partial_n^+ U, \partial_n^- U) & \text{on } \partial\{U > 0\}. \end{cases}$$


Again, the resulting operator H enjoys the GCP, is translation invariant, nonlinear, nonlocal, and the existence/comparison results in Sections 8-9 and regularity propagation result in Section 10 apply. Furthermore, as in Section 6, the reader will see that H is just a combination of two operators like the I from the first example, but interacting through the function, G .

3.3. Prandtl-Batchelor as a non-linear integro-differential equation on the sphere. This example illustrates how the ideas in this paper could possibly be applied to free boundaries which are not given by a graph over \mathbb{R}^d , but rather a graph over a sphere. The Prandtl-Batchelor is a two dimensional model in fluid mechanics that models a vortex patch occupying a convex region, the patch being surrounded by a steady flow (see [1], [7], [8], [17]). The interface of the vortex patch is what plays the role of the free boundary. Although not covered directly by our results, it is close to them in spirit, and most of the tools we use can be applied to the situation of operators that act on functions over a reference manifold.

In the Prandtl-Batchelor flow we denote the stream function of the flow by U , and it satisfies the following

$$\begin{cases} \Delta U = 0 & \text{in } \{U > 0\}, \\ \Delta U = 1 & \text{in } \{U < 0\}, \\ |\partial_n^+ U|^2 - |\partial_n^- U|^2 = 1 & \text{on } \partial\{U > 0\}. \end{cases}$$

The set $\{U < 0\}$ corresponds to the vortex patch.



This problem can be recast as a (steady state) non-linear integro-differential equation on the sphere (or circle). To see this, consider the set of continuous functions $f : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ such that

$$0 < f(x) \leq \bar{f}(x) \text{ in } \mathbb{S}^{d-1},$$

where \bar{f} is a given positive, continuous function. For each such f , we define the sets

$$\begin{aligned} D_f^- &:= \{x \in \mathbb{R}^d \mid x = re \text{ where } e \in \mathbb{S}^{d-1} \text{ and } 0 \leq r < f(e)\}, \\ D_f^+ &:= D_{\bar{f}}^- \setminus D_f^-, \end{aligned}$$

furthermore, we define functions $U_f^\pm : D_f^\pm \rightarrow \mathbb{R}$, each given as the unique solution to the Dirichlet problem


$$\begin{cases} \Delta U^- = 1 & \text{in } D_f^-, \\ U^- = 0 & \text{on } \partial D_f^-, \end{cases}$$

and

$$\begin{cases} \Delta U^+ = 0 & \text{in } D_f^+, \\ U^+ = 0 & \text{on } \partial D_f^-, \\ U^+ = 1 & \text{on } \partial D_{\bar{f}}. \end{cases}$$

Then, we define an operator $P : C^2(\mathbb{S}^{d-1}) \rightarrow C^0(\mathbb{S}^{d-1})$ by

$$P(f, x) := |\partial_n^+ U_f^+(x, f(x))|^2 - |\partial_n^- U_f^-(x, f(x))|^2$$



The operator $P(f, x)$ admits the GCP. Moreover, for sufficiently smooth f , solving

$$P(f, x) = 1 \text{ in } x \in \mathbb{S}^{d-1},$$

means that the (radial) graph of f is the boundary of a vortex patch in a Prandtl-Batchelor flow, and that U_f^\pm gives the positive and negative phases of the stream function. We note that thanks to the invariance of the Laplacian and the boundary conditions by rotations of the domain, the operator, P , will also be invariant under the action of rotations on $C^2(\mathbb{S}^{d-1})$. This rotational invariance would play the same role in the min-max and existence-uniqueness theory as does the translation invariance that we use in Sections 7, 8, 10.

4. BACKGROUND

As mentioned above, our goal is to show how some free boundary problems correspond to parabolic integro-differential equations for a scalar over a submanifold. In some aspects, this observation is not new, but in other aspects it does seem new— we will try to put it in the context of some existing results. For the purposes of the following discussion we will focus solely on free boundary problems involving one scalar and one hypersurface. This, of course, is not the case for all free boundary problems, but will be so for all of the free boundary problems we consider here. We will also limit our discussion to those free boundary problems that non-stationary.

Examples of free boundary problems include a large class of fluid problems such as cavitation [9] obstacle problems [15, 52], the porous medium equation [13, 59], the Muskat problem [56], problems with surface tension [51], singular perturbation problems [18], Prandtl-Batchelor flows [1], [7], [8], [17], sharp interface limits of phase field models [2, 19, 20], and more.

Many of these problems admit a variational formulation, often in terms of variational inequalities (see for instance Duvaut and Lions [35] or Kinderlehrer and Stampacchia [48] for further discussion of such problems). In many situations, not necessarily mutually exclusive with the former, the problem has a comparison principle at the level of the scalar field. Such a comparison principle has made it possible to define viscosity solutions, and the origins of this seem to be in the work of Caffarelli for the stationary two-phase problem in [14]. This notion of solution was then used for two-phase Stefan problems in Athanasopoulos-Caffarelli-Salsa in [5], and was later adapted to the Porous Medium equation in work of Caffarelli-Vazquez [13], as well as for the Hele-Shaw problem in work of Kim [46] and the two-phase Stefan problem of Kim-Pozar [47]. In particular, the work of [46] shows existence, uniqueness, and a comparison principle for viscosity solutions for the one phase Hele-Shaw problems. Regularity for these free boundary problems was studied, for Stefan, in [5], and for Hele-Shaw in [27], [28], [45]. It is interesting to point out that although in a slightly different fashion than what we treat here, there was an occurrence of something resembling an integro-differential equation in [27]. In [27, Lemma 10.4], for the one-phase Hele-Shaw flow, they establish that (using our notation above) when

$\partial\{U > 0\}$ is Lipschitz with a small enough Lipschitz norm, U satisfies an equation of the form

$$\partial_t U = \int_{\partial\{U > 0\}} |\nabla U(x, t)|^2 k(x, t; y) d\sigma(y);$$

where k is the Poisson kernel for $\partial\{U > 0\}$, and σ is the surface measure on $\partial\{U > 0\}$.

There are free boundary problems where the scalar is constant (say, zero) along the free boundary. The free boundary is then a level set of the solution of a PDE, and accordingly, statements about the free boundary can be reduced to statements about the scalar function itself; such as in the Porous Medium equation or the Hele-Shaw equation. Matters are different in problems where the free boundary is not a level set of the scalar, such as the Muskat problem, and where the above approach is not as directly applicable. Accordingly, the emphasis does not reside exclusively on the scalar or underlying potential function, but on the free boundary itself, and on how its shape determines the scalar and then its evolution.

A representative example of this latter situation can be found in work of Cordoba and Gancedo for the Muskat problem. In [31], the authors use contour-integration to describe the evolution equation for the interface for the Muskat problem. This requires in particular that the interface at time t is given by the graph of a function $f(x, t)$ defined over \mathbb{R}^d . Then, the scalar f is shown to solve the integro-differential equation [31, Section 2, Equation 13]

$$\partial_t f(x, t) = \frac{\rho_2 - \rho_1}{4\pi} \text{P.V.} \int_{\mathbb{R}^2} \frac{(\nabla f(x, t) - \nabla f(x - y, t)) \cdot y}{(|y|^2 + (f(x, t) - f(x - y, t)))^{\frac{2}{3}}} dy.$$

Here, ρ_1 and ρ_2 represent the respective (constant) densities of each of the fluids. As observed in [31, Section 2], the linearization of the above equation at a constant f is the fractional heat equation.

The Muskat problem, also known as the Muskat-Leibenzon problem, describes the interface bounding a fluid, see [56] for a thorough discussion and further references. The Muskat problem is an accurate model for the two-phase regime of a “Hele-Shaw cell”, and is accordingly known also as the two-phase Hele-Shaw problem. In this problem the fluid is assumed to be irrotational, but the respective pressure is no longer assumed to be constant along the free boundary. In [4] Ambrose obtained local well-posedness for the Muskat problem with initial data in H^2 that satisfies the Rayleigh-Taylor condition. Siegel, Caffisch, and Howison [56] studied global solutions with periodic boundary conditions and initial datum close to equilibrium.

The approach in [31] was used in several subsequent results, among which we highlight a few. Constantin et al. [30] showed the existence of global weak solutions which are globally Lipschitz in space as long as the initial data had a Lipschitz constant strictly smaller than 1. In contrast, Castro et al [22, Theorem 2.1] showed the 2D Muskat problem may develop a singularity (in C^1) in finite time, even if the initial data is smooth. More recently, Cameron [21], studied a closely related problem, using integro-differential methods combined with the modulus preservation technique of [49].

An approach amenable to many problems involves changing variables, either to Lagrangian coordinates (for fluid problems) or through the Hanzawa transform, pulling back the free boundary onto a fixed, reference interface, and writing the free boundary problem as a system on a fixed domain. The book of Prüss and Simonett [54] has a thorough presentation of this approach in combination with L^p -maximal regularity. The Hanzawa transform, for instance, entails fixing a reference free boundary Γ_0 , and considering interfaces giving as a normal graph over Γ_0

$$\Gamma(t) = \{x + h(x, t)\nu_{\Gamma_0} \mid x \in \Gamma_0\}.$$

The function $h : \Gamma_0 \times [0, T] \rightarrow \mathbb{R}$ is known as a *height* function. The PDE in “the bulk” of the new domain will have coefficients determined by the change of variables, and the boundary conditions on this new domain are represented by a coupling between h and the fields along the fixed boundary [54, p. 33]). Having set up the problem in the fixed domain one can pose this as a non-linear evolution problem in some properly chosen Banach space. The problem can be written, under the right circumstances, as a non-linear perturbation of a linear problem where the linear operator is elliptic. Elliptic here meaning

in terms of a Fourier symbol, and operators of order higher than 2 are allowed (this, in particular makes it possible to treat problems with surface tension). With this approach it is possible to prove short time existence for smooth initial data, global existence near equilibrium, stability of equilibria, and more. This is done for a large class of problems, including two-phase Stefan problems [53], problems involving surface tension [36], and problems where the underlying fields are not necessarily scalar. For works making use of a Lagrangian approach to analyze free boundary problems, we mention work of Hadzic and Shkoller [39] where they obtain global stability for the classical Stefan problem, and Cheng, Coutand, and Shkoller [26] where the authors study the Hele-Shaw problem with surface tension.

We see there is a vast literature where free boundary problems are treated by putting the focus on the free boundary in one way or another. Either by assuming it is given by a global graph, and using contour integration to represent the free boundary condition as non-linear integro-differential equation (as done in the Muskat problem) or by representing it as the normal graph over a reference interface (e.g. using the Hanzawa transform) and pulling this back to a reference configuration to obtain a coupled system in a fixed domain. Finally, we note a result that is similar to our own in both spirit and results: using a blow up argument and a change of variables, two of the authors in [25] were able to deduce some regularity results of Hele-Shaw flow by invoking recent results for integro-differential equations.

The approach in this current paper shares features with both of these previous approaches— there is a nontrivial overlap as they all involve a “height function” h to represent the free boundary. However, the work we present here differs a bit in the sense that it is based on the structure of the underlying operators that is enforced by the comparison principle. The result is a description of (certain classes of) problems in terms of a single a scalar solving an equation with non-linear integro-differential operator reminiscent of the Dirichlet to Neumann map. This operator satisfies the global comparison principle so it can be studied via viscosity solutions methods for nonlocal equations and hence also opens up the possibility to subsequently apply non-divergence regularity results obtained in recent years.

5. THE FREE BOUNDARY OPERATOR

This section is dedicated to a study of something that we call a “free boundary” operator, which is defined in (1.5) and (1.6). Eventually we will prove many properties of I , including its Lipschitz nature as a function from special convex subsets of $C^{0,1}$ to $C^{\gamma'}$. The results of this section lead in two different, but related, directions with more or less common goals. The first is to be able to establish a min-max integro-differential representation for I and subsequently derive some basic properties of the linear operators that make up this min-max (e.g. no second order terms, and negative zero-order terms). The second direction is to use the Lipschitz nature of I , with some special functions, to derive a comparison theorem for sub and super solutions of the fractional parabolic problem, (1.2). Both of these inquiries are useful in their own right, and of course, they overlap at the stage of the comparison theorem. They will be developed in the sequels, Sections 7 and 8.

5.1. Setup. As above, we assume that F is uniformly elliptic and satisfies standard assumptions for existence and uniqueness of viscosity solutions, listed in section 2.2.

Given an $f \in C^{1,\gamma}(\mathbb{R}^d)$ with $\inf f > 0$ we assign U_f , the unique viscosity solution to

$$\begin{cases} F(D^2U_f, \nabla U_f) = 0 & \text{in } D_f, \\ U_f = 0 & \text{on } \Gamma_f, \\ U_f = 1 & \text{on } \Gamma_0. \end{cases} \quad (5.1)$$

We recall that for shorthand purposes, in this section, $\Gamma_f = \text{graph}(f)$ and $\Gamma_0 = \mathbb{R}^d \times \{0\}$ (Section 2.3 has this and other notations). We mention some sufficient conditions for the existence of such a U_f below. We note that, thanks to the boundary gradient estimates in [58, Remark 3.2], for some $\gamma' \in (0, \gamma]$, this U_f will enjoy a global in D_f estimate, $U_f \in C^{1,\gamma'}(\overline{D_f})$. Thus, we may define the

operator

$$I(f, x) := \partial_n U_f(x, f(x)), \quad (\text{equivalently, } I(f, x) = |\nabla U_f(x, f(x))|). \quad (5.2)$$

We see that for this γ' , we have

$$I : C^{1,\gamma}(\mathbb{R}^d) \cap \{f : \inf f > 0\} \rightarrow C^{\gamma'}(\mathbb{R}^d).$$

The study of this operator will be our chief concern. It turns out that there are more relaxed situations in which I is well defined, and we build these results below. They will use a generalization of semi-concavity that is well known in the field of optimal transport (usually referred to as c -convexity), and a special case is the next definition.

Definition 5.1 ($C^{1,\gamma}$ -semi-concave). *Let $\gamma \in (0, 1]$ and $m > 0$. A Lipschitz function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ will be said to be $C^{1,\gamma}$ -semi-concave with constant m if there is a real valued function, $r(y)$, such that*

$$f(x) = \inf_{y \in \mathbb{R}^d} \{r(y) + m|x - y|^{1+\gamma}\}.$$

A function f is said to be $C^{1,\gamma}$ -semi-convex with constant m if $(-f)$ is $C^{1,\gamma}$ -semi-concave with constant m .

We also need the notion of a function to be pointwise “ $C^{1,\gamma}$ ”.

Definition 5.2 (Pointwise $C^{1,\gamma}$). *As above, let $\gamma \in (0, 1]$ and $m > 0$ be fixed. We say that $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is pointwise m - $C^{1,\gamma}$ at x_0 , denoted $f \in m$ - $C^{1,\gamma}(x_0)$, if $\nabla f(x_0)$ exists, there exists $r > 0$, such that*

$$\begin{aligned} |f(x_0)| &\leq m, \quad |\nabla f(x_0)| \leq m, \\ \text{and } \forall x \in B_r(x_0), \quad &|f(x) - f(x_0) - \nabla f(x_0) \cdot (x - x_0)| \leq m|x - x_0|^{1+\gamma}. \end{aligned}$$

For $m, \delta > 0$ and $\gamma \in (0, 1)$, we consider the convex set of functions

$$\mathcal{K}(\gamma, \delta, m) := \{f \in C^{1,\gamma}(\mathbb{R}^d) \mid f(x) > \delta \ \forall x \in \mathbb{R}^d, \ \|f\|_{C^{1,\gamma}(\mathbb{R}^d)} < m\}, \quad (5.3)$$

as well as a respective convex sets of “semi-concave/convex” functions, which are larger,

$$\mathcal{K}^*(\gamma, \delta, m) := \{f \in C^{0,1}(\mathbb{R}^d) \mid f(x) > \delta \ \forall x \in \mathbb{R}^d, \ f \text{ is } C^{1,\gamma}\text{-semi-concave with constant } m\}, \quad (5.4)$$

$$\mathcal{K}_*(\gamma, \delta, m) := \{f \in C^{0,1}(\mathbb{R}^d) \mid f(x) > \delta \ \forall x \in \mathbb{R}^d, \ f \text{ is } C^{1,\gamma}\text{-semi-convex with constant } m\}. \quad (5.5)$$

It is clear that we have the inclusion

$$\mathcal{K}(\gamma, \delta, m) \subset \mathcal{K}^*(\gamma, \delta, m) \cap \mathcal{K}_*(\gamma, \delta, m).$$

Our goal, as noted above, is that we intend to use I to define (degenerate) parabolic equations, and to show a comparison theorem for viscosity solutions of these equations. This means that there are, more or less, three primary concerns:

- (1) For a fixed $x \in \mathbb{R}^d$, for which functions, f , is $I(f, x)$ classically defined? Certainly, we will require that there is a $\delta > 0$ so that $f \geq \delta$. Furthermore, it is not too hard to show, and we do below, that $I(f, x)$ is well defined if also for some $r > 0$, $f \in C^{0,1}(\mathbb{R}^d) \cap C^{1,\gamma}(B_r(x))$. However, we prefer to have a slightly weaker situation, and indeed, we are able to show that $I(f, x)$ is still well defined when $f \in C^{0,1}(\mathbb{R}^d) \cap (m$ - $C^{1,\gamma}(x))$.
- (2) Over which collection of functions does I enjoy the GCP? (The GCP appears in Definition 2.2.)
- (3) Over which set is the mapping I Lipschitz? We will show that I is locally Lipschitz on the convex sets given by $\mathcal{K}^*(\gamma, \delta, m)$ above, and that the Lipschitz norm grows as either δ decreases or m increases. (As a corollary, I will be locally Lipschitz on $\mathcal{K}(\gamma, \delta, m)$.)

5.2. Basic properties. We have defined $I(f, x)$ in (5.2) for functions which are globally of class $C^{1,\gamma}$ for some $\gamma \in (0, 1)$. We now carefully check the existence and uniqueness for U_f as well as the well-posedness of $I(f, x)$ in some less restrictive situations. Eventually, we show that I is well defined whenever f is either Lipschitz and locally $C^{1,\gamma}$ in a neighborhood of the point of evaluation, or when f is Lipschitz and $C^{1,\gamma}$ -semi-concave.

Proposition 5.3. *If $f \in C^{0,1}(\mathbb{R}^d)$ and $f \geq \delta$, then there exists a unique $U_f \in C(\overline{D_f})$ that is the viscosity solution of (5.1) and continuously attains its boundary values.*

Proof of Proposition 5.3. This proposition is more or less standard for viscosity solutions, but we were unable to find a standard reference that contained the particular situation stated above. Thus, we have included some of the main details.

So long as we can construct a barrier for U_f to force it to attain its boundary values on Γ_f continuously, we will have the existence and uniqueness from the results by, e.g. Ishii [42]. This is a consequence of the fact that the Perron Method will produce a solution in the interior, and so we may assume that U_f is already defined in D_f . We just focus on the boundary values.

We first note that the fact that $\Gamma_0 = \mathbb{R}^d \times \{0\}$ is flat (hence C^2), the existence of lower and upper barriers attaining the value 1 on Γ_0 is standard. We instead focus our attention on barriers at the upper boundary, Γ_f .

The Lipschitz nature of f gives D_f an exterior cone condition, and so it is possible to construct an upper barrier for (5.1) (we note that the constant, zero, function serves as a lower barrier). Let $X_0 = (x_0, f(x_0)) \in \Gamma_f$ be fixed. We will construct a function ψ_{X_0} that serves as an upper barrier for U_f at X_0 . Let ϕ be a continuous function such that

$$\phi(X_0) = 0, \quad \phi \geq 0, \quad \phi(Y) \geq 1 \text{ for } Y \in (\partial Q_\delta(X_0) \cap \overline{D_f}),$$

where we use $Q_\delta(X_0)$ to be the cube of side length, δ , centered at X_0 . We note that since D_f enjoys the exterior cone condition, then so does $Q_\delta(X_0) \cap D_f$. Thus, by [10, Corollary 3.10], there exists a unique solution, $\psi_{X_0} \in C^{2,\gamma}(Q_\delta(X_0) \cap D_f) \cap C(\overline{Q_\delta(X_0) \cap D_f})$, to the extremal equation,

$$\begin{cases} \mathcal{M}^+(D^2\psi_{X_0}) + |\nabla\psi_{X_0}| = 0 & \text{in } Q_\delta(X_0) \cap D_f \\ \psi_{X_0} = \phi & \text{on } \partial(Q_\delta(X_0) \cap D_f). \end{cases}$$

From the definition of uniform ellipticity, it follows that this ψ_{X_0} is a viscosity supersolution of $F(D^2U, \nabla U) = 0$ in the domain $Q_\delta(X_0) \cap D_f$.

Because we know that $U_f \leq 1$ in $\overline{D_f}$ and $U_f = 0$ on Γ_f , we see that $U_f \leq \psi$ on $\partial(Q_\delta(X_0) \cap D_f)$. Thus, by comparison, we see that

$$0 \leq U_f(X_0) \leq \psi(X_0) = 0.$$

Hence, $U_f(X_0) = 0$. Since X_0 was generic, we conclude that such a U_f exists and continuously attains $U_f = 0$ on Γ_f . □

Proposition 5.4. *If $f, g \in C^{0,1}(\mathbb{R}^d)$, $f \geq \delta$, $g \geq \delta$, and $f \leq g$ in \mathbb{R}^d , then for U_f, U_g solving (5.1),*

$$U_f \leq U_g \text{ in } D_f.$$

Proof of Proposition 5.4. First, we note that $f \leq g$ implies that $D_f \subset D_g$. Furthermore, since $f = 0$ on Γ_f and since $g \geq 0$ in D_f , we see that U_g is a supersolution of (5.1) for f . Hence, $U_f \leq U_g$ is a direct application of the comparison theorem for elliptic equations in [42]. □

Proposition 5.5. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a bounded, globally Lipschitz function such that $f \geq \delta$ in \mathbb{R}^d for some $\delta > 0$, and let $x_0 \in \mathbb{R}^d$. If f is $C^{1,\gamma}(B_r(x_0))$, for some fixed $r \in (0, 1)$ then $I(f, x_0)$ is well*

defined. Moreover, there is a constant C , depending on d, Λ, λ and the $C^{1,\gamma}(B_r(x_0))$ norm of f , such that

$$|I(f, x_0)| \leq Cr^{-1}.$$

Proof. The assumptions on f have already been shown in Proposition 5.3 to give the existence and uniqueness of U_f . In particular, U_f satisfies in the viscosity sense

$$F(D^2U_f, \nabla U_f) = 0 \text{ in } D_f \cap B_r^{d+1}((x_0, f(x_0))).$$

Since ∂D_f is of class $C^{1,\gamma}$, from [58, Remark 3.2] it follows that U_f is of class $C^{1,\gamma'}$ in the smaller domain $D_f \cap B_{\frac{r}{2}}^{d+1}((x_0, f(x_0)))$, with the estimate

$$\|\nabla U_f\|_{L^\infty(D_f \cap B_{\frac{r}{2}}^{d+1}((x_0, f(x_0))))} \leq Cr^{-1} \|U_f\|_{L^\infty(D_f)} \leq Cr^{-1}.$$

□

Lemma 5.6. *Assume that $x_0 \in \mathbb{R}^d$, $\delta > 0$, and $r > 0$ are fixed. The map, I , has the global comparison property based at x_0 (Definition 2.2) for functions, f, g that satisfy*

$$f, g \in \left(C^{0,1}(\mathbb{R}^d) \cap C^{1,\gamma}(B_r(x_0)) \cap \{h : \mathbb{R}^d \rightarrow \mathbb{R} : h \geq \delta\} \right).$$

Consequently I also enjoys the GCP for functions in $\mathcal{K}(\gamma, \delta) = \bigcup_m \mathcal{K}(\gamma, \delta, m)$.

Proof. Let $f, g \in C^{0,1}(\mathbb{R}^d) \cap C^{1,\gamma}(B_r(x_0))$ and $x_0 \in \mathbb{R}^d$ be such that

$$f(x) \leq g(x) \quad \forall x \in \mathbb{R}^d, \quad f(x_0) = g(x_0).$$

It is immediate that

$$D_f \subset D_g, \quad \text{and} \quad X_0 := (x_0, f(x_0)) = (x_0, g(x_0)) \in \Gamma_f \cap \Gamma_g.$$

Now, since the boundary values of U_g are nonnegative, the maximum principle shows that

$$U_g \geq 0 \text{ in } D_g.$$

In particular, since $\Gamma_f \subset D_f$, we have $U_g \geq 0$ on Γ_f . Thus, $U_g \geq U_f$ on the boundary of D_f . Then, from the comparison principle, it follows that

$$U_g \geq U_f \quad \text{in } D_f.$$

As both functions vanish at X_0 , we conclude that at X_0

$$|\nabla U_g| \geq |\nabla U_f|,$$

in other words, $I(f, x_0) \leq I(g, x_0)$, as we wanted.

□

The next proposition says the translation invariance of our setup implies, as one would expect, that the operator I itself is translation invariant.

Proposition 5.7. *For each $\gamma \in (0, \infty)$ and $\delta > 0$, $I : \mathcal{K}(\gamma, \delta) \rightarrow C^0$ is translation invariant; i.e. for f given, then for any $h \in \mathbb{R}^d$ that is fixed, we have*

$$\forall x \in \mathbb{R}^d, \quad [\tau_h I(f)](x) = I(\tau_h f, x).$$

Proof. For $h \in \mathbb{R}^d$, fixed, let us extend the translation operator, τ_h , to act on functions on \mathbb{R}^{d+1} as

$$W : \mathbb{R}^{d+1} \rightarrow \mathbb{R}, \quad (x, x') \in \mathbb{R}^{d+1}, \quad \tau_h W(x, x') := W(x + h, x').$$

The proposition will follow immediately from the observation

$$\tau_h U_f = U_{\tau_h f},$$

which itself is a consequence of the fact that the operator, F , and the lower boundary of D_f are translation invariant. Indeed, if we define $W = \tau_h U_f$, we see that

$$F(D^2W(x, x'), \nabla W(x, x')) = 0, \quad \text{whenever } (x + h, x') \in D_f;$$

and furthermore,

$$W(x, f(x + h)) = U_f(x + h, f(x + h)) = 0 \quad \text{and} \quad W(x, 0) = U_f(x, 0) = 0.$$

Hence W solves (5.1) in the domain $D_{\tau_h f}$. By the uniqueness of solutions of (5.1), we see that we have

$$\tau_h U_f = W = U_{\tau_h f}.$$

Since the operator ∂_n commutes with τ_h , we conclude that

$$\tau_h(\partial_n U_f)(x) = \partial_n(\tau_h U_f)(x) = \partial_n U_{\tau_h f}(x),$$

whence

$$\tau_h[I(f)](x) = I(\tau_h f, x).$$

□

The next two propositions say respectively that U_f is monotone increasing in f , and that if f and g are close in L^∞ , then U_f and U_g are also close in L^∞ in their common domain.

Proposition 5.8. *There is a constant $C = C(d, \lambda, \Lambda, \delta, \gamma, m)$ such that for all $f \in \mathcal{K}^*(\gamma, \delta, m) \cap C^{0,1}(\mathbb{R}^d)$,*

$$\forall (x, x_{d+1}) \in D_f, \quad U_f(x, x_{d+1}) \leq C(f(x) - x_{d+1}).$$

Furthermore,

$$\nabla U_f \in L^\infty(D_f) \quad \text{and} \quad 0 \leq \partial_n U_f(x, f(x)) \leq C, \quad \text{whenever } \partial_n U_f \text{ exists.}$$

Proof of Proposition 5.8. Assume that x_0 is fixed. Since f is $C^{1,\gamma}$ -semi-concave, there exists a $\psi \in C^{1,\gamma}(\mathbb{R}^d)$ so that $f \leq \psi$ and $f(x_0) = \psi(x_0)$, and $\|\psi\|_{C^{1,\gamma}}$ depends only on δ and m . Thus, $D_f \subset D_\psi$. Furthermore, we know already, from [58, Remark 3.2] that there is some γ' so that $U_\psi \in C^{1,\gamma'}(\overline{D_\psi})$. In particular, U_ψ is globally Lipschitz in $\overline{D_\psi}$. Finally, because of the ordering of $f \leq \psi$ and $f(x_0) = \psi(x_0)$, we see that both $U_f \leq U_\psi$ (from Proposition 5.4) and $0 \leq \partial_n U_f(x_0, f(x_0)) \leq \partial_n U_\psi(x_0, \psi(x_0))$. The $C^{1,\gamma}$ nature of ψ that depends only on δ and m means that the Lipschitz norm of U_ψ depends only on $d, \lambda, \Lambda, \delta, \gamma$, and m . Since this Lipschitz property of U_ψ implies the result of the lemma with U_f replaced by U_ψ , we conclude by the previously noted ordering of $U_f \leq U_\psi$, that the outcome of the lemma is valid for U_f as well. □

Proposition 5.9. *There is a constant $C = C(d, \lambda, \Lambda, \delta, \gamma, m)$ such that if $f, g \in \mathcal{K}^*(\gamma, \delta, m)$, then*

$$\|U_f - U_g\|_{L^\infty(D_f \cap D_g)} \leq C\|f - g\|_{L^\infty(\mathbb{R}^d)}.$$

Proof. Let us show there is a $C = C(\lambda, \Lambda, \delta, \gamma, m)$ such that if $f \in \mathcal{K}^*(\delta, \gamma, m)$ and $s > 0$, then

$$U_{f+s} \leq U_f + Cs \quad \text{in } D_f. \tag{5.6}$$

Indeed, by Proposition 5.8

$$U_{f+s}(x, x_{d+1}) \leq C(f(x) + s - x_{d+1})_+ \quad \forall (x, x_{d+1}) \in D_{f+s},$$

In particular, $U_{f+s}(x, f(x)) \leq Cs$ for every $x \in \mathbb{R}^d$, which is the same as

$$U_{f+s} \leq Cs \quad \text{on } \Gamma_f.$$

Then, if $\tilde{U} := U_f + Cs$ with this same C , we have

$$U_{f+s} \leq \tilde{U} \quad \text{on } \partial D_f.$$

Furthermore, \tilde{U} is a viscosity solution of $F(D^2U, \nabla U) = 0$ in D_f , so by the comparison principle, $U_{f+s} \leq \tilde{U}$ everywhere in D_f , and (5.6) is proved. Now, given a second function $g \in \mathcal{K}^*(\delta, \gamma, m)$, let $s = \|f - g\|_\infty$, so that

$$g \leq f + s.$$

Then, by Proposition 5.4 we have $U_g \leq U_{f+s}$ in D_g . Applying (5.6), it follows that

$$U_g \leq U_f + Cs \text{ in } D_g.$$

Arguing in the exact same manner but reversing the roles of f and g , we conclude that (with the same constant C as before)

$$U_f \leq U_g + Cs \text{ in } D_f,$$

and this proves the proposition. \square

In the next proposition, we state (with an abbreviated proof) a basic estimate for a one parameter family of barrier functions $\{H_s\}_{s>0}$.

Proposition 5.10. *Fix $w \in \mathcal{K}(\gamma, \delta, m)$ and $s \in (0, \infty)$. Let $H_s : D_w \rightarrow \mathbb{R}$ be the unique viscosity solution of*

$$\begin{cases} F(D^2H_s, \nabla H_s) = 0 & \text{in } D_w, \\ H_s = 0 & \text{on } \Gamma_w, \\ H_s = s & \text{on } \Gamma_0. \end{cases}$$

Then, for some universal $\gamma' \in (0, \gamma]$ and a constant $C = C(d, \lambda, \Lambda, \gamma, \delta, m)$, we have for $X_0 \in \Gamma_w$ and $X \in \overline{D_w}$

$$\begin{aligned} |H_s(X) - (\nabla H_s(X_0), X - X_0)| &\leq Cs|X - X_0|^{1+\gamma'}, \\ C^{-1}s &\leq |\nabla H_s(X_0)| \leq Cs. \end{aligned}$$

Moreover, there is a constant C such that

$$|H_{s_1}(X) - H_{s_2}(X)| \leq C|s_1 - s_2|d(X, \Gamma_w).$$

Proof. We just provide a small sketch of the details. The first two claims are immediate from the $C^{1, \gamma'}$ regularity of solutions in the domain, D_w , also using lower and upper barriers to bound the gradient along Γ_w for the second claim. The lower bound on $|\nabla H_s|$ uses the Hopf principle for fully nonlinear equations.

The third assertion follows from the fact that, if we assume that $s_1 \geq s_2$, then we have $0 \leq H_{s_1} - H_{s_2} \leq \psi_{up}$, where ψ_{up} is a barrier function that solves

$$\mathcal{M}^+(D^2\psi_{up}) + \Lambda|\nabla\psi_{up}| = 0 \text{ with } \psi_{up} = s_1 - s_2 \text{ on } \Gamma_0, \text{ and } \psi_{up} = 0 \text{ on } \Gamma_w.$$

Furthermore, standard regularity theory, e.g. [58], shows that $\psi_{up} \in C^{0,1}(\overline{D_w})$, with

$$0 \leq \psi_{up} \leq s_1 - s_2, \text{ and } |\nabla\psi_{up}| \leq C\|\psi_{up}\|_{L^\infty}.$$

The claim follows by using the barrier up to the boundary at Γ_w . \square

The following Lemma follows an argument about the behavior of harmonic functions near regular points of their boundary. Here we adapt the details from, e.g. [11, Lemma 11.17].

Lemma 5.11. *Assume f is such that $\inf f > 0$ and f is differentiable at x_0 , and furthermore that f satisfies for some $\gamma \in (0, 1)$ and $C > 0$*

$$|f(x) - f(x_0) - (\nabla f(x_0), x - x_0)| \leq C|x - x_0|^{1+\gamma}.$$

Then, the function U_f is differentiable at $X_0 = (x_0, f(x_0))$, in the sense that for some $\alpha > 0$

$$U_f(X) = \alpha(n(X_0), X - X_0) + o(|X - X_0|),$$

as $X \rightarrow X_0$ non-tangentially in D_f . Here $n(X_0)$ denotes the inner normal to D_f at X_0 .

Proof. Let $\delta = \inf f > 0$. Consider the function defined as

$$w(x) = \rho_0(f(x_0) + (\nabla f(x_0), x - x_0) - C|x - x_0|^{1+\gamma}),$$

for x close to x_0 , where ρ_0 denotes a smooth, monotone function of one variable such that


$$\begin{aligned} \rho_0(t) &= t \text{ for } t \geq \delta, \\ \rho_0(t) &\leq \delta \text{ for } t \leq \delta, \\ \rho_0(t) &= \delta/2 \text{ for } t \leq \delta/2. \end{aligned}$$

Then, it is clear that $w \in \mathcal{K}(\gamma, \delta/2, m')$ for some $m' = m'(\gamma, \delta, C)$. Moreover, from the assumption on f (that $f(x) \geq f(x_0) + (\nabla f(x_0), x - x_0) - C|x - x_0|^{1+\gamma}$), we see that $w(x_0) = f(x_0)$ and $w \leq f$. Thus,

$$D_w \subset D_f \text{ and } X_0 \in \partial D_w \cap D_f.$$

Let $H_s(X)$ be as in Proposition 5.10, for the domain given by D_w . For $k \in \mathbb{N}$ sufficiently large (so that $2^{-k} \leq \delta$) we define

$$\alpha_k = \sup\{s \mid U_f \geq H_s \text{ in } B_{2^{-k}}(X_0) \cap D_w\}.$$



It is clear that the sequence α_k is non-decreasing with respect to k . Therefore, there is some real number α_* such that $\alpha_* = \lim \alpha_k$ or else U_f has superlinear growth at X_0 . The latter option is impossible, since U_f is Lipschitz at X_0 ; seen by using a $C^{1,\gamma}$ function, ψ , that touches f from above at x_0 and arguing as in Proposition 5.8.

Let $\{X_m\}_m$ be any sequence converging to X_0 in D_w , and let us write

$$U_f(X_m) - H_{\alpha_*}(X_m) = U_f(X_m) - H_{\alpha_k}(X_m) + (H_{\alpha_k}(X_m) - H_{\alpha_*}(X_m)),$$

for any k , and let k_m be defined by the inequalities

$$2^{-k_m-2} \leq r_m < 2^{-k_m-1}, \quad r_m := |X_m - X_0|.$$

Then, for every m we have

$$U_f(X_m) - H_{\alpha_*}(X_m) \geq H_{\alpha_{k_m}}(X_m) - H_{\alpha_*}(X_m).$$

On the other hand, thanks to Proposition 5.10, there is $C > 0$ such that

$$|H_{\alpha_{k_m}}(X_m) - H_{\alpha_*}(X_m)| \leq C|\alpha_{k_m} - \alpha_*| \cdot |X_m - X_0|.$$

We know that $\alpha_* - \alpha_{k_m} \rightarrow 0$ as $m \rightarrow \infty$, so the right hand side is $o(|X_m - X_0|)$. Since the sequence X_m was arbitrary, we have proved that

$$U_f(X) \geq H_{\alpha_*}(X) + o(|X - X_0|), \quad X \in D_w.$$

Thus, using Proposition 5.10, we can establish a further characterization of α_* , which is

$$\alpha_* = \sup \{s \mid U_f(X) \geq H_s(X) + o(|X - X_0|), \text{ as } X \rightarrow X_0 \text{ non-tangentially}\}.$$

Let us show that in fact

$$U_f(X) = H_{\alpha_*}(X) + o(|X - X_0|), \text{ as } X \rightarrow X_0 \text{ non-tangentially.} \quad (5.7)$$

We argue by contradiction. If (5.7) does not hold, there is a sequence $X_m \in D_w$ converging non-tangentially to X_0 along which $U_f - H_{\alpha_*}$ is larger than a quantity comparable to the distance to X_0 . In other words, there is some $\theta > 0$ such that

$$X_m \rightarrow X_0, \quad |X_m - X_0| \geq \theta |X'_m - X_0|, \quad X_m \in D_w,$$

(X'_m is the projection of X_m onto the tangent hyperplane to Γ_w at X_0) and at the same time

$$U_f(X_m) \geq H_{\alpha_*}(X_m) + \beta |X_m - X_0|,$$

for some $\beta > 0$ independent of m . To take advantage of this, let us define for $k \in \mathbb{N}$ the function

$$W_k(X) := U_f(X) - H_{\alpha_k}(X), \quad \text{for } X \in D_w \setminus D_{w-r_0}.$$


Since $\alpha_k \leq \alpha_*$ for all k , we have

$$W_k(X_m) \geq \beta |X_m - X_0| \text{ and } W_k \geq 0 \text{ in } B_{2^{-k}}(X_0) \cap D_w;$$

furthermore, W_k , satisfies (in the viscosity sense)

$$\mathcal{M}^+(D^2W_k) + \Lambda |\nabla W_k| \geq -C_1,$$

$$\mathcal{M}^-(D^2W_k) - \Lambda |\nabla W_k| \leq C_1.$$



Here we make use of the non-tangential convergence of the sequence: there is a $c_1 > 0$ such that

$$B_{c_1 r_m}(X_m) \subset B_{2^{-k_m}}(X_0).$$

Applying the Harnack inequality (e.g. a rescaled statement of [16, Theorem 4.3] or [60, Theorem 4.18], which includes the gradient term) it follows that (for some $\beta_0 > 0$)

$$W_{k_m}(X) \geq \beta_0 r_m \text{ in } B_{\frac{1}{2}c_1 r_m}(X_m) \cap D_w.$$

This together with $W_{k_m} \geq 0$ in $B_{\frac{1}{2}c_1 r_m}(X_m) \cap D_w$, guarantees (via a standard barrier argument) that

$$W_{k_m}(X) \geq c_2 \beta_0 d(x, \partial D_w) \text{ in } B_{2^{-k_m}}(X_0) \cap D_w,$$

for some constant $c_2 = c_2(d, \lambda, \Lambda, \gamma, C)$. In terms of U_f , this says that

$$U_f(X) \geq H_{\alpha_{k_m}}(X) + c_2\beta_0 d(X, \partial D_w) \text{ in } B_{2^{-k_m}}(X_0) \cap D_w.$$

On the other hand, by Proposition 5.10,

$$H_\alpha(X) - H_{\alpha_{k_m}}(X) \leq |\alpha - \alpha_{k_m}| d(X, \partial D_w).$$

Therefore, choosing some $\alpha > \alpha_{k_m}$ sufficiently close to α_{k_m} (i.e. $|\alpha - \alpha_{k_m}| \leq c_2\beta_0$) and $\alpha > \alpha_*$, we have

$$U_f(X) \geq H_\alpha(X) \text{ in } B_{2^{-k_m}}(X_0) \cap D_w,$$

which is impossible because α_* is the supremum over all such α . Therefore (5.7) holds and the Lemma is proved. \square

The following is a useful corollary of Lemma 5.11 that we state without a proof.

Corollary 5.12. *The value, $I(f, x_0)$, is well defined whenever f is pointwise m - $C^{1,\gamma}(x_0)$.*

5.3. The Lipschitz property. In this section, we show that I is a Lipschitz mapping on $\mathcal{K}^*(\gamma, \delta, m)$, which are convex subsets of $C^{0,1}$. This Lipschitz property will have two very important consequences: a representation of I via integro-differential operators; and a comparison theorem within $\mathcal{K}^*(\gamma, \delta, m) \cap \mathcal{K}_*(\gamma, \delta, m)$ and hence for general viscosity solutions. The main result is the following theorem and its corollary.

Theorem 5.13. *There is a constant $C = C(d, \lambda, \Lambda, \delta, \gamma, m)$ such that if f, g satisfy*

$$f, g \in \mathcal{K}^*(\delta, \gamma, m) \text{ and } f - g \in C^{1,\gamma}(\mathbb{R}^d),$$

then for any $x \in \mathbb{R}^d$ at which $I(f, x)$ and $I(g, x)$ are both defined classically, we have

$$I(g, x) - I(f, x) \leq C \|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

Corollary 5.14. *If f and g satisfy $f \in \mathcal{K}_*(\delta, \gamma, m)$ and $g \in \mathcal{K}^*(\delta, \gamma, m)$ as well as $\phi \in C^{1,\gamma}(\mathbb{R}^d)$ with $g + \phi \in \mathcal{K}^*(\delta, \gamma, m)$, and all are such that*

$$(f - g) - \phi \text{ has a non-negative global maximum at } x_0 \in \mathbb{R}^d,$$

then, with $C = C(d, \lambda, \Lambda, \delta, \gamma, m)$, we have

$$I(f, x_0) - I(g, x_0) \leq C \|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

Similarly, as above, if $g - \phi \in \mathcal{K}^(\delta, \gamma, m)$ and*

$$(g - f) - \phi \text{ has a non-positive global minimum at } x_0 \in \mathbb{R}^d,$$

then

$$I(f, x_0) - I(g, x_0) \leq C \|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

The proofs of Theorem 5.13 and Corollary 5.14 will require some preliminary propositions and lemmas. We will finish the proofs of the Theorem and Corollary at the end of this section.

In estimating $I(g, x) - I(f, x)$ at a given x , it will be necessary to reduce matters to the situation where $f = g$ at x . This is the purpose of the next result.

Lemma 5.15. *There is a $C = C(d, \lambda, \Lambda, \delta, \gamma, m)$ such that if $s \geq 0$ and f is a function such that $f, f + s \in \mathcal{K}^*(\delta, \gamma, m)$, then for all x such that $I(f, x)$ is well defined,*

$$I(f + s, x) \leq I(f, x) \leq I(f + s, x) + Cs, \quad \forall x \in \mathbb{R}^d.$$

Proof. Let us compare U_f to a vertical shift of U_{f+s} . We use the coordinates, $(x, x_{d+1}) \in \mathbb{R}^d \times [0, \infty)$. Define $\tilde{U} : D_f \rightarrow \mathbb{R}$ by

$$\tilde{U}(x, x_{d+1}) := U_{f+s}(x, x_{d+1} + s).$$

Observe that $\tilde{U} = U = 0$ on Γ_f while $1 - Cs \leq \tilde{U} \leq 1$ on Γ_0 . Furthermore, the translation invariance of (5.1) ensures that \tilde{U} also solves (5.1) in a translated domain. Thus, by the comparison principle

$$\tilde{U} \leq U_f \text{ and } U_f - \tilde{U} \leq Cs \text{ in } D_f.$$

The first inequality implies that

$$|\nabla \tilde{U}| \leq |\nabla U_f| \text{ on } \Gamma_f,$$

while the second inequality, together with the same upper gradient estimate at the boundary used in Proposition 5.8 (which uses that $f \in \mathcal{K}^*$), implies that

$$|\nabla(U_f - \tilde{U})| \leq Cs \text{ on } \Gamma_f.$$

In conclusion, we have

$$|\nabla \tilde{U}| \leq |\nabla U_f| \leq |\nabla \tilde{U}| + Cs \text{ on } \Gamma_f.$$

Noting that $\nabla \tilde{U}(x, f(x)) = \nabla U_{f+s}(x, f(x) + s)$, we conclude that

$$I(f + s, x) \leq I(f, x) \leq I(f + s, x) + Cs.$$

□

As we saw in Proposition 5.9, the function U_f depends in a Lipschitz manner on the function f , as long as f lies in $\mathcal{K}^*(\delta, \gamma, m)$. The next Lemma will produce a barrier that will allow us to translate pointwise control of U_f in D_f into pointwise control of ∇U_f along Γ_f , this will be the key step in the proof of Theorem 5.13.

Lemma 5.16. *Fix $w \in \mathcal{K}^*(\delta, \gamma, m) \cap (m-C^{1,\gamma}(x_0))$, $X_0 \in \Gamma_w$, and $r_0 \in (0, \delta/2)$. Let $W : D_w \cap B_{r_0}(X_0) \rightarrow \mathbb{R}$ be a continuous function such that in the viscosity sense,*

$$\mathcal{M}^+(D^2W) + \Lambda |\nabla W| \geq -C_1 \text{ in } D_w \cap B_{r_0}(X_0),$$

$$\mathcal{M}^-(D^2W) - \Lambda |\nabla W| \leq C_1 \text{ in } D_w \cap B_{r_0}(X_0),$$

$$|W| \leq C_1 |X - X_0|^{1+\gamma} \text{ on } \partial(D_w \cap B_{r_0}(X_0)).$$

Then, with a constant $C = C(d, \lambda, \Lambda, \gamma, \delta, m, r_0)$ we have

$$|\nabla W(X_0)| \leq C \cdot C_1.$$

Proof. Consider the $C^{1,\gamma}(\mathbb{R}^d)$ function given by

$$\hat{w}^+(x) := \eta_\delta(w(x_0) + Dw(x_0) \cdot (x - x_0) + m|x - x_0|^{1+\gamma}),$$

where $\eta_\delta(t) := \delta\eta(t/\delta)$ and $\eta : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth non-decreasing function such that

$$\eta(t) = t \text{ for } t \geq 1, \quad \eta(t) = 1/2 \text{ for } t \leq 1/2.$$

This guarantees that $\hat{w}^+(x) \geq \delta/2$ and $\hat{w}^+(x) \geq w(x)$ for all $x \in \mathbb{R}^d$ with equality for $x = x_0$. In particular, note that $\Gamma_w \subset D_{\hat{w}^+}$. Then, let \hat{W}^+ be the unique viscosity solution of

$$\begin{cases} \mathcal{M}^+(\hat{W}^+) + \Lambda |\nabla \hat{W}^+| = -1 & \text{in } D_{\hat{w}^+} \cap B_{r_0}(X_0), \\ \hat{W}^+ = |X - X_0|^{1+\gamma} & \text{on } \partial(D_{\hat{w}^+} \cap B_{r_0}(X_0)). \end{cases}$$

Using $w \in m-C^{1,\gamma}(x_0)$ and a standard barrier argument we have that for c_0, r_1 small (depending on λ, Λ, m, r_0 , and γ) we have

$$\hat{W}^+(X) \geq c_0 |X - X_0|^{1+\gamma} \text{ on } \partial(D_w \cap B_{r_1}(X_0)). \quad (5.8)$$

From [58, Remark 3.2] we know that $\hat{W}^+ \in C^{1,\gamma'}(\overline{D_{\hat{w}^+} \cap B_{r_0/2}(X_0)})$; and in particular, that

$$|\nabla \hat{W}^+(X_0)| \leq C.$$

From (5.8) it follows that $W \leq c_0^{-1} C_1 \hat{W}^+$ on $\partial(D_w \cap B_{r_1}(X_0))$ and since without loss of generality we may assume $c_0 \in (0, 1)$

$$\Lambda \left| \nabla(C_1 \hat{W}^+) \right| + \mathcal{M}^+(D^2(C_1 \hat{W}^+)) = -c_0^{-1} C_1 \leq -C_1 \leq \mathcal{M}^+(D^2 W) + \Lambda |\nabla W| \text{ in } D_w \cap B_{r_1}(X_0),$$

in the viscosity sense. Thus, the comparison principle says that

$$W \leq \hat{W}^+ \text{ in } D_w \cap B_{r_1}(X_0).$$

We can repeat this argument from below, using $\hat{W}^- := -\hat{W}^+$ as a barrier, so $|W| \leq \hat{W}^+$ in $D_w \cap B_{r_1}(X_0)$. Since we also have $W(X_0) = \hat{W}^\pm(X_0) = 0$, it follows that $|\nabla W(X_0)| \leq |\nabla \hat{W}^+(X_0)|$, and from the bound above it follows that

$$|\nabla W(X_0)| \leq (c_0^{-1} C) \cdot C_1,$$

which proves the lemma.

(Proof of (5.8): since the domain $D_{\hat{w}^+}$ is of class $C^{1,\gamma}$ we can apply the Hopf Lemma at $X_0 = (x_0, f(x_0))$ as well as global C^{1,γ^-} estimates for \hat{W}^+ to conclude that

$$-\partial_y \hat{W}^+ \geq 0 \text{ in } D_{\hat{w}^+} \cap B_{r_1}(X_0)$$

for some $r_1 \in (0, r_0/2)$ is sufficiently small (depending only on λ, Λ, m, r_0 , and γ). Then, for every $X' \in \Gamma_{\hat{w}^+} \cap B_{r_1}(X_0)$ and $X \in D_{\hat{w}^+} \cap B_{r_1}(X_0)$ of the form $X := X' - se_{d+1}$ ($s > 0$) we have

$$\hat{W}^+(X) \geq \hat{W}^+(X') \geq |X' - X_0|^{1+\gamma}.$$

This means that if $X = (x, y) \in D_{\hat{w}^+} \cap B_{r_1}(X_0)$ then

$$\hat{W}^+(X) \geq (|x - x_0|^2 + (\hat{w}^+(x) - \hat{w}^+(x_0))^2)^{(1+\gamma)/2}.$$

At the same time if $X \in \Gamma_w$ then $y = w(x)$ we have (recall $\hat{w}^+(x_0) = w(x_0)$)

$$\begin{aligned} |y - \hat{w}^+(x_0)| &= |w(x) - \hat{w}^+(x_0)| \\ &\leq |w(x) - \hat{w}^+(x)| + |\hat{w}^+(x) - \hat{w}^+(x_0)| \\ &\leq m|x - x_0|^{1+\gamma} + |\hat{w}^+(x) - \hat{w}^+(x_0)|. \end{aligned}$$

from where it follows that

$$|X - X_0| \leq C(m)|X' - X_0|.$$

This shows that for some $C(m) > 0$ we have

$$\hat{W}^+(X) \geq C(m)^{-1}|X - X_0|^{1+\gamma} \text{ on } \Gamma_w \cap B_{r_1}(X_0).$$

Using once again that $-\partial_y \hat{W}^+ \geq 0$ in $\hat{w}^+ \cap B_{r_1}(X_0)$ and Harnack inequality it is not difficult to show that $\hat{W}^+ \geq C$ on $D_w \cap \partial B_{r_1}(X_0)$, where $C = C(\lambda, \Lambda, d, m, \gamma, r_0)$. This completes the proof (5.8). \square

The next two lemmas are concerned with how much the operator, I , can deviate when it is evaluated on two domains that touch at a point with C^1 contact. Originally, it was thought that Lemma 5.17 (below) would be sufficient, but this subsequently turned out not to be the case. We have decided to leave it here because we believe it could be useful for future investigations.

Lemma 5.17. *Let f, g be such that $f, g \in \mathcal{K}^*(\delta, \gamma, m) \cap (m\text{-}C^{1,\gamma}(x_0))$, and $f - g \in C^{1,\gamma}(\mathbb{R}^d)$. Assume that $x_0 \in \mathbb{R}^d$ be a point such that $f(x_0) = g(x_0)$ and $\nabla f(x_0) = \nabla g(x_0)$. In this case, there exists a $C = C(d, \lambda, \Lambda, \delta, \gamma, m)$ such that*

$$|I(f, x_0) - I(g, x_0)| \leq C \|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

Proof of Lemma 5.17. The assumptions of the lemma allow us to assert that

$$\forall x \in \mathbb{R}^d, \quad f(x) \geq g(x) - \|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)} |x - x_0|^{1+\gamma},$$

(Recall, we use the convention that $X \in \mathbb{R}^{d+1}$ whereas $x \in \mathbb{R}^d$.) We will modify this function on the right hand side of the inequality to define a function w ,

$$w(x) := \eta_\delta(g(x) - \|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)} |x - x_0|^{1+\gamma}),$$

here $\eta_\delta(t) := \delta\eta(t/\delta)$ where $\eta : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth, non-decreasing function such that

$$\eta(t) = t \text{ for } t \geq 1, \quad \eta(t) = 1/2 \text{ for } t \leq 1/2.$$

This guarantees that $w(x) \geq \delta/2$ and that $f(x)$ and $g(x)$ are both greater than $w(x)$ for all $x \in \mathbb{R}^d$. For notational convenience, let us call

$$X_0 = (x_0, f(x_0)) = (x_0, g(x_0)) \in \Gamma_f \cap \Gamma_g.$$

Thus, by this choice of w , we have,

$$D_w \subset D_f \quad \text{and} \quad D_w \subset D_g.$$

Recalling Proposition 5.9, it follows that

$$\|U_g - U_f\|_{L^\infty(D_w)} \leq C\|f - g\|_{L^\infty(\mathbb{R}^d)}.$$

Thanks to the globally Lipschitz nature of U_f and U_g , and the definition of w , we have in fact the following pointwise estimate

$$|U_g(X) - U_f(X)| \leq C\|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)} |X - X_0|^{1+\gamma}, \quad \text{for } X \in \Gamma_w \cap B_r(X_0).$$

(This follows immediately once we keep in mind the zero boundary values of U_f, U_g on Γ_f, Γ_g , and that both top boundaries separate from Γ_w like $\|f - g\|_{C^{1,\gamma}} |X - X_0|^{1+\gamma}$.)

Next, we can define the function on D_w , $\tilde{U} = U_f - U_g$. We see that \tilde{U} is a viscosity solution of

$$\mathcal{M}^+(D^2\tilde{U}) + \Lambda |\nabla\tilde{U}| \geq 0 \quad \text{and} \quad \mathcal{M}^-(D^2\tilde{U}) - \Lambda |\nabla\tilde{U}| \leq 0.$$

Therefore, we may apply Lemma 5.16, with $C_1 = C\|f - g\|_{C^{1,\gamma}}$, in order to deduce that (recall that $\|U_f - U_g\|_{L^\infty} \leq \|U_f - U_g\|_{C^{1,\gamma}}$)

$$|\nabla\tilde{U}(X_0)| \leq C\|f - g\|_{C^{1,\gamma}}.$$

Hence, by collecting the previous estimates we see that

$$|\nabla U_f(X_0) - \nabla U_g(X_0)| \leq C\|f - g\|_{C^{1,\gamma}}.$$

□

This next lemma is similar to the previous, except that we adapt the assumptions to allow that the equations for the respective solutions need not be the same for U_f, U_g .

Lemma 5.18. *Let f, g be such that $f, g \in \mathcal{K}^*(\delta, \gamma, m) \cap (m-C^{1,\gamma}(x_0))$, and $f - g \in C^{1,\gamma}(\mathbb{R}^d)$, and let $x_0 \in \mathbb{R}^d$ be a point such that $f(x_0) = g(x_0)$ and $\nabla f(x_0) = \nabla g(x_0)$. Further assume the existence $r > 0$ and of functions, V_f and V_g that satisfy in the viscosity sense, for $X_0 = (x_0, f(x_0)) = (x_0, g(x_0))$,*

$$\text{in } D_f \cap B_r(X_0), \quad |F(D^2V_f, \nabla V_f)| \leq C_1$$

$$\text{in } D_g \cap B_r(X_0), \quad |F(D^2V_g, \nabla V_g)| \leq C_1.$$

Then, there exists a $C = C(d, \lambda, \Lambda, \delta, \gamma, m, r)$ such that

$$|\nabla V_f(X_0) - \nabla V_g(X_0)| \leq C(\|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)} + C_1).$$

Proof of Lemma 5.18. The proof of Lemma 5.18 follows nearly identically to that of Lemma 5.17. We simply note that the auxiliary function, \tilde{U} will be defined as

$$\tilde{U} = V_f - V_g.$$

Because of the assumptions on f, g, V_f, V_g , we see that \tilde{U} satisfies in the viscosity sense,

$$\mathcal{M}^+(D^2\tilde{U}) + \Lambda \left| \nabla \tilde{U} \right| \geq -2C_1 \quad \text{and} \quad \mathcal{M}^-(D^2\tilde{U}) - \Lambda \left| \nabla \tilde{U} \right| \leq 2C_1.$$

The rest of the argument is identical, as this situation also falls within the scope of Lemma 5.16. \square

The next lemma, will be used in the proof of Theorem 5.13; it is a simple consequence of the definition of viscosity solutions, the uniform ellipticity of F , and the fact that F is assumed to be rotationally invariant in the D^2U argument. We omit the proof.

Lemma 5.19. *If F satisfies the assumption of rotational invariance in Section 2.2-(c), Ω is an open set, \mathcal{R} is a rotation, $U \in C^{0,1}(\Omega)$ is a viscosity solution of*

$$F(D^2U, \nabla U) = 0 \text{ in } \Omega,$$

then for $W = U \circ \mathcal{R}$, W solves in the viscosity sense

$$\left| F(D^2W, \nabla W) \right| \leq C\Lambda \left| \mathcal{R} - \text{Id} \right| \cdot \|\nabla U\|_{L^\infty}, \text{ in } \mathcal{R}^{-1}\Omega.$$

We are finally ready to prove Theorem 5.13.

Proof of Theorem 5.13. Let x_0 be an arbitrary point in \mathbb{R}^d , let f, g be such that $f, g \in \mathcal{K}^*(\delta, \gamma, m)$ and $f - g \in C^{1,\gamma}(\mathbb{R}^d)$, we proceed in four steps.

Step 0. ($f - g$ is small.) We note that it will suffice to prove the result under the assumption that $\|f - g\|_{L^\infty(\mathbb{R}^d)} \leq \frac{\delta}{2}$. Otherwise, we can simply use the fact that the constant, C , naturally depends upon δ , hence by the definition of I , (note, $I \geq 0$)

$$I(g, x) - I(f, x) \leq I(g, x) + I(f, x) \leq \frac{2\tilde{C}}{(\delta/2)}(\delta/2) \leq \frac{2\tilde{C}}{(\delta/2)}\|f - g\|_{L^\infty} \leq \frac{2\tilde{C}}{(\delta/2)}\|f - g\|_{C^{1,\gamma}}.$$

Thus, from here on, we will assume that $\|f - g\|_{L^\infty(\mathbb{R}^d)} \leq \frac{\delta}{2}$.

Step 1. (Reduction to $f(x_0) = g(x_0)$.)

Consider the function $\hat{f}(x) = f(x) + (g(x_0) - f(x_0))$. By step 0, we know $\hat{f} \in \mathcal{K}^*(\gamma, \delta/2, m)$. By Lemma 5.15, we have

$$\left| I(f, x_0) - I(\hat{f}, x_0) \right| \leq C|f(x_0) - g(x_0)| \leq C\|f - g\|_{L^\infty(\mathbb{R}^d)}.$$

Therefore, we have that $\hat{f}(x_0) = g(x_0)$ and

$$I(g, x_0) - I(f, x_0) \leq C\|f - g\|_{L^\infty(\mathbb{R}^d)} + \|I(g, x_0) - I(\hat{f}, x_0)\|.$$

From this observation, we conclude that to prove the theorem it is sufficient to show that

$$I(g, x_0) - I(f, x_0) \leq C\|f - g\|_{C^{1,\gamma}(\mathbb{R}^d)} \text{ at } x_0 \text{ s.t. } f(x_0) = g(x_0). \quad (5.9)$$

Accordingly, for the rest of the proof we assume that $x_0 \in \mathbb{R}^d$ is such that $f(x_0) = g(x_0)$, and focus on proving (5.9).

Step 2. (Rotation to achieve $\nabla f(x_0) = \nabla g(x_0)$.)

Let us denote by n_f and n_g respectively the outer normals to Γ_f and Γ_g at the point $(x_0, f(x_0))$. Let \mathcal{R} denote the rotation in \mathbb{R}^{d+1} given by

$$\mathcal{R}n_f = n_g, \text{ and } \mathcal{R}v = v \text{ if } v \perp \text{span}\{n_f, n_g\}.$$

We note that \mathcal{R} is a rotation that sends the tangent plane to Γ_f at X_0 to the tangent plane of Γ_g at X_0 . Furthermore, by definition, $|\mathcal{R} - \text{Id}| \leq C|n_f - n_g|$, we see that

$$|\mathcal{R} - \text{Id}| \leq C|n_f - n_g| \leq C|\nabla f(x_0) - \nabla g(x_0)| \leq C\|\nabla f - \nabla g\|_{L^\infty}. \quad (5.10)$$

Then, we denote by \mathcal{T} the isometry $\mathcal{T}X := \mathcal{R}(X - X_0) + X_0$. We note that we can choose r_0 small enough so that we can choose a function, w , that satisfies

$$B_{r_0}(X_0) \cap \mathcal{T}(D_f) = B_{r_0}(X_0) \cap D_w,$$

and we can still ensure $w \in K^*(\gamma, \frac{1}{2}\delta, 2m) \cap (m-C^{1,\gamma}(x_0))$. We then define a function \tilde{U} by

$$\tilde{U}(X) = U_f(\mathcal{T}^{-1}X) \quad \forall x \in B_{r_0}(X_0) \cap D_w.$$

Observe that

$$\nabla \tilde{U}(X_0) = \mathcal{R}\nabla U_f(X_0).$$

Since $|\nabla U_f(X_0)| \leq C(d, \lambda, \Lambda, \delta, M)$, it follows that

$$|\nabla \tilde{U}(X_0) - \nabla U_f(X_0)| \leq |(\mathcal{R} - \text{Id})\nabla U_f(X_0)| \leq C|\mathcal{R} - \text{Id}|,$$

and we conclude, per (5.10), that

$$|\nabla \tilde{U}(X_0) - \nabla U_f(X_0)| \leq C|\nabla f(x_0) - \nabla g(x_0)|. \quad (5.11)$$

Step 3. We first confirm the equation satisfied by \tilde{U} in $B_{r_0}(X_0) \cap D_w$, and then we will invoke Lemma 5.18. Indeed, since $\nabla U_f \in L^\infty(D_f)$ (via, e.g. Proposition 5.8), we see, by the rotational invariance assumed for F and Lemma 5.19, that in the viscosity sense, by Lemma 5.19,

$$\left| F(D^2\tilde{U}, \nabla \tilde{U}) \right| \leq C\Lambda|\mathcal{R} - \text{Id}| \cdot \|\nabla U_f\|_{L^\infty}, \text{ in } B_{r_0}(X_0) \cap D_w.$$

This shows that both \tilde{U} and U_g satisfy the assumptions of Lemma 5.18 (applied to the respective domains of D_w and D_g), and so we can conclude that

$$\left| \nabla \tilde{U}(X_0) - \nabla U_g(X_0) \right| \leq C(\|w - g\|_{C^{1,\gamma}} + \tilde{C}).$$

Here, we have used that

$$\tilde{C} = C\Lambda|\mathcal{R} - \text{Id}| \cdot \|\nabla U\|_{L^\infty}.$$

Collecting all of the above estimates, we see that under the assumption that $f(X_0) = g(X_0)$ and $\nabla f(X_0) = \nabla g(X_0)$, the estimate holds,

$$|\nabla U_f(X_0) - \nabla U_g(X_0)| \leq C\|f - g\|_{C^{1,\gamma}},$$

where we note, by construction of the function, w , that we can choose w so that $\|w - g\|_{C^{1,\gamma}} \leq \|f - g\|_{C^{1,\gamma}}$. Hence, the conclusion of the theorem under the step 3 assumption holds. Hence, combining with steps 1 and 2 give the original result. \square

Proof of Corollary 5.14. First, we give the argument for the case when $f - g - \phi$ attains a maximum. Indeed, we have for $s = f(x_0) - g(x_0) - \phi(x_0) \geq 0$ that

$$\begin{aligned} f - \phi - s &\leq g \text{ in } \mathbb{R}^d, \\ f - \phi - s &= g \text{ at } x_0. \end{aligned}$$

First, we note that by the assumptions that $f \in \mathcal{K}_*(\gamma, \delta, m)$, $g \in \mathcal{K}^*(\gamma, \delta, m)$, and $\phi \in C^{1,\gamma}$, this inequality shows that in fact $g, f \in m-C^{1,\gamma}(x_0)$. Hence, by Lemma 5.11, we know that $I(f, x_0)$ and $I(g, x_0)$ are defined classically.

Then, the GCP says that

$$I(f, x_0) \leq I(g + \phi + s, x_0).$$

Also since $s \geq 0$ and $g + \phi \in \mathcal{K}^*(\gamma, \delta, m)$, Lemma 5.15 yields

$$I(g + \phi + s, x_0) \leq I(g + \phi, x_0).$$

Furthermore, by Theorem 5.13 we know that

$$I(g + \phi, x_0) - I(g, x_0) \leq C\|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

Hence, putting this all together,

$$I(f, x_0) - I(g, x_0) \leq I(g + \phi + s, x_0) - I(g, x_0) \leq I(g + \phi, x_0) - I(g, x_0) \leq C\|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

The second part of the result, when $g - f - \phi$ attains a minimum is analogous. In this case, $s \leq 0$, and the GCP says that

$$I(f, x_0) \leq I(g - \phi - s, x_0).$$

Lemma 5.15 then shows that

$$I(g - \phi - s, x_0) \leq I(g - \phi, x_0).$$

The rest follows the same steps, invoking Theorem 5.13. \square

The next Lemma quantifies how the dependence of $I(f, x)$ on values of f far away from x decays as the distance to x grows (see also [38, Proposition 3.8]). It culminates in Lemma 5.22, which is a requirement for producing a min-max result of the form promised in Theorem 1.4, and it appears in Section 7 as Assumption 7.1-(vii).

Lemma 5.20. *There is a $C = C(d, \lambda, \Lambda, \delta, \gamma, m, L)$, such that given $f, g \in \mathcal{K}^*(\delta, \gamma, m) \cap m\text{-}C^{1,\gamma}(x_0)$, with $f, g \leq L$, $x_0 \in \mathbb{R}^d$, and $R > 1$ all such that*

$$f \equiv g \text{ in } B_R(x_0),$$

and if a.e. $B_R(x_0)$, then we have

$$|I(f, x_0) - I(g, x_0)| \leq CR^{-2}\|f - g\|_{L^\infty(\mathbb{R}^d)} + CR^{-1-\gamma}.$$

Proof of Lemma 5.20. First, we note that the assumption that $f, g \in m\text{-}C^{1,\gamma}(x_0)$ ensures that $I(f, x_0)$ and $I(g, x_0)$ are well defined, via Corollary 5.12.

We will proceed in two steps. The first step is the result under the additional assumption that the maximum of f over $B_R(x_0)$ occurs at x_0 . The second step is to use the \mathcal{K}^* upper bound for f , with Theorem 5.13, to reduce to the first step, incurring an extra error.

We will only show half of the inequality in that

$$I(f, x_0) - I(g, x_0) \leq CR^{-2}\|f - g\|_{L^\infty(\mathbb{R}^d)} + CR^{-1-\gamma}. \quad (5.12)$$

The reverse inequality follows by constructing an appropriate subsolution, where below we treat the case of a supersolution. The modifications are standard.

Step 1 (Assume $f(x_0)$ is the maximum of f .)

Let us assume, without loss of generality, that $x_0 = 0$ and for $|x| \leq R$, $f(x) \leq f(0)$. Let us consider then, the set

$$T := \{(x, x_{d+1}) \in \mathbb{R}^{d+1} \mid 0 \leq x_{d+1} \leq f(x), x \in B_{R/2}(0)\}.$$

We will construct a barrier that traps f on the top boundary, at $(0, f(0))$. For a function $\phi_0(y)$ ($y \in [0, L]$) to be determined, let

$$\psi_R(x, x_{d+1}) = \tilde{\psi}(x/R) + R^{-2}\phi_0(x_{d+1}), \quad \tilde{\psi}(x) := |x|^2(1 + |x|^2)^{-1}.$$

A straightforward computation shows that

$$\nabla\psi_R(x, x_{d+1}) = (R^{-1}(\nabla\tilde{\psi})(x/R), R^{-2}\phi'_0(x_{d+1}))$$

and

$$D^2\psi_R(x, x_{d+1}) = \begin{pmatrix} R^{-2}(D^2\tilde{\psi})(x/R) & 0 \\ 0 & R^{-2}\phi_0''(x_{d+1}) \end{pmatrix}.$$

Then, if $\phi_0(x_{d+1})'' \leq 0$ and $\phi_0'(x_{d+1}) \leq 0$ for all $x_{d+1} \in [0, L]$, we have

$$\mathcal{M}^+(D^2\psi_R) + \Lambda|\nabla\psi_R| \leq \Lambda R^{-2}\|D^2\tilde{\psi}\|_\infty + \lambda R^{-2}\phi_0''(x_{d+1}) + \Lambda R^{-2}\|\nabla\tilde{\psi}\|_\infty - \Lambda R^{-2}\phi_0'(x_{d+1}).$$

Therefore,

$$\mathcal{M}^+(D^2\psi_R) + \Lambda|\nabla\psi_R| \leq R^{-2} \left\{ \lambda\phi_0''(x_{d+1}) - \Lambda\phi_0'(x_{d+1}) + \Lambda\|\tilde{\psi}\|_{C^2} \right\}. \quad (5.13)$$

With the goal in mind that is to force this inequality to be non-positive (i.e. ψ_R should be a super solution), we can now choose ϕ_0 . Let us take, for some $M, b, c > 0$, and restricting $y \in [0, c]$,

$$\phi_0(y) = M(1 - e^{-b(c-y)}).$$

Then, for every $y \geq 0$ we have

$$\phi_0'(y) = -bMe^{-b(c-y)} \leq 0, \quad \phi_0''(y) = -b^2Me^{-b(c-y)} \leq 0,$$

and thus

$$\lambda\phi_0''(y) - \Lambda\phi_0'(y) + \Lambda\|\tilde{\psi}\|_{C^2} = (-\lambda b^2 + \Lambda b)Me^{-b(c-y)} + \Lambda\|\tilde{\psi}\|_{C^2}.$$

We first choose $b > 0$ to be large enough so that

$$-\lambda b^2 + \Lambda b \leq -1.$$

Furthermore, we see that for $y \in [0, c]$, and $c \leq L$

$$e^{-b(c-y)} \geq e^{-bc} \geq e^{-bL},$$

thus with our choice of b ,

$$(-\lambda b^2 + \Lambda b)Me^{-b(c-y)} \leq (-\lambda b^2 + \Lambda b)Me^{-bL}.$$

Now, we can choose M large enough, depending only on λ, Λ, b, L , so that

$$(-\lambda b^2 + \Lambda b)Me^{-bL} \leq -\Lambda\|\tilde{\psi}\|_{C^2}.$$

Finally, we set $c = f(0)$. Hence, we have attained

$$\mathcal{M}^+(D^2\psi_R) + \Lambda|\nabla\psi_R| \leq 0 \text{ in } D_L,$$

and we recall that by our assumption that $f = g$ in B_R , $D_f \cap T = D_g \cap T$, and that $D_f \subset D_L$.

We gather the following properties of ψ_R

$$\begin{aligned} \psi_R &\geq \tilde{\psi}(x/R) \text{ on } \Gamma_f, \\ \psi_R &\geq c_0 > 0 \text{ on } \{y = 0\}, \\ \psi_R &\geq c_0 > 0 \text{ on } \{(x, y) : |x| = R/2\}. \end{aligned}$$

We further recall that by Proposition 5.9 we have

$$\|U_f - U_g\|_{L^\infty(D_f \cap D_g)} \leq C\|f - g\|_{L^\infty(\mathbb{R}^d)}.$$

Thus, this means that for a universal choice of C , we can show that

$$\text{on } \partial T, \quad U_f - U_g \leq C\|f - g\|_{L^\infty}\psi_R.$$

Since U_f and U_g are viscosity solutions of the same equation, we see that $W = U_f - U_g$ is a viscosity subsolution of

$$\mathcal{M}^+(D^2W) + \Lambda|\nabla W| \geq 0.$$

Using the comparison theorem in T with the functions W and $C\|f - g\|_{L^\infty}\psi_R$, hence

$$\forall X \in T, \quad U_f(X) - U_g(X) \leq C\|f - g\|_{L^\infty}\psi_R(X).$$

By construction, we have that $U_f(0, f(0)) = U_g(0, f(0)) = \psi_R(0, f(0)) = 0$, and so

$$\partial_n U_f(0, f(0)) - \partial_n U_g(0, f(0)) = \partial_n W(0, f(0)) \leq \partial_n \psi_R(0, f(0)) \leq CR^{-2} \|f - g\|_{L^\infty(\mathbb{R}^d)},$$

which gives the desired result in (5.12). This completes the proof in the setting of Step 1.

Step 2 (reducing to x_0 is a max of f in B_R).

Given that we are assuming $f \in K^*(\gamma, \delta, m)$, we know that for an appropriate choice of c , depending upon m ,

$$\forall x \in B_R(x_0), \quad f(x) \leq f(x_0) + Cm \left| \frac{x - x_0}{R} \right|^{1+\gamma}.$$

Hence, replacing both f and g by

$$\tilde{f}(x) = f(x) - Cm \left| \frac{x - x_0}{R} \right|^{1+\gamma} \quad \text{and} \quad \tilde{g}(x) = g(x) - Cm \left| \frac{x - x_0}{R} \right|^{1+\gamma},$$

we see that we satisfy the assumptions of Step 1.

Furthermore, we know, from Theorem 5.13, after extending the function $Cm \left| \frac{x - x_0}{R} \right|^{1+\gamma}$ to all of \mathbb{R}^d in a way that does not increase its $C^{1,\gamma}$ norm by a factor of more than, 2, that we have

$$\left| I(f, x_0) - I(\tilde{f}, x_0) \right| \leq CmLR^{-1-\gamma}.$$

This, combined with the result in Step 1, shows the desired estimate. \square

Remark 5.21. *It is worth comparing this decay rate with what one observes for the Dirichlet-to-Neumann map in the half-space for the Laplacian. Of course, one obtains as the D-to-N for the Harmonic extension the half Laplacian, $-(-\Delta)^{1/2}f$. Because the harmonic extension satisfies an equation in half-space, a simple rescaling argument can be shown that in the context of the previous Lemma (with $f \equiv g$ in $B_R(x_0)$),*

$$|(-\Delta)^{1/2}(f - g)(x_0)| \leq CR^{-1} \|f - g\|_{L^\infty}.$$

This is, of course, different from what we have obtained above. The discrepancy can be attributed to two factors: we work in a finite width domain, with a Dirichlet condition at $y = 0$; and our equation does not rescale in a positively homogeneous way (different powers from the Hessian and gradient). Thus, as seen from our barrier, ψ_R , the term R^{-2} is not surprising. The way in which this estimate is invoked is completely unaffected by the actual decay on the right hand side of the estimate.

Next, we remove the assumption from the previous lemma that required $f \equiv g$ in $B_R(x_0)$.

Lemma 5.22. *If $C = C(d, \lambda, \Lambda, \delta, \gamma, m, L)$ is as in Lemma 5.20, that $f, g \in \mathcal{K}^*(\delta, \gamma, m)$ and $f - g \in C^{1,\gamma}(B_{2R}(x_0))$, with $f, g \leq L$, $x_0 \in \mathbb{R}^d$, and $R > 1$, then*

$$|I(f, x_0) - I(g, x_0)| \leq C \|f - g\|_{C^{1,\gamma}(B_{2R}(x_0))} + CR^{-2} \|f - g\|_{L^\infty(\mathbb{R}^d)} + CR^{-1-\gamma}.$$

Proof of Lemma 5.22. Let ψ be a smooth function such that $0 \leq \psi \leq 1$, $\psi \equiv 1$ in $B_R(x_0)$, $\psi \equiv 0$ outside $B_{2R}(x_0)$, and such that

$$\|\nabla \psi\|_{L^\infty} \leq CR^{-1}, \quad \|D^2 \psi\|_{L^\infty} \leq CR^{-2}.$$

Then, let

$$\hat{f} = \psi f + (1 - \psi)g,$$

In particular, $\hat{f} \equiv g$ outside $B_{2R}(x_0)$. Therefore, since $f - g \in C^{1,\gamma}(B_{2R}(x_0))$, we have $\hat{f} - g \in C^{1,\gamma}(\mathbb{R}^d)$ (as $\hat{f} - g = \psi \cdot (f - g)$). Thus, by Theorem 5.13, it follows that

$$\begin{aligned} |I(g, x) - I(\hat{f}, x)| &\leq C \|g - \hat{f}\|_{C^{1,\gamma}(\mathbb{R}^d)} \\ &= C \|\psi \cdot (g - f)\|_{C^{1,\gamma}(B_{2R}(x_0))} \\ &\leq C \|\psi\|_{C^{1,\gamma}} \|f - g\|_{C^{1,\gamma}(B_{2R}(x_0))} \end{aligned}$$

Meanwhile, since $\hat{f} \equiv f$ inside $B_R(x_0)$, by Lemma 5.20, we have

$$|I(f, x) - I(\hat{f}, x)| \leq CR^{-2} \|f - \hat{f}\|_{L^\infty(\mathbb{R}^d)} + CR^{-1-\gamma}$$

However, note that

$$\|f - \hat{f}\|_{L^\infty(\mathbb{R}^d)} = \|(1 - \psi)(f - g)\|_{L^\infty(\mathbb{R}^d)} \leq \|f - g\|_{L^\infty(\mathbb{R}^d)}.$$

Then, by the triangle inequality, we conclude that

$$|I(f, x_0) - I(g, x_0)| \leq C \|f - g\|_{C^{1,\gamma}(B_{2R}(x_0))} + CR^{-2} \|f - g\|_{L^\infty(\mathbb{R}^d)} + CR^{-1-\gamma}.$$

□

Proposition 5.23. *If x_0 is fixed, $f, g \in \mathcal{K}^*(\gamma, \delta, m)$ and $f - g \in C^{1,\gamma}(B_{2R}(x_0))$, $f, g \leq L$, and $f \geq g$, then there exists $C = C(d, \lambda, \Lambda, \delta, \gamma, m, L)$ so that for $R > 1$,*

$$I(f, x_0) \leq I(g, x_0) + C \left(\operatorname{osc}_{B_{2R}(x_0)}(f - g) + \|\nabla f - \nabla g\|_{C^\gamma(B_{2R}(x_0))} + R^{-2} \operatorname{osc}_{L^\infty(\mathbb{R}^d)}(f - g) + R^{-1-\gamma} \right).$$

Proof of Proposition 5.23. Fix $x_0 \in \mathbb{R}^d$ and $R > 1$. Let $c := \inf_{B_{2R}(x_0)}(f - g)$ and note that

$$\begin{aligned} \|f - c - g\|_{L^\infty(B_{2R}(x_0))} &\leq \operatorname{osc}_{B_{2R}(x_0)}(f - g) \\ \|f - c - g\|_{L^\infty(\mathbb{R}^d)} &\leq \operatorname{osc}_{\mathbb{R}^d}(f - g) \end{aligned}$$

Since, by assumption, $c \geq 0$, Lemma 5.15 shows that

$$I(f, x_0) \leq I(f - c, x_0) \quad \forall x \in \mathbb{R}^d.$$

Then, bounding $I(f - c, x_0) - I(g, x_0)$ from above using Lemma 5.22, it follows that

$$I(f, x_0) - I(g, x_0) \leq C \|f - c - g\|_{C^{1,\gamma}B_{2R}(x_0)} + CR^{-2} \|f - c - g\|_{L^\infty(\mathbb{R}^d)} + CR^{-1-\gamma},$$

and using the above bounds on $\|f - c - g\|_{L^\infty}$ in B_{2R} and \mathbb{R}^d with the respective oscillation of $f - g$, we have

$$I(f, x_0) - I(g, x_0) \leq C \left(\operatorname{osc}_{B_{2R}(x_0)}(f - g) + \|\nabla f - \nabla g\|_{C^\gamma(B_{2R}(x_0))} \right) + CR^{-2} \operatorname{osc}_{L^\infty(\mathbb{R}^d)}(f - g) + CR^{-1-\gamma}.$$

□

A useful corollary of Proposition 5.23 is that we can construct a bump function that does not increase the values of I too much.

Corollary 5.24. *Let $\phi(x) = |x|^2(1 + |x|^2)^{-1}$ and for $R > 1$ and $C \geq 0$, define $\phi_R(x) = C + \phi(x/R)$. Given any $\varepsilon > 0$, there exists $C_1(\varepsilon)$, such that for all $R > C_1(\varepsilon)$ for all $g \in \mathcal{K}^*(\gamma, \delta, m) \cap (m - C^{1,\gamma}(x_0))$ with $g + \phi_R \in \mathcal{K}^*(\gamma, \delta, m)$,*

$$I(g + \phi_R, x_0) - I(g, x_0) \leq \varepsilon.$$

Proof of Corollary 5.24. We observe that

$$\begin{aligned} \operatorname{osc}_{\mathbb{R}^d} \phi_R &= 1 \\ \|\nabla \phi_R(x)\|_{L^\infty(\mathbb{R}^d)} &\leq R^{-1} \|\nabla \phi_1\|_{L^\infty(\mathbb{R}^d)} \\ [\nabla \phi_R(x)]_{C^\gamma(\mathbb{R}^d)} &\leq R^{-1-\gamma} [\nabla \phi_1(x)]_{C^\gamma(\mathbb{R}^d)}. \end{aligned}$$

Next, we note that for all $R > 1$, $g + \phi_R \in \mathcal{K}(\gamma, \delta, m + 1)$. Let us fix x , and let us introduce a temporary parameter, $\rho > 1$. We can invoke Proposition 5.23 for B_ρ , to obtain

$$\forall x \in \mathbb{R}^d, \quad I(g + \phi_R, x) - I(g, x) \leq C \left(\operatorname{osc}_{B_{2\rho}(x)} \phi_R + \|\nabla \phi_R\|_{C^\gamma(B_{2\rho}(x))} + \rho^{-2} + \rho^{-1-\gamma} \right).$$

First, we can choose $C_1(\varepsilon)$ large enough so that for all $\rho > C_1(\varepsilon)$, we have

$$C(\rho^{-2} + \rho^{-1-\gamma}) < \frac{\varepsilon}{2}.$$

Next we take ρ to be fixed, and we can choose $C_2(\varepsilon, \rho)$ large enough so that

$$C \left(\operatorname{osc}_{B_{2\rho}(x)} \phi_R + \|\nabla \phi_R\|_{C^\gamma(B_{2\rho}(x))} \right) < \frac{\varepsilon}{2},$$

which is possibly because of the decay of $\nabla \phi_R$ that is uniform in x . □

6. THE TWO-PHASE OPERATOR

Let us now return to the analysis for the two phase operator. We will show that the two-phase boundary condition can be completely characterized as a combination of two operators in the same category as I , as in Section 5.

We recall that the sets, $\mathcal{K}(\gamma, \delta, m)$, $\mathcal{K}^*(\gamma, \delta, m)$, $\mathcal{K}_*(\gamma, \delta, m)$ were defined in (5.3)–(5.5). For $L > \delta$ fixed, we consider a modification of these sets as follows:

$$\mathcal{K}(\gamma, \delta, m, L) = \{f \in \mathcal{K}(\gamma, \delta, m) : f \leq L - \delta\}; \quad (6.1)$$

and similarly,

$$\begin{aligned} \mathcal{K}^*(\gamma, \delta, m, L) &= \{f \in \mathcal{K}^*(\gamma, \delta, m) : f \leq L - \delta\} \\ \mathcal{K}_*(\gamma, \delta, m, L) &= \{f \in \mathcal{K}_*(\gamma, \delta, m) : f \leq L - \delta\}. \end{aligned}$$

For f in any of these sets, we define the domains,

$$\begin{aligned} D_f^+ &:= D_f = \{(x, x_{d+1}) : 0 < x_{d+1} < f(x)\} \\ D_f^- &:= \{(x, x_{d+1}) : f(x) < x_{d+1} < L\}. \end{aligned}$$

Furthermore, we define U_f as the unique continuous viscosity solution to the equations

$$\begin{cases} F_1(D^2 U_f, \nabla U_f) = 0 & \text{in } D_f^+, \\ F_2(D^2 U_f, \nabla U_f) = 0 & \text{in } D_f^-, \\ U_f = 0 & \text{on } \Gamma_f, \\ U_f = 1 & \text{on } \Gamma_0, \\ U_f = -1 & \text{on } \Gamma_L, \end{cases} \quad (6.2)$$

which will have a unique solution whenever $f \in C^{0,1}(\mathbb{R}^d)$ and $\delta \leq f \leq L - \delta$, see Proposition 5.3. (We recall that the notation for $\Gamma_f, \Gamma_0, \Gamma_L$ is in Section 2.3.)

In order to state the free boundary condition, we assume that we are given a Lipschitz continuous function

$$G : (0, \infty)^2 \rightarrow \mathbb{R},$$

which satisfies the following uniform monotonicity conditions for a.e. (a, b) ,

$$\lambda_0 \leq \frac{\partial}{\partial a} G(a, b) \leq \Lambda_0, \quad \lambda_0 \leq -\frac{\partial}{\partial b} G(a, b) \leq \Lambda_0. \quad (6.3)$$

The function U_f defined in (6.2) is always continuous across Γ_f , but in general it will not be differentiable across it. However, in most situations, it will be differentiable separately on each side of Γ_f . In particular, when $f \in \mathcal{K}(\gamma, \delta, m, L)$, the following two limits exist

$$\begin{aligned} \nabla U_f^+(X) &:= \lim_{X' \rightarrow X, X' \in D_f^+} \nabla U_f(X'), \\ \nabla U_f^-(X) &:= \lim_{X' \rightarrow X, X' \in D_f^-} \nabla U_f(X'). \end{aligned}$$

When these operations do exist, they coincide with the slightly more general operators, $\partial_n^\pm U$, which we define as:

$$\text{for } X_0 \in \Gamma_f, \text{ and } n(X_0) \text{ the unit normal derivative to } \Gamma_f, \text{ pointing into the set } D^+ f, \\ \partial_n^+ U(X_0) := \lim_{t \rightarrow 0} \frac{U(X_0 + tn(X_0)) - U(X_0)}{t} \quad \text{and} \quad \partial_n^- U(X_0) = -\lim_{t \rightarrow 0} \frac{U(X_0 - tn(X_0)) - U(X_0)}{t}. \quad (6.4)$$

We note that $\partial_n^\pm U$ are normalized so that in the context of U_f solving (6.2), we have that both $\partial_n^+ U_f$ and $\partial_n^- U_f$ are positive, and we also note that the inward normal to D_f^- is the vector $(-n(X_0))$.

We define the two-phase analogue to the one-phase operator (5.2) by

$$H(f, x) := G(\partial_n^+ U_f(x, f(x)), \partial_n^- U_f(x, f(x))). \quad (6.5)$$

As we are about to show, this ‘‘two phase’’ operator, (6.5), may be expressed as a composition with two one-phase operators of the form (5.2). This fact makes it easy to extend basically all of the results in the previous section to the operator in (6.5). Indeed, to (6.5) we associate two operators I^+ and I^- of the form (5.2), as follows. First, given f we define U_f^+ as the unique solution of

$$\begin{cases} F_1(D^2 U_f^+, \nabla U_f^+) = 0 & \text{in } D_f^+, \\ U_f^+ = 0 & \text{on } \Gamma_f, \\ U_f^+ = 1 & \text{on } \Gamma_0. \end{cases} \quad (6.6)$$

Similarly, we can define U_f^- as the unique solution of

$$\begin{cases} F_2(D^2 U_f^-, \nabla U_f^-) = 0 & \text{in } D_f^-, \\ U_f^- = 0 & \text{on } \Gamma_f, \\ U_f^- = -1 & \text{on } \Gamma_L. \end{cases} \quad (6.7)$$

Using as notation $n^-(X_0)$ to be the inward normal to the set, D_f^- , we define the operators I^\pm as the inward normals to their respective phases,

$$I^+(f, x) := \partial_n U_f^+(x, f(x)), \quad I^-(f, x) := -\partial_{n^-} U_f^-(x, f(x)) = \partial_n^- U_f^-(x, f(x)). \quad (6.8)$$

This means that we will also write the two-phase operator as

$$H(f, x) := G(I^+(f, x), I^-(f, x)). \quad (6.9)$$

We note again we enforce the convention that we seek operators such that both I^\pm are non-negative quantities. However, thanks to the boundary condition $U_f^- = -1$ on Γ_L , it is not hard to check that $-I^-$ obeys the GCP over the set, $\mathcal{K}(\gamma, \delta, m, L)$.

In order to make an exact analog between I^- defined in (6.8) and the operator, I , defined in (5.2), we introduce a transformation so that I^- can be recognized as an operator that uses a solution to

an equation in $D_f^+ = D_f$ (in the notation of Section 5). To this end, we need to transform both the equation for the negative phase and the lower boundary that was previously Γ_f . Thus, we define

$$\tilde{F}(Q, p) := -F(-Q, p), \quad \text{and} \quad \tilde{f}(x) := f(-x),$$

and we take \tilde{U}_f^- to be the unique solution of

$$\begin{cases} \tilde{F}_2(D^2\tilde{U}_f^-, \nabla\tilde{U}_f^-) = 0 & \text{in } D_{L-\tilde{f}}, \\ \tilde{U}_f^- = 0 & \text{on } \Gamma_{L-\tilde{f}}, \\ \tilde{U}_f^- = 1 & \text{on } \Gamma_0. \end{cases} \quad (6.10)$$

The natural relationship between $I^-(f, x)$ and \tilde{U}_f^- is verified by the following lemma.

Lemma 6.1. *For all $f \in \mathcal{K}(\gamma, \delta, m, L)$, if $I^-(f, x)$ is defined in (6.8), $I_{\tilde{F}_2}$ is defined as in (5.1) and (5.2) with F replaced by \tilde{F}_2 , and \tilde{U}_f^- is defined in (6.10), then*

$$I^-(f, x) = -\partial_n \tilde{U}_f^-(-x, L - \tilde{f}(-x)) = -I_{\tilde{F}_2}(L - \tilde{f}, -x), \quad (6.11)$$

and $-I_{\tilde{F}_2}$ is an operator that satisfies the definitions and assumptions of Section 5.

Proof of Lemma 6.1. First observe that if $y \geq f(x)$, then $L - y \leq L - \tilde{f}(-x)$, and in fact

$$(x, y) \in D_f^- \iff (-x, L - y) \in D_{L-\tilde{f}}.$$

Furthermore, we list some similar and useful related observations:

$$\begin{aligned} (x, y) \in \Gamma_{L-\tilde{f}} &\iff (-x, L - y) \in \Gamma_f, \\ (x, y) \in \Gamma_f &\iff (-x, L - y) \in \Gamma_{L-\tilde{f}}, \\ n_f(x, f(x)) &= n_{L-\tilde{f}}(-x, L - \tilde{f}(-x)). \end{aligned}$$

Thus, given U_f as in (6.2), define $V : D_{L-\tilde{f}} \rightarrow \mathbb{R}$ by

$$V(x, y) = -U(-x, L - y).$$

The function V solves

$$\begin{cases} -F_2(-D^2V, \nabla V) = 0 & \text{in } D_{L-\tilde{f}}, \\ V = 0 & \text{on } \Gamma_{L-\tilde{f}}, \\ V = 1 & \text{on } \Gamma_0. \end{cases}$$

In other words by uniqueness of solutions, we have $V = \tilde{U}_f^-$.

Using that if $(x, y) \in \Gamma_f$, then

$$\nabla U_f(x, y) = \nabla V(-x, L - y),$$

we see that

$$\nabla U_f(x, y) \cdot n_f(x, f(x)) = \nabla V(-x, L - y) \cdot n_{L-\tilde{f}}(-x, L - \tilde{f}(-x)).$$

We also observe that if F_2 satisfies the (λ, Λ) ellipticity of Definition 2.4, then \tilde{F}_2 enjoys the same ellipticity constants and extremal operators. Furthermore, by definition (5.1) and (5.2) with F replaced by \tilde{F}_2 we see that by construction,

$$\partial_n \tilde{U}_f^-(-x, L - \tilde{f}(-x)) = I_{\tilde{F}_2}(L - \tilde{f}, -x)$$

□

We conclude this section with the analogous statements from Section 5 (Theorem 5.13 and Corollary 5.14), but tailored to the two-phase operator, H .

Theorem 6.2. *If f and g satisfy $f \in \mathcal{K}_*(\delta, \gamma, m, L)$ and $g \in \mathcal{K}^*(\delta, \gamma, m, L)$ as well as $\phi \in C^{1,\gamma}(\mathbb{R}^d)$, $g + \phi \in \mathcal{K}^*(\delta, \gamma, m)$, $f - \phi \leq L - \delta$, $f - \phi$ is m - $C^{1,\gamma}$ -semi-convex, and all are such that*

$$(f - g) - \phi \text{ has a non-negative global maximum at } x_0 \in \mathbb{R}^d,$$

then, with $C = C(d, \lambda, \Lambda, \Lambda_0, \delta, \gamma, m)$, we have

$$H(f, x_0) - H(g, x_0) \leq C \|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

Proof. First of all, observe that (6.3) implies that

$$G(a, b) - G(a', b') \leq \Lambda_0(a - a')_+ + \Lambda_0(b' - b)_+,$$

for all $a, a', b, b' \in (0, \infty)$. Therefore, recalling the definitions of I^\pm in (6.8) and the monotonicity of (6.3), we see that

$$\begin{aligned} H(f, x_0) - H(g, x_0) &= G(I^+(f, x_0), I^-(g, x_0)) - G(I^+(f, x_0), I^-(g, x_0)) \\ &\leq \Lambda_0(I^+(f, x) - I^+(g, x))_+ + \Lambda_0(I^-(g, x) - I^-(f, x))_+. \end{aligned}$$

Now, according to Corollary 5.14,

$$I^+(f, x_0) - I^+(g, x_0) \leq C \|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

Lemma 6.1 and Corollary 5.14 (applied when a minimum is attained) show that

$$(I^-)(g, x_0) - (I^-)(f, x_0) = -I_{\tilde{F}_2}(L - \tilde{g}, -x_0) - (-I_{\tilde{F}_2}(L - \tilde{f}, -x_0)) \leq C \|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

(Here we note that we used $f - \phi \leq L - \delta$ and $f - \phi$ m - $C^{1,\gamma}$ -semi-convex to obtain $L - \tilde{f} - (-\tilde{\phi}) \in \mathcal{K}^*(\gamma, \delta, m)$, and we used that $(L - \tilde{f}) - (L - \tilde{g}) - (-\tilde{\phi})$ attains a minimum at x_0 .) It then follows that

$$H(g, x_0) - H(f, x_0) \leq C \Lambda_0 \|\phi\|_{C^{1,\gamma}(\mathbb{R}^d)}.$$

□

7. AN INTEGRO-DIFFERENTIAL REPRESENTATION OF I AND H

This section proves that the free boundary operator, I , defined in (5.2), and the two-phase operator, H , defined in (6.5), are in fact integro-differential operators, and they can be represented, via a min-max procedure, as claimed in Theorem 1.4. Although the perceptive reader will have noticed, a uniqueness theorem for parabolic equations involving I or H can be deduced from Corollary 5.24, we still think it is useful to pursue the integro-differential development for I . We would like to point out that, in our opinion, we would have not have realized the properties of I in Section 5 (especially in Corollary 5.14 and Proposition 5.23) without expecting that I is a min-max of integro-differential operators. In particular, the integro-differential framework was essential for our choice to pursue I as a Lipschitz mapping on subsets of $C^{1,\gamma}$, instead of the possibly more obvious space of $C^{1,1}$. This distinction is significant, as the reader will see in this section, because it means that the representative integro-differential operators will not contain any second order terms— a welcome simplification for using the integro-differential theory. Furthermore, as one will subsequently see, the preservation of modulus (claimed in Theorem 1.1) can be deduced purely from a comparison theorem for these fractional parabolic equations. However, we hope that the integro-differential framework will be useful to gain higher regularity results. In future contexts, it seems that, e.g. Theorem 6.2 will not be enough, but rather one will need more refined information as hinted by Theorem 1.4. Thus, we have chosen to study some initial properties of the integro-differential representation of I here.

The results of this section do not rely on the underlying free boundary problem, and so we treat these operators as a class of their own.

7.1. The general class of operators. To this end, we assume that J is an operator that acts on functions on \mathbb{R}^d and that satisfies the following assumptions.

Assumption 7.1. J has the following properties:

- (i) $0 < \gamma < 1$ is fixed;
- (ii) $J : \left(\bigcup_{\delta > 0} \bigcup_{m > \delta} \mathcal{K}(\gamma, \delta, m) \right) \rightarrow C^0(\mathbb{R}^d)$.
- (iii) For each δ and m , J is a Lipschitz mapping on the sets $\mathcal{K}(\gamma, \delta, m)$, whose Lipschitz constant, $C(\delta, m)$ increases as δ decreases or m increases.
- (iv) J satisfies the GCP (Definition 2.2).
- (v) J is translation invariant.
- (vi) If $f \in \mathcal{K}(\gamma, \delta, m)$ and $c > 0$ is a constant, then

$$\forall x \in \mathbb{R}^d, J(f + c, x) \leq J(f, x).$$
- (vii) J enjoys the operator splitting property: $\exists C = C(\gamma, \delta, m)$ and $\exists \omega$ (a modulus with $\omega(R) \rightarrow 0$ as $R \rightarrow \infty$), such that $\forall f, g \in \mathcal{K}(\gamma, \delta, m)$, for $R > 1$

$$\|J(f, \cdot) - J(g, \cdot)\|_{L^\infty(B_R)} \leq C(\|f - g\|_{C^{1,\gamma}(B_{2R})} + \omega(R)\|f - g\|_{L^\infty(\mathbb{R}^d)} + \omega(R)).$$

Remark 7.2. We note that as presented, the assumptions for J only include operators like $I(f)$, $G(I(f))$, $G(I(f)) \cdot \sqrt{1 + |\nabla f|^2}$, which appear in the one-phase problem. However, thanks to Lemma 6.1, we see that the operator, $H(f)$, which is required for the two-phase problem, reduces to the case of $I(f)$, with the additional constraint that all of the constants involved will also depend upon the height of the upper boundary, which is given by L in Section 6.

The first results we will show are that any J , as in Assumption 7.1, can be represented as a min-max of integro-differential operators, followed by proving some properties of the corresponding extremal operators for the class that includes J . The basis of the results in this section is taken from the main result in [37], and for convenience, we restate it here, in the context of the operators, J .

Theorem 7.3 (from Theorem 1.6, Proposition 1.7 of [37]). *If J satisfies items (iii), (iv), (vii) of Assumption 7.1, then*

$$\forall f \in \mathcal{K}(\gamma, \delta, m), J(f, x) = \min_{g \in \mathcal{K}(\gamma, \delta, m)} \max_{L \in \mathcal{L}(\mathcal{K}(\gamma, \delta, m))} \{J(g, x) + L(f - g, x)\}, \quad (7.1)$$

where $\mathcal{L}(\mathcal{K}(\gamma, \delta, m))$ is a collection of linear operators, $L : C^{1,\gamma} \rightarrow C^0$, that enjoy the following form:

$$L(f, x) = c(x)f(x) + b(x) \cdot \nabla f(x) + \int_{\mathbb{R}^d} (f(x+h) - f(x) - \mathbb{1}_{B_1}(h)\nabla f(x) \cdot h) \mu(x, dh), \quad (7.2)$$

and satisfy for some C , uniformly depending on the norm $\|J\|_{\mathcal{K}(\gamma, \delta, m) \rightarrow C^{\gamma'}}$,

$$\|c\|_{L^\infty}, \|b\|_{L^\infty} \leq C \quad \text{and} \quad \sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} \min\{|h|^{1+\gamma}, 1\} \mu(x, dh) \leq C. \quad (7.3)$$

Finally, the Lévy measures satisfy, for a uniform constant across the family, \mathcal{L} ,

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_R} \mu(x, dh) \leq C\omega(R). \quad (7.4)$$

Although for many purposes, this representation in (7.1) and (7.2) will suffice, in our current context we will require (and prove) something slightly more precise. It turns out that since J is translation invariant, one can take a different approach to the representation in (7.1) and obtain more detail on the collection of L in (7.2).

A key building block that goes into the proof of Theorem 7.3 is the structure of linear functionals on $\mathcal{K}(\gamma, \delta, m)$ that enjoy the GCP. We state it explicitly, and refer the proof to modifications of [32] or [37].

Proposition 7.4. *If $\ell : \mathcal{K}(\gamma, \delta, m) \rightarrow \mathbb{R}$ is a bounded linear functional that enjoys the GCP based at $x_0 = 0$ (Definition 2.2), then there exists $b \in \mathbb{R}$, $c \in \mathbb{R}$, and a measure, μ , so that $\forall f \in \mathcal{K}(\gamma, \delta, m)$,*

$$\ell(f) = cf(0) + b \cdot \nabla f(0) + \int_{\mathbb{R}^d} (f(h) - f(0) - \mathbb{1}_{B_1}(h) \nabla f(0) \cdot h) \mu(dh),$$

where b , c , and μ satisfy the same bounds as in Theorem 7.3, but none of them depend on x .

We will state the following theorem, and the proof will appear after two other results are established.

Theorem 7.5 (Translation invariant min-max). *If J is as in Assumption 7.1, then J admits a min-max representation as in (7.1) in which each L is also translation invariant; i.e. $L \in \mathcal{L}_{inv}$, and \mathcal{L}_{inv} is the class which contains all linear operators of the form*

$$L(f, x) = cf(x) + b \cdot \nabla f(x) + \int_{\mathbb{R}^d} (f(x+h) - f(x) - \mathbb{1}_{B_1}(h) \nabla f(x) \cdot h) \mu(dh), \quad (7.5)$$

where each of c , b , μ are independent of x . Furthermore, given a $C_1 > 0$, there exists a $C_2 > 0$ such that for all J with a Lipschitz norm bounded by C_1 , all such c , b , and μ , resulting from an $L \in \mathcal{L}_{inv}$, we have $|c| \leq C_2$, $|b| \leq C_2$,

$$\int_{\mathbb{R}^d} \max(|h|^{1+\gamma}, 1) \mu(dh) \leq C_2, \quad \text{and} \quad \int_{\mathbb{R}^d \setminus B_R} \mu(dh) \leq C_2 \omega(R).$$

The class of operators, \mathcal{L}_{inv} in (7.5), and the min-max in (7.1) both depend on γ , δ , and m , via $\mathcal{K}(\gamma, \delta, m)$.

There are two important consequences of Theorem 7.5. First, following as in [37, Section 4.6 and Proposition 4.35], this identifies the natural class of extremal operators for J as

$$M_{inv}^+(f, x) = \max_{L \in \mathcal{L}_{inv}} (L(f, x)) \quad \text{and} \quad M_{inv}^-(f, x) = \min_{L \in \mathcal{L}_{inv}} (L(f, x)). \quad (7.6)$$

These extremal operators are defined specifically to produce, for all $f, g \in \mathcal{K}(\gamma, \delta, m)$, the inequalities:

$$M_{inv}^-(f - g, x) \leq J(f, x) - J(g, x) \leq M_{inv}^+(f - g, x). \quad (7.7)$$

We note that the translation invariance shows that

$$\forall x \in \mathbb{R}^d, \quad M_{inv}^\pm(f, x) = M_{inv}^\pm(\tau_x f, 0).$$

Second, the translation invariance of J shows that all of the desired properties of \mathcal{L}_{inv} can be obtained from studying the (nonlinear) functional

$$j : \mathcal{K}(\gamma, \delta, m) \rightarrow \mathbb{R} \quad \text{with} \quad j(f) := J(f, 0).$$

This functional, j , also satisfies the GCP based at $x_0 = 0$ (Definition 2.2). As will be shown in the remainder of this subsection, this means that once J is fixed, the class \mathcal{L} can be restricted even further so that one uses only those L in the Clarke differential of j , i.e. those L such that

$$L(f, x) = \ell(\tau_x f), \quad \text{for some choice of } \ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)},$$

where we recall in Definition 2.6 the notation, $[\partial j]_{\mathcal{K}(\gamma, \delta, m)}$. Thus, we can define a different extremal operator, depending explicitly on J and $\mathcal{K}(\gamma, \delta, m)$ as

$$M_{J, \mathcal{K}(\gamma, \delta, m)}^+(f, x) = \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (\ell(\tau_x f)) \quad \text{and} \quad M_{J, \mathcal{K}(\gamma, \delta, m)}^-(f, x) = \min_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (\ell(\tau_x f)), \quad (7.8)$$

and the inequalities in (7.7) still hold. We note, by definition, that $M_{J, \mathcal{K}(\gamma, \delta, m)}^\pm$ are translation invariant. It is also true that these $M_{J, \mathcal{K}(\gamma, \delta, m)}^\pm$ serve as extremal operators for J , which we record here.

Proposition 7.6. *If J is fixed, and $M_{J, \mathcal{K}(\gamma, \delta, m)}^\pm$ are defined in (7.8), then J and $M_{J, \mathcal{K}(\gamma, \delta, m)}^\pm$ also obey the inequalities in (7.7) with M_{inv}^\pm replaced by $M_{J, \mathcal{K}(\gamma, \delta, m)}^\pm$. Furthermore, $M_{J, \mathcal{K}(\gamma, \delta, m)}^\pm$ are Lipschitz functions, as mappings of $C^{1, \gamma}(\mathbb{R}^d) \rightarrow C^0(\mathbb{R}^d)$ with a Lipschitz norm bounded by that of J .*

Proof of Proposition 7.6. This proof is a direct consequence of the fact that $[\partial j]_{\mathcal{K}(\gamma, \delta, m)}$ is non-empty ([29, Proposition 2.1.2]), combined with a mean value property enjoyed by j and ∂j , due to Lebourg (see, e.g. [29, Theorem 2.3.7]). It says that given any $f, g \in \mathcal{K}(\gamma, \delta, m)$, there exists an element, $\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}$, so that

$$j(f) - j(g) = \ell(f - g).$$

Hence, taking a max over $[\partial j]_{\mathcal{K}(\gamma, \delta, m)}$, (we note as in [29, Proposition 2.1.2], $[\partial j]_{\mathcal{K}(\gamma, \delta, m)}$ is weak-* compact)

$$\forall f, g \in \mathcal{K}(\gamma, \delta, m), \quad j(f) - j(g) \leq \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (\ell(f - g)).$$

A similar argument produces the lower inequality in (7.7). The Lipschitz nature of $M_{J, \mathcal{K}(\gamma, \delta, m)}^{\pm}$ was already apparent from the original invocation of [37, Theorem 1.6, Proposition 1.7], which appeared in (7.1)–(7.3); correspondingly, this implies that the bounds on the linear operators are uniform over these ingredients. \square

Finally, we give the one last argument for the proof of Theorem 7.5.

Proof of Theorem 7.5. Following that last step of the proof of Proposition 7.6, we see that

$$\forall f, g \in \mathcal{K}(\gamma, \delta, m), \quad j(f) \leq \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (j(g) + \ell(f - g)).$$

Now, taking a minimum over g , gives

$$j(f) \leq \min_{g \in \mathcal{K}(\gamma, \delta, m)} \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (j(g) + \ell(f - g)),$$

and evaluating when $g = f$, shows

$$\min_{g \in \mathcal{K}(\gamma, \delta, m)} \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (j(g) + \ell(f - g)) \leq j(f).$$

Now, recalling the fact that J is translation invariant, we see that

$$J(f, x) = j(\tau_x f),$$

and now the theorem follows from that fact that any $\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}$ also enjoys the comparison principle, and thus must be of the form (7.5), per Proposition 7.4. \square

Corollary 7.7. *For each γ, δ , and m fixed, the operator, J , can be extended– with respect to the set, $\mathcal{K}(\gamma, \delta, m)$ – to a function on all of $C^{1, \gamma}(\mathbb{R}^d)$, instead of just $\mathcal{K}(\gamma, \delta, m) \subset C^{1, \gamma}$, and this extension is still Lipschitz and enjoys the GCP. Specifically, if we define*

$$\tilde{J}_{\mathcal{K}(\gamma, \delta, m)} : C^{1, \gamma}(\mathbb{R}^d) \rightarrow C^0(\mathbb{R}^d) \quad \text{via} \quad \tilde{J}_{\mathcal{K}(\gamma, \delta, m)}(f, x) = \min_{g \in \mathcal{K}(\gamma, \delta, m)} \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma, \delta, m)}} (j(g) + \ell((\tau_x f) - g)), \quad (7.9)$$

then \tilde{J} is Lipschitz, $\tilde{J} = J$ on $\mathcal{K}(\gamma, \delta, m)$, and \tilde{J} enjoys the GCP.

Corollary 7.8. *If x is fixed, γ, δ, m are given, and f is m - $C^{1, \gamma}(x)$, then $\tilde{J}_{\mathcal{K}(\gamma, \delta, m)}(f, x)$ is classically defined via (7.9).*

Proof of Corollary 7.8. If f is m - $C^{1, \gamma}(x)$, then this means that there are two functions, f^+ and f^- , such that $f^{\pm} \in C^{1, \gamma}(\mathbb{R}^d)$ and for all y , $f^-(y) \leq f(y) \leq f^+(y)$, with $f^-(x) = f(x) = f^+(x)$. (Note, these f^{\pm} would of course depend on the parameter, r , in Definition 5.2, but that is irrelevant.) Hence, the formula in (7.9) holds classically by Proposition 7.4 and the estimate on μ that appears in (7.3). \square

An important consequence of the Assumption 7.1-(vi) is that it shows that $c^{ij} \leq 0$, which we state and prove here.

Proposition 7.9. *If J is as in Assumption 7.1 and $M_{J,\mathcal{K}(\gamma,\delta,m)}^+$ is the extremal operator defined in (7.8), then*

$$\forall x \in \mathbb{R}^d, \quad M_{J,\mathcal{K}(\gamma,\delta,m)}^+(1, x) \leq 0.$$

As a consequence, we see that necessarily for all $L \in \mathcal{L}_{inv}$ in (7.5), we have $c \leq 0$.

Proof. By the translation invariance of $M_{J,\mathcal{K}(\gamma,\delta,m)}^+$, it suffices to show that

$$M_{J,\mathcal{K}(\gamma,\delta,m)}^+(1, 0) \leq 0.$$

Let us write m_0 to denote the Lipschitz functional $m_0(f) = M_{J,\mathcal{K}(\gamma,\delta,m)}^+(f, 0)$ (the Lipschitz character is part of Proposition 7.6), and recall that

$$m_0(f) = M_{J,\mathcal{K}(\gamma,\delta,m)}^+(f, 0) = \max_{\ell \in [\partial j]_{\mathcal{K}(\gamma,\delta,m)}} \ell(f) \quad \text{where } j \text{ is } j(f) = J(f, 0).$$

Assumption 7.1-(vi) will show that all $\ell \in [\partial j]_{\mathcal{K}(\gamma,\delta,m)}$ will have a sign for $\ell(1)$. Indeed, if $f \in C^{1,\gamma}$, then by Assumption 7.1-(vi), we have

$$\limsup_{s \rightarrow 0^+} \frac{j(f + s \cdot 1) - j(f)}{s} \leq 0.$$

From Definition 2.6 for $[\partial j]_{\mathcal{K}(\gamma,\delta,m)}$, we conclude that

$$\text{if } \ell \in [\partial j]_{\mathcal{K}(\gamma,\delta,m)}, \text{ then } \ell(1) \leq 0.$$

Thus, via the definition of $M_{J,\mathcal{K}(\gamma,\delta,m)}^+$ in (7.8), we see that $M_{J,\mathcal{K}(\gamma,\delta,m)}^+(1, 0) = m_0(1) \leq 0$, as we wanted. \square

Finally, to end this section, we will mention how the results of Sections 5 and 6 ensure that the one-and-two phase operators are covered by the assumptions of this section.

Theorem 7.10. *The operators, I and H , defined respectively in (5.2) and (6.5) satisfy Assumption 7.1. In the case of H , all of the constants and Lipschitz norms will depend on $\mathcal{K}(\gamma, \delta, m, L)$, instead of just $\mathcal{K}(\gamma, \delta, m)$ (where L is the height of the top boundary in Section 6). Furthermore, the same is true as well for the operators, for respectively the one-phase and two-phase evolutions:*

$$G(I(f)) \cdot \sqrt{1 + |\nabla f|^2} \quad \text{and} \quad H(f) \cdot \sqrt{1 + |\nabla f|^2}.$$

Proof. For I , this is a direct consequence of the results in Theorem 5.13, Lemma 5.6, Proposition 5.7, Lemma 5.15, and Lemma 5.22. Hence for the map $G(I(f))$, this follows from the Lipschitz nature of G . For H , this follows immediately from the Lipschitz nature of G , combined with the observation of Lemma 6.1. Indeed, I^+ already has the desired properties (as in Section 5), and by Lemma 6.1 I^- does as well. Everything can be checked for H with a calculation similar to that of the proof of Theorem 6.2.

Finally, for the cases of

$$G(I(f)) \cdot \sqrt{1 + |\nabla f|^2} \quad \text{and} \quad H(f) \cdot \sqrt{1 + |\nabla f|^2},$$

we note that the mapping,

$$f \mapsto \sqrt{1 + |\nabla f|^2},$$

is bounded and Lipschitz on each of the sets, $\mathcal{K}(\gamma, \delta, m, L)$. This, combined with the local nature of $\sqrt{1 + |\nabla f|^2}$, is enough to preserve all of the assumptions listed in 7.1, as soon as I or H satisfy them as well. \square

8. COMPARISON THEOREM FOR PARABOLIC VISCOSITY SOLUTIONS

This section is dedicated to the uniqueness and existence of viscosity solutions for the class of fractional parabolic equations that contain those in Assumption 7.1. As we just saw in Theorem 7.10, this contains the operators $G(I(f)) \cdot \sqrt{1 + |\nabla f|^2}$ and $H(f) \cdot \sqrt{1 + |\nabla f|^2}$ for the free boundary flow. We want to emphasize that this section should be viewed as a collection of small modifications to the arguments found in the paper of Silvestre, [57, Appendix A], for some similar integro-differential equations. At the end of this section, we will show how a similar comparison result can be deduced from Theorem 5.13, Lemma 5.11, and Corollary 5.24.

8.1. Using the assumptions of Section 7. In this part, we show the uniqueness for parabolic equations governed by general operators satisfying the assumptions of Section 7. For such J that satisfy Assumption 7.1, we prove uniqueness for

$$\begin{cases} \partial_t f = J(f) & \text{in } \mathbb{R}^d \times (0, T) \\ f(\cdot, 0) = f_0 & \text{on } \mathbb{R}^d \times \{0\}. \end{cases} \quad (8.1)$$

Remark 8.1. *We present all the proofs in a fashion that applies to the one-phase evolution, using the operator, e.g. $J(f) = G(I(f)) \cdot \sqrt{1 + |\nabla f|^2}$. This is apparent in the need to reference sets such as $\mathcal{K}(\gamma, \delta, m)$, and the dependence of arguments on the parameters, δ and m . All of these results hold for the two-phase problem, e.g. $J(f) = H(f) \cdot \sqrt{1 + |\nabla f|^2}$ as well, with the additional constraint that constants and arguments will also depend upon L and $\mathcal{K}(\gamma, \delta, m, L)$.*

It turns out that since J is translation invariant, the viscosity solutions theory is well developed and straightforward to implement. The key step is the regularization via the operations of inf-convolution and sup-convolution (also known earlier as the Moreau-Yosida approximation on Hilbert spaces). These operations were expanded upon by Lasry-Lions [50]; first utilized for uniqueness of elliptic equations by Jensen in [44]; and improved upon by Ishii in [42]. A good demonstration of their utility appears in the work of Jensen-Lions-Souganidis [43]. As mentioned above, most of the arguments that we need that pertain to the nonlocal terms come nearly verbatim from the work of Silvestre in [57, Appendix A], and we will refer most of the details to that work. We begin by recognizing the fact that for equations whose order is strictly less than 2, the inf/sup-convolution technique works even more seamlessly than in the case of second order equations. This is due to the fact that a function with $C^{1,1}$ regularity in space is good enough to classically evaluate J . Such benefits of working with sub and super solutions that can be evaluated classically can be seen in Caffarelli-Silvestre [12, Section 5] and Barles-Imbert [6], as well as in [57]. For a general overview of viscosity solutions, we suggest the User's Guide [33], and for an introduction of these techniques in the context of parabolic equations, we suggest the lecture notes of Imbert-Silvestre [40].

First, we must recall the definition of a viscosity solution of the parabolic equation. For our context, this is equivalent to any of the definitions that appear in Barles-Imbert [6], Caffarelli-Silvestre [12], or Silvestre [57].

Definition 8.2. *We say that f is a viscosity subsolution of (8.1) if f is upper semi-continuous, that $\inf f > 0$, and f has the property that for all $(x, t) \in \mathbb{R}^d \times (0, T)$ for which there exists a function, ϕ , so that for some $\beta > 0$ with $\gamma + \beta < 1$, $\phi \in C^{1, \gamma + \beta}(\mathbb{R}^d)$ in space and C^1 in time, and for some $t > r > 0$, $f - \phi$ attains a global maximum over $\mathbb{R}^d \times (t - r, t)$, then ϕ must satisfy*

$$\partial_t \phi(x, t) \leq J(\phi, x).$$

We say that g is a viscosity supersolution if g is lower semi-continuous, $\inf g > 0$, and the above properties are replaced by $g - \phi$ attains a minimum for $\inf \phi > 0$, and

$$\partial_t \phi(x, t) \geq J(\phi, x).$$

We say that f is a viscosity solution if f is both a subsolution and a supersolution.

Remark 8.3. We note that we have chosen the space of test functions to have the natural regularity in space that is associated to Assumption 7.1; this is instead of the usual C^2 regularity that typically appears in other works that use viscosity solutions for integro-differential equations.

Remark 8.4. We note that definition 8.2 is not identically the one in [57, Section 2], however, it is equivalent, as evidenced in Barles-Imbert [6].

Thanks to the results in Section 7, it is easy to show that the definition of viscosity solutions extends to a class of functions with less regularity in space, namely test functions that, for some $\beta > 0$ with $\gamma + \beta < 1$, are punctually $C^{1,\gamma+\beta}$ in space at the point of contact will produce the same set of solutions.

Lemma 8.5. (This is the analog of the result of, e.g. [12, Lemma 4.3].) If f is a viscosity subsolution of (8.1) and $(x, t) \in \mathbb{R}^d \times (0, T)$ is a point such that $f - \phi$ attains a maximum at (x, t) , for some ϕ that is punctually $C^{1,\gamma+\beta}(x)$ in space and C^1 in time, with $\beta > 0$ and $\gamma + \beta < 1$, then there exists a choice of δ_0 small enough and m_0 large enough (depending on ϕ), so that $\tilde{J}_{\mathcal{K}(\gamma, \delta_0, m_0)}(\phi, x)$ is defined classically and

$$\partial_t \phi(x, t) \leq \tilde{J}_{\mathcal{K}(\gamma, \delta_0, m_0)}(\phi, x).$$

(\tilde{J} is defined in (7.9) and its appearance is because it is not required that ϕ is in the set, $\mathcal{K}(\gamma, \delta_0, m_0)$.) The analogous result holds for g that are supersolutions and for those ϕ that also additionally satisfy $\inf \phi > 0$. (The inequality also holds for $\delta < \delta_0$ and $m > m_0$.)

Proof. As we mentioned in the statement of the lemma, this is basically the result that is presented in [12, Lemma 4.3]. The proof is also very similar. We provide some details for the sake of readability. First of all, the well defined nature of $\tilde{J}_{\mathcal{K}(\gamma, \delta, m)}(\phi, x)$ is a consequence of Corollary 7.8, from which it is useful to note does not require the extra $\gamma + \beta$ regularity of ϕ at x . The extra regularity is required to ensure that $\tilde{J}_{\mathcal{K}(\gamma, \delta, m)}(\phi, x)$ actually obeys the correct inequality.

We observe that by the assumption on ϕ , there exist functions, f^+ , f^- , both in $C^{1,\gamma+\beta}(\mathbb{R}^d)$, and that satisfy

$$\forall y \in \mathbb{R}^d, \quad f^-(y) \leq \phi(y) \leq f^+(y), \quad \text{and} \quad f^-(x) = \phi(x) = f^+(x).$$

For each $r > 0$, we can define the function, ϕ_r , as

$$\phi_r(y) = \begin{cases} f^+(y) & \text{if } y \in B_r(x) \\ \phi(y) & \text{otherwise.} \end{cases}$$

Furthermore, we note that we also have the ordering

$$f^- \leq \phi \leq \phi_r \leq f^+,$$

and that

$$\nabla f^-(x) = \nabla \phi(x) = \nabla \phi_r(x) = \nabla f^+(x).$$

Finally, we can assume without loss of generality that

$$\partial_t \phi(x, t) = \partial_t f^+(x, t).$$

The definition of viscosity solution shows that

$$\partial_t \phi(x, t) = \partial_t f^+(x, t) \leq J(f^+, x).$$

Now we can use Corollaries 7.7 and 7.8, as well as the relevant extremal operators to see that since there exists some δ_0 and m_0 so that $f^+ \in \mathcal{K}(\gamma, \delta_0, m_0)$, which we simply call $\mathcal{K}(\gamma, \delta_0, m_0) = \mathcal{K}$,

$$\begin{aligned} J(f^+, x) &= \tilde{J}_{\mathcal{K}}(f^+, x) \leq \tilde{J}_{\mathcal{K}}(\phi_r, x) + M_{J, \mathcal{K}}^+(f^+ - \phi_r, x) \\ &\leq \tilde{J}_{\mathcal{K}}(\phi, x) + M_{J, \mathcal{K}}^+(\phi_r - \phi, x) + M_{J, \mathcal{K}}^+(f^+ - \phi_r, x). \end{aligned}$$

Again, we recall that all of the operators involving $\tilde{J}_{\mathcal{K}}$ and $M_{J, \mathcal{K}}^+$ are well defined on each of these functions, as they are all punctually $C^{1,\gamma}(x)$. The only question now is whether or not the last two

terms vanish as $r \rightarrow 0$. This is where the slightly higher regularity is used. Indeed, because the set, \mathcal{K} , is now fixed, and thanks to the bounds in Theorem 7.5, as well as the definition of $M_{J,\mathcal{K}}^+$, we see that all of these terms include a measure μ and an integral that are dominated by

$$\int_{B_r} C_\phi |y|^{1+\gamma+\beta} \mu(dy),$$

where C_ϕ is a constant depending the fact that $\phi \in C^{1,\gamma+\beta}(x)$. Furthermore, we know that $M_{J,\mathcal{K}}^+$ is a max over a family of these μ , all of the μ (arising from the set, $[J]_{\mathcal{K}}$), satisfy the uniform bound (7.3). For simplicity, let us call $meas(\mathcal{K})$ to be the set of all measures, μ , so that there exists $\ell \in [J]_{\mathcal{K}}$ so that μ is the corresponding measure for ℓ as in Proposition 7.4. Hence, the remaining terms above, that depend on r , are all bounded by

$$\int_{B_r} C_\phi |y|^{1+\gamma+\beta} \mu(dy) \leq C_\phi r^\beta \sup_{\mu \in meas(\mathcal{K})} \int_{\mathbb{R}^d} \min\{|y|^{1+\gamma}, 1\} \mu(dy) \leq C_2 \cdot C_\phi \cdot r^\beta.$$

Hence, taking $r \rightarrow 0$ shows that $M_{J,\mathcal{K}}^+(\phi_r - \phi) + M_{J,\mathcal{K}}^+(f^+ - \phi) \rightarrow 0$, and this finishes the claim of the lemma. \square

Following either [40] or [57], we see that the inf and sup convolutions that are appropriate for parabolic equations are the following:

$$\phi^\varepsilon(x, t) = \sup_{y \in \mathbb{R}^d, s \in [0, \infty)} \phi(y, s) - \frac{1}{2\varepsilon}(t-s)^2 - \frac{1}{2\varepsilon}|x-y|^2 \quad (8.2)$$

$$\phi_\varepsilon(x, t) = \inf_{y \in \mathbb{R}^d, s \in [0, \infty)} \phi(y, s) + \frac{1}{2\varepsilon}(t-s)^2 + \frac{1}{\varepsilon}|x-y|^2. \quad (8.3)$$

We will also need to know how the family of operators as in Assumption 7.1 operate on rescaled versions of smooth bump functions. In particular, it is important to know that the following.

Lemma 8.6. *If Φ and Φ_R are the smooth functions defined as*

$$\Phi(x) = \frac{|x|^2}{1+|x|^2}, \quad \text{and} \quad \Phi_R(x) = \Phi\left(\frac{x}{R}\right),$$

then, for a fixed J , given any $\delta > 0$, $m > \delta$, $\rho > 0$, there exists $R > 1$, with $R = R(J, \rho, \delta, m)$, so that

$$\sup_{x \in \mathbb{R}^d} M_{J,\mathcal{K}(\gamma, \delta, m)}^+(\Phi_R, x) \leq \rho.$$

Proof of Lemma 8.6. First, for a fixed $t > 1$ to be made precise later, we collect a few facts about Φ_R :

$$\begin{aligned} 0 \leq \Phi_R(x) \leq 1 \quad \text{and} \quad \|\nabla \Phi_R\|_{L^\infty} &\leq \frac{C}{R} \\ |\Phi_R(x+h) - \Phi_R(x) - \mathbb{1}_{B_1}(h) \nabla \Phi_R(x) \cdot h| \mathbb{1}_{B_1}(h) &\leq \frac{C|h|^2}{R^2}, \\ |\Phi_R(x+h) - \Phi_R(x) - \mathbb{1}_{B_1}(h) \nabla \Phi_R(x) \cdot h| \mathbb{1}_{B_t \setminus B_1}(h) &\leq \frac{Ct}{R} \\ |\Phi_R(x+h) - \Phi_R(x) - \mathbb{1}_{B_1}(h) \nabla \Phi_R(x) \cdot h| \mathbb{1}_{\mathbb{R}^d \setminus B_t}(h) &\leq 1. \end{aligned}$$

Now, using the uniform estimates on the Lévy measures, μ , we can first choose t large enough according to Theorem 7.5, so that, uniformly across $[\partial j]_{\mathcal{K}(\gamma, \delta, m)}$,

$$\int_{\mathbb{R}^d \setminus B_t} |\Phi_R(x+h) - \Phi_R(x) - \mathbb{1}_{B_1}(h) \nabla \Phi_R(x) \cdot h| \mu(dh) \leq \frac{\rho}{3}.$$

Next, given this t , we can choose R large enough so that uniformly across b from $[\partial j]_{\mathcal{K}(\gamma, \delta, m)}$

$$b \cdot \nabla \Phi_R \leq \frac{\rho}{3}$$

and

$$\int_{B_t} |\Phi_R(x+h) - \Phi_R(x) - \mathbb{1}_{B_1}(h) \nabla \Phi_R(x) \cdot h| \mu(dh) \leq \frac{\rho}{3}.$$

Finally, thanks to Proposition 7.9, we see that for any of the constants, c , appearing in (7.5), we have

$$c\Phi_R(x) \leq 0.$$

Hence, by the definition of $M_{J, \mathcal{K}(\gamma, \delta, m)}^+$ we have obtained the desired inequality. \square

The previous lemma allows us to construct a smooth strict supersolution, that also acts a bump function.

Lemma 8.7. *Given any $\delta > 0$, $m > \delta$, $\rho > 0$, and $C > 0$ there exists a choice of $R = R(J, \rho, \delta, m)$ so that for all $h > 0$,*

$$\forall (x, t) \in \mathbb{R}^d \times [0, T], \quad \Psi(x, t) = C + h\Phi_R(x) + h\rho t,$$

is a classical, strict super solution of

$$\partial_t \Psi > M_{J, \mathcal{K}(\gamma, \delta, m)}^+(\Psi).$$

Proof of Lemma 8.7. This is a direct calculation, but it is essential to invoke Proposition 7.9 to ensure the desired inequality. In particular, Proposition 7.9, or rather that $c \leq 0$ in the definition of $M_{J, \mathcal{K}(\gamma, \delta, m)}^+$, show that since $C \geq 0$,

$$M_{J, \mathcal{K}(\gamma, \delta, m)}^+(C + h\Phi_R) \leq hM_{J, \mathcal{K}(\gamma, \delta, m)}^+(\Phi_R).$$

Thus, with ρ given, we can invoke Lemma 8.6 with $(\rho/2)$ to get the desired inequality. \square

The first comparison result is for subsolutions of an extremal equation. Although we do not invoke this result in the same way as in [57], we think it is useful to have as a tool. The result that is truly essential to this work is the next one, which is Proposition 8.9.

Lemma 8.8. *If $\delta > 0$, $m > \delta$ are fixed, and $w : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$ is a bounded, upper semi-continuous function such that, in the viscosity sense*

$$\partial_t w \leq M_{J, \mathcal{K}(\gamma, \delta, m)}^+(w),$$

then, $\sup_{\mathbb{R}^d \times [0, T]} w_+ \leq \sup_{\mathbb{R}^d} w_+(\cdot, 0)$

Proof. First, we will provide the proof when $\sup_{\mathbb{R}^d} w(\cdot, 0) = 0$. Given the result of Lemma 8.7, we see that the proof of [57, Lemma 3.3] applies directly. Indeed if we assume, for the sake of contradiction, $\sup_{\mathbb{R}^d \times [0, T]} w > 0$, then for some choice of $C \geq 0$, and $h > 0$, $w - \Psi$ (where Ψ is as in Lemma 8.7) will attain a global max for $t > 0$. The definition of viscosity solution applied to Ψ as a test function gives a contradiction.

Next, if we do not necessarily assume that $\sup_{\mathbb{R}^d} w(\cdot, 0) = 0$, we can replace w by the function,

$$\tilde{w} = w - \sup_{\mathbb{R}^d} w(\cdot, 0).$$

Thus, if $c = \sup_{\mathbb{R}^d} w(\cdot, 0) \geq 0$, we see that $w = \tilde{w} + c$, and so in the viscosity sense, $\partial_t w = \partial_t \tilde{w}$. Furthermore, since $c \geq 0$, by Assumption 7.1 (vi) and Proposition 7.9, in the viscosity sense,

$$M_{J, \mathcal{K}(\gamma, \delta, m)}^+(w) = M_{J, \mathcal{K}(\gamma, \delta, m)}^+(\tilde{w} + c) \leq M_{J, \mathcal{K}(\gamma, \delta, m)}^+(\tilde{w}).$$

Hence, in the viscosity sense,

$$\partial_t \tilde{w} = \partial_t w \leq M_{J, \mathcal{K}(\gamma, \delta, m)}^+(w) \leq M_{J, \mathcal{K}(\gamma, \delta, m)}^+(\tilde{w}),$$

and the first case is applicable. \square

Proposition 8.9. *If $f, g : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$ are bounded and, with for some $\delta > 0$, $f, g \geq \delta$, are respectively a sub and a supersolution, in the viscosity sense, of*

$$\partial_t u = J(u),$$

and $f(x, 0) \leq g(x, 0)$ for all x , then,

$$f(x, t) \leq g(x, t) \quad \forall (x, t) \in \mathbb{R}^d \times [0, T].$$

Proof of Proposition 8.9. We first note that thanks to the definition of viscosity solution and the translation invariance of J , that if we define the sup-convolution, f^ε , and the inf-convolution, g_ε , then the following two inequalities are satisfied in the viscosity sense:

$$\partial_t f^\varepsilon \leq J(f^\varepsilon), \quad \text{and} \quad \partial_t g_\varepsilon \geq J(g_\varepsilon).$$

We let $\varepsilon \in (0, 1)$ be fixed. We will assume for the sake of contradiction that there exists a $t^* \in (0, T)$ such that

$$0 < m_0 = \sup_{(x, t) \in \mathbb{R}^d \times [0, T]} (f(x, t) - g(x, t)) = \sup_{x \in \mathbb{R}^d} (f(x, t^*) - g(x, t^*)).$$

Now, since ε is fixed, and m_0 is given, it is possible to invoke Lemma 8.7 with $m = c\varepsilon^{-1}$ (and $C = cm_0$), in order to ensure the existence of Ψ such that the following inequality holds classically for all $(x, t) \in \mathbb{R}^d \times [0, T]$:

$$\partial_t \Psi > M_{J, \mathcal{K}(\gamma, \delta/2, c\varepsilon^{-1})}^+(\Psi).$$

Furthermore, by construction, C can be chosen so that there exists $(x_0^\varepsilon, t_0^\varepsilon) \in \mathbb{R}^d \times (0, T]$ so that

$$f^\varepsilon - g_\varepsilon \leq \Psi, \quad \text{and} \quad (f^\varepsilon - g_\varepsilon)(x_0^\varepsilon, t_0^\varepsilon) = \Psi(x_0^\varepsilon, t_0^\varepsilon).$$

By definition, both f^ε and $-g_\varepsilon$ are semi-convex in space and time. Thus, since $f^\varepsilon - g_\varepsilon$ is touched from above by Ψ at $(x_0^\varepsilon, t_0^\varepsilon)$, we have that $f^\varepsilon, g_\varepsilon \in (c\varepsilon^{-1})\text{-}C^{1,1}(x_0^\varepsilon)$. This also implies $f^\varepsilon, g_\varepsilon \in (c\varepsilon^{-1})\text{-}C^{1,\gamma}(x_0^\varepsilon)$, and that $\partial_t f^\varepsilon, \partial_t g_\varepsilon, \nabla f^\varepsilon, \nabla g_\varepsilon$ all exist at $(x_0^\varepsilon, t_0^\varepsilon)$. (Here we have implicitly used that without loss of generality, $\Psi \in (c\varepsilon^{-1})\text{-}C^{1,1}(x_0^\varepsilon)$) Thus, by Lemma 8.5 and Corollary 5.12, we see that the equations hold classically for f^ε and g_ε . Furthermore, by Proposition 7.6 and Lemma 8.5, we see that classically, (for $c = \|f\|_{L^\infty} + \|g\|_{L^\infty}$)

$$\begin{aligned} \partial_t f^\varepsilon - \partial_t g_\varepsilon &\leq \tilde{J}_{\mathcal{K}(\gamma, \delta, c\varepsilon^{-1})}(f^\varepsilon, x_0^\varepsilon) - \tilde{J}_{\mathcal{K}(\gamma, \delta, c\varepsilon^{-1})}(g_\varepsilon, x_0^\varepsilon) \\ &\leq M_{J, \mathcal{K}(\gamma, \delta, c\varepsilon^{-1})}^+(f^\varepsilon - g_\varepsilon, x_0^\varepsilon) \leq M_{J, \mathcal{K}(\gamma, \delta, c\varepsilon^{-1})}^+(\Psi, x_0^\varepsilon), \end{aligned}$$

where the last inequality resulted from the fact that $M_{J, \mathcal{K}(\gamma, \delta, c\varepsilon^{-1})}^+$ enjoys the GCP. Since at $(x_0^\varepsilon, t_0^\varepsilon)$ we know that $\partial_t(f^\varepsilon - g_\varepsilon) = \partial_t \Psi$, this contradicts the fact that

$$\partial_t \Psi > M_{J, \mathcal{K}(\gamma, \delta/2, c\varepsilon^{-1})}^+.$$

Thus, we can conclude that for each ε fixed, we know that

$$\forall (x, t) \in \mathbb{R}^d \times [0, T], \quad f^\varepsilon(x, t) \leq g_\varepsilon(x, t).$$

Taking $\varepsilon \rightarrow 0$ yields the result of the proposition. \square

8.2. Existence. The method of existence will, again, follow [57, Appendix A] (which comes from Ishii's adaptation of Perron's method, e.g. [41], [42], or [33]), provided we can construct certain barrier functions. Indeed, nearly the same barriers from [57] will work, with a minor modification.

Assume that $f_0 : \mathbb{R}^d \rightarrow \mathbb{R}$ is uniformly continuous and $f_0 \geq \delta$. First, we construct two smooth functions to serve as lower and upper initial data. Assume that $b : \mathbb{R}^d \rightarrow \mathbb{R}$, b is smooth, $0 \leq b \leq 1$, $b(0) = 1$, and $\text{support}(b) \subset B_1$.

Fix x_0 . We will produce lower and upper barriers for f_0 at the point x_0 . Define, for $\rho > 0$,

$$b_\rho(x) := b\left(\frac{x - x_0}{\rho}\right).$$

We claim that given any $\varepsilon > 0$, we can produce a choice of ρ so that the following functions will trap f_0 . We define the upper and lower barrier's for f_0 at x_0 as

$$\begin{aligned} U_0(x) &:= b_\rho(x)f_0(x_0) + (1 - b_\rho(x)) \sup_{\mathbb{R}^d}(f_0) + \varepsilon \\ L_0(x) &:= b_\rho(x)f_0(x_0) + (1 - b_\rho(x)) \inf_{\mathbb{R}^d}(f_0) - \varepsilon. \end{aligned}$$

Specifically, given $\varepsilon > 0$, we can find a $\rho > 0$ so that

$$\forall x \in \mathbb{R}^d \quad U_0(x) \geq f_0(x) \quad \text{and} \quad L_0(x) \leq f_0(x).$$

With these barriers in hand, we see that Perron's method for viscosity solutions applies to the equation (8.1). We summarize this as

Theorem 8.10. *If f_0 is uniformly continuous and bounded on \mathbb{R}^d , then there exists a unique viscosity solution, f , that solves (8.1).*

8.3. Comparison using the free boundary operators directly. In this part, we show how the previous results indeed do work, with basic modifications, under the assumptions of the free boundary operators, I and H , above. These arguments will be valid regardless of whether or not one has proved that I has an integro-differential representation as in Section 7.

Theorem 8.11. *If I is as in (5.2) or H as in (6.5), $f, g : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$ are bounded and, for some $\delta > 0$, $f, g \geq \delta$, and for H also satisfy $f, g \leq L - \delta$, are respectively a sub and a supersolution, in the viscosity sense, of*

$$\partial_t u = G(I(u)) \cdot \sqrt{1 + |\nabla u|^2}, \quad \text{or} \quad \partial_t u = H(u) \cdot \sqrt{1 + |\nabla u|^2}$$

and $f(x, 0) \leq g(x, 0)$ for all x , then,

$$f(x, t) \leq g(x, t) \quad \forall (x, t) \in \mathbb{R}^d \times [0, T].$$

The proof proceeds nearly identically to the arguments in Proposition 8.9 and Lemma 8.8, except that we don't invoke the extremal operator, $M_{J, \mathcal{K}(\gamma, \delta, m)}^+$, and we use Corollary 5.24 instead of Lemma 8.7.

First we will demonstrate the case for the equation for the one phase case, $G(I(u))$. To this end, we again take f^ε and g_ε to be the sup-convolution and inf-convolution of f and g . We assume, for the sake of contradiction, that $\sup_{\mathbb{R}^d \times [0, T]} f^\varepsilon - g_\varepsilon > 0$. As above, this means that for some $C \geq 0$, $\rho > 0$, and $h > 0$, there will exist \hat{x} and $\hat{t} > 0$ such that $f^\varepsilon - g_\varepsilon - \Psi$ attains a zero maximum at (\hat{x}, \hat{t}) (Ψ is taken to be as in Lemma 8.7). Furthermore, we can choose the parameters C, ρ, h so that we also maintain

$$\delta \leq g_\varepsilon + \Psi, \quad \text{and} \quad f + \Psi \leq L - \frac{\delta}{2},$$

which is a technical requirement for applying Corollary 5.24; the restriction relative to f^ε and Ψ is only relevant for the case of the operator, H , which we treat second.

This means that at (\hat{x}, \hat{t}) , both f^ε and g_ε have classical derivatives with respect to both x and t at (\hat{x}, \hat{t}) and also that $I(f^\varepsilon, \hat{x})$, $I(g_\varepsilon, \hat{x})$ are classically defined. We can use the Corollary 5.24 to get

a contradiction. Indeed, assume that $s > 0$ is generic. We see that by the Lipschitz nature of G and $\sqrt{1 + |\nabla f|}$,

$$\begin{aligned} & G(I(g_\varepsilon + \Psi))\sqrt{1 + |\nabla g_\varepsilon + \nabla \Psi|^2} - G(I(g_\varepsilon))\sqrt{1 + |\nabla g_\varepsilon|^2} \\ & \leq C \left\| \left(\sqrt{1 + |\nabla g_\varepsilon + \nabla \Psi|^2} \right) \right\|_{L^\infty} \cdot |I(g_\varepsilon + \Psi) - I(g_\varepsilon)| + C \|G(I(g_\varepsilon))\|_{L^\infty} |\nabla \Psi| \end{aligned}$$

Hence for a sufficiently large R , using the definition of Ψ and Corollary 5.24, we can force the right hand side to be controlled by s . We note that this choice of R will depend upon ε , δ , and m , which is suitable since ε is fixed (as in the proof of Proposition 8.9). Thus, at (\hat{x}, \hat{t}) we first use the GCP, followed by the previous inequality to obtain

$$\begin{aligned} \partial_t(g_\varepsilon + \Psi) & \leq \partial_t f^\varepsilon \leq G(I(f^\varepsilon, \hat{x}))\sqrt{1 + |\nabla f^\varepsilon(\hat{x})|^2} \leq G(I(g_\varepsilon + \Psi, \hat{x}))\sqrt{1 + |\nabla g_\varepsilon(\hat{x}) + \nabla \Psi(\hat{x})|^2} \\ & \leq G(I(g_\varepsilon, \hat{x}))\sqrt{1 + |\nabla g_\varepsilon(\hat{x})|^2} + s \leq \partial_t g_\varepsilon + s. \end{aligned}$$

(Here, we note the first inequality is only to include the case that $\hat{t} = T$, otherwise if $\hat{t} < T$, we would have equality.) In other words, this implies that

$$h\rho \leq s.$$

Since any $s > 0$ is admissible, once ρ and h have been fixed, we see, by choosing $s = \frac{1}{2}h\rho$, that this gives a contradiction.

For the case of $H(u) = G(I^+(u), I^-(u))$, we would give nearly the same proof, except that we need to use Corollary 5.24 twice, and we follow the argument for Theorem 6.2. Here is a sketch of the argument, following as above, and the notation for \tilde{f} , \tilde{F}_2 , etc... are as that preceding Lemma 6.1. The term we need to focus on is

$$H(f^\varepsilon, \hat{x}) - H(g_\varepsilon, \hat{x}).$$

Indeed, we have

$$H(f^\varepsilon, \hat{x}) - H(g_\varepsilon, \hat{x}) \leq \Lambda_0(I^+(f^\varepsilon, \hat{x}) - I^+(g_\varepsilon, \hat{x}))_+ + \Lambda_0(-I_{\tilde{F}_2}(L - \tilde{g}_\varepsilon, \hat{x}) + I_{\tilde{F}_2}(L - \tilde{f}^\varepsilon, \hat{x}))_+.$$

At this point, the first term is handled exactly as above, using the fact that f^ε is touched from above at \hat{x} by the function $g^\varepsilon + \Psi$. For the second term, we use the fact that our assumptions on f^ε , g_ε , and Ψ , are such that

$$(L - \tilde{f}^\varepsilon) + (-\tilde{\Psi}) \text{ is touched from below by } (L - \tilde{g}_\varepsilon) \text{ at } \hat{x}.$$

Furthermore, the requirement that $f + \Psi \leq L - \frac{\delta}{2}$ ensures that both $(L - \tilde{f}^\varepsilon)$ and $(L - \tilde{f}^\varepsilon) + (-\tilde{\Psi})$ are in $\mathcal{K}^*(\gamma, \delta/2, m\varepsilon)$ (as f^ε is semi-convex, so $-f^\varepsilon$ is semi-concave). By the GCP,

$$I_{\tilde{F}_2}((L - \tilde{f}^\varepsilon) + (-\tilde{\Psi}), \hat{x}) \geq I_{\tilde{F}_2}(L - \tilde{g}_\varepsilon, \hat{x}),$$

and by Corollary 5.24,

$$I_{\tilde{F}_2}(L - \tilde{f}^\varepsilon, \hat{x}) - I_{\tilde{F}_2}((L - \tilde{f}^\varepsilon) + (-\tilde{\Psi}), \hat{x}) \leq s.$$

These two inequalities are enough to finish the proof as in the first case.

9. FREE BOUNDARY VISCOSITY SOLUTIONS

In this section we study various notions of viscosity solutions of free boundary problems like (1.1). Because we have in mind a one-to-one correspondence between viscosity solutions of the free boundary problem and viscosity solutions of a parabolic equation, we will introduce a definition of viscosity solution that is different from the standard one – as presented in, e.g. [5], [14], [46], [47]. We will then show that our definition contains the existing ones when the assumptions overlap.

9.1. Viscosity solutions for the free boundary evolution (1.1). By a *test interface* in $[a, b]$, we shall mean the following: a smooth hypersurface in $S \subset \Sigma_0 \times [a, b]$ such that each time slice $S(t)$ separates Σ_0 ($\Sigma_0 = \mathbb{R}_+^{d+1}$ or $\Sigma_0 = \mathbb{R}^d \times [0, L]$) into two connected components, that is $\Sigma_0 \setminus S(t) = S(t)^+ \cup S(t)^-$ where $S(t)^+$ and $S(t)^-$ are open sets, $S(t)$ is a positive distance away from $\partial\Sigma_0$, and

$$\begin{aligned} \Gamma_0 &\subset \overline{S(t)^+} && \text{(one phase case),} \\ \Gamma_0 &\subset \overline{S(t)^+}, \Gamma_L \subset \overline{S(t)^-} && \text{(two phase case).} \end{aligned}$$

Given a test interface S in $[a, b]$, we define $U_S : \Sigma_0 \times [a, b] \rightarrow \mathbb{R}$ as the function which for each time $t \in [a, b]$ is the unique solution to one of the following Dirichlet problems; in the two-phase case ($\Sigma_0 = \mathbb{R}^d \times (0, L)$), it solves the equation,

$$\begin{cases} F_1(D^2U_S, \nabla U_S) = 0 & \text{in } S^+, \\ F_2(D^2U_S, \nabla U_S) = 0 & \text{in } S^-, \\ U_S = 0 & \text{on } S, \\ U_S = -1 & \text{on } \Gamma_L, \\ U_S = 1 & \text{on } \Gamma_0, \end{cases} \quad (9.1)$$

and in the one-phase case ($\Sigma_0 = \mathbb{R}_+^{d+1}$), it solves the equation,

$$\begin{cases} F(D^2U_S, \nabla U_S) = 0 & \text{in } S^+, \\ U_S = 0 & \text{on } S, \\ U_S = 1 & \text{on } \Gamma_0, \end{cases} \quad (9.2)$$

Let us recall the definition of classical subsolutions and supersolutions.

Definition 9.1. Let $U : \Sigma_0 \times [a, b] \rightarrow \mathbb{R}$. This function is said to be a *classical subsolution* (respectively *supersolution*) of (1.1) if

- (1) The set $\partial\{U > 0\}$ is a codimension 1 differentiable submanifold of $\Sigma_0 \times [a, b]$ such that each time slice $\partial\{U > 0\} \cap \Sigma_0 \times \{t\}$ ($a \leq t \leq b$) is a codimension 1 differentiable submanifold of Σ_0 , and U is twice differentiable in space and differentiable in time in $\Sigma_0 \times [a, b] \setminus \{U \neq 0\}$.
- (2) For every fixed t , the function $U = U(\cdot, t)$ solves, in the viscosity sense,

$$\begin{aligned} F_1(D^2U, \nabla U) &\geq 0 \text{ in } \{U > 0\} \text{ (resp. } \leq), \\ F_2(D^2U, \nabla U) &\geq 0 \text{ in } \{U < 0\} \text{ (resp. } \leq), \end{aligned}$$

and if V denotes the normal velocity of $\partial\{U > 0\}$ (in the outer normal direction), then

$$V \leq G(\partial_\nu^+ U, \partial_\nu^- U) \text{ along } \partial\{U > 0\} \text{ (resp. } \geq).$$

Furthermore, U is said to be a *classical solution* if it is both a subsolution and a supersolution.

The definition for the one-phase is entirely analogous and we omit it.

Definition 9.2. A function U is said to be *touched from above* at (X_0, t_0) by S if S is a test interface in $[t_0 - \tau, t_0]$ for some $\tau > 0$ such that

$$\{U > 0\} \cap \{t_0 - \tau \leq t \leq t_0\} \subset \overline{S^+} \text{ and } (X_0, t_0) \in \partial\{U > 0\} \cap S.$$

The notion of viscosity solution we will be using is the following.

Definition 9.3. A *viscosity subsolution* (respectively *supersolution*) of (1.1) in $[a, b]$ is an upper semi-continuous function (resp. lower semi continuous function)

$$U : \Sigma_0 \times [a, b] \rightarrow \mathbb{R},$$

which is required to have the following properties. It satisfies the pointwise bounds

$$\begin{aligned} U &\leq 1 \text{ (resp. } U \geq 1) \text{ on } \Gamma_0, U \leq -1; \text{ (resp. } U \geq -1) \text{ on } \Gamma_L \text{ (two-phase case),} \\ U &\leq 1 \text{ (resp. } U \geq 1) \text{ on } \Gamma_0 \text{ (one-phase case);} \end{aligned}$$

and satisfies the following relations in the viscosity sense: in the two-phase case they are

$$\begin{aligned} F_1(D^2U, \nabla U) &\geq 0 \text{ in } \{U > 0\}^0 \text{ (resp. } \leq), \\ F_2(D^2U, \nabla U) &\geq 0 \text{ in } \{U < 0\}^0 \text{ (resp. } \leq), \end{aligned}$$

and in the one-phase case they are

$$F(D^2U, \nabla U) \geq 0 \text{ in } \{U > 0\}^0 \text{ (resp. } \leq).$$

Last but not least, for any test interface S touching U from above at $(X_0, t_0) \in \partial\{U > 0\}$ we have

$$\begin{aligned} V_S(X_0, t_0) &\leq G(\partial_\nu^+ U_S, \partial_\nu^- U_S)(X_0, t_0) \text{ (resp. } \geq), \text{ (two-phase case),} \\ V_S(X_0, t_0) &\leq G(\partial_\nu^+ U_S)(X_0, t_0) \text{ (resp. } \geq), \text{ (one-phase case).} \end{aligned}$$

It should be useful to discuss the notion of viscosity solutions between what we have proposed above and the definition given by Kim in [46] for the one-phase Hele-Shaw problem. Here we recall the definition in [46].

Definition 9.4. A non-negative upper semicontinuous function $U : \Sigma_0 \times [0, \infty) \rightarrow \mathbb{R}$ is said to be a viscosity subsolution to the one-phase Hele-Shaw problem, if

- (i) $U = U_0$ at $t = 0$, $U \leq 1$ on Γ_0 ;
- (ii) $\overline{\{U > 0\}} \cap \{t = 0\} = \overline{\{U(X, 0) > 0\}}$;
- (iii) for each $T \geq 0$ the set $\overline{\{U > 0\}} \cap \{t \leq T\}$ is bounded and
- (iv) for every $\Phi \in C_{x,t}^{2,1}$ that has a local maximum of

$$U - \Phi \text{ in } \overline{\{U > 0\}} \cap \{t \leq t_0\} \cap Q \text{ at } (X_0, t_0),$$

- (a) $-\Delta\Phi(x_0, t_0) \leq 0$ if $U(X_0, t_0) > 0$,
- (b) $\min(-\Delta\Phi, \partial_t\Phi - |\nabla\Phi|^2)(X_0, t_0) \leq 0$, if $(X_0, t_0) \in \partial\{U > 0\}$, $U(X_0, t_0) = 0$.

A non-negative lower semicontinuous function $U : \Sigma_0 \times [0, \infty) \rightarrow \mathbb{R}$ is said to be a viscosity supersolution to the one-phase Hele-Shaw problem, if

- (i) $U = U_0$ at $t = 0$, $U \geq 1$ on Γ_0 ;
- (ii) for every $\Phi \in C_{X,t}^{2,1}$ that has a local minimum of

$$U - \Phi \text{ in } \overline{\{U > 0\}} \cap \{t \leq t_0\} \cap Q \text{ at } (x_0, t_0),$$

- (a) $-\Delta\Phi(x_0, t_0) \geq 0$ if $(X_0, t_0) \in \{U > 0\}$,
- (b) $\max(-\Delta\Phi, \partial_t\Phi - |\nabla\Phi|^2)(X_0, t_0) \geq 0$, if $(X_0, t_0) \in \partial\{U > 0\}$, $|\nabla\Phi(X_0, t_0)| \neq 0$ and $\{\Phi > 0\} \cap \{U > 0\} \cap B \neq \emptyset$

for any ball B centered at (X_0, t_0) .

Then, U is a viscosity solution if it is both a viscosity supersolution and its upper semicontinuous envelope U_* is a viscosity subsolution.

Lemma 9.5. For the one-phase Hele-Shaw problem, if U is a viscosity subsolution (supersolution) in the sense of Definition 9.4 then it is a viscosity subsolution (supersolution) in the sense of Definition 9.3.

Proof. Let U be a solution in the sense of Definition 9.4. Fix $(X_0, t_0) \in \partial\{U > 0\}$ and let S be a test interface in $[t_0 - \tau, t_0]$ touching U from above at (X_0, t_0) , for some small $\tau > 0$. Let U_S be as given by (9.2) with $F(D^2U, \nabla U) = \Delta U$.

Fix $\varepsilon > 0$ and let $\tilde{\Phi}_\varepsilon : \Sigma_0 \times [t_0 - \tau, t_0]$ be the function given by

$$\begin{cases} \Delta\tilde{\Phi}_\varepsilon + \varepsilon = 0 \text{ in } S^+, \\ \tilde{\Phi}_\varepsilon = 0 \text{ on } S, \\ \tilde{\Phi}_\varepsilon = 1 \text{ on } \Gamma_0. \end{cases}$$

In particular, $\tilde{\Phi}_\varepsilon$ is superharmonic in S^+ and that the following pointwise limits hold

$$\lim_{\varepsilon \rightarrow 0} \tilde{\Phi}_\varepsilon = U_S \text{ in } S^+, \quad \lim_{\varepsilon \rightarrow 0} \partial_n^+ \tilde{\Phi}_\varepsilon = \partial_n^+ U_S \text{ on } S.$$

On the other hand, it is clear that $\tilde{\Phi}_\varepsilon$ is $C^{2,1}$ in S^+ . Let Φ_ε be a smooth extension of $\tilde{\Phi}_\varepsilon$ to $\Sigma_0 \times [t_0 - \tau, t_0]$.

By construction, for any ε we have that $U - \Phi_\varepsilon$ has a local maximum at some point $(X_0, t_0) \in S$ (seen as a function in $\{U > 0\} \cap \{t \leq t_0\}$). Applying Definition 9.4, it follows that at (X_0, t_0) at least one of the following inequalities must hold

$$\Delta \Phi_\varepsilon \geq 0 \quad \text{or} \quad \partial_t \Phi_\varepsilon \leq |\nabla \Phi_\varepsilon|^2.$$

Since $\Delta \Phi = -\varepsilon < 0$ along S , we conclude that

$$\partial_t \Phi_\varepsilon \leq |\nabla \Phi_\varepsilon|^2 \text{ at } (X_0, t_0), \quad \text{and hence } V_S(X_0, t_0) \leq \partial_n^+ \Phi_\varepsilon(X_0, t_0).$$

Letting $\varepsilon \rightarrow 0$, we conclude that

$$V_S(X_0, t_0) \leq \partial_n^+ U_S(X_0, t_0).$$

and we conclude that U is a solution in the sense of Definition 9.3. \square

Remark 9.6. *The use of a slightly superharmonic function $\tilde{\Phi}_\varepsilon$ instead of the harmonic function U_S is a type of argument familiar from the theory of viscosity solutions when dealing with boundary conditions, cf. [38, Proposition A.2] where the Neumann condition for a non-linear problem is studied. A similar situation is seen in free boundary problems, see for instance the discussion following [34, Definition 2.5].*

It will be necessary to understand how $\partial_n^\pm U_S$ varies with vertical shifts of S , and accordingly we introduce some notation for such shifts. Given a test interface S and $h \in \mathbb{R}$, we define

$$S_h := \{X = (x, x_{d+1}) \mid (x, x_{d+1} - h) \in S\}. \quad (9.3)$$

In other words S_h is the surface resulting from shifting S in the upward direction by h (if $h < 0$, then S_h is S shifted down by $|h|$). The following Lemma is, essentially, an extension of Lemma 5.15 with a slightly different set up that will be convenient in the next subsection.

Lemma 9.7. *Let S be a test interface. There is a constant $C = C_S$ such that for all sufficiently small h , and any $(X, t) \in S_h$ (S_h as in (9.3)), we have*

$$|\partial_n^+ U_{S_h}(X - he_{d+1}, t) - \partial_n^+ U_S(X, t)| \leq C|h|.$$

Here, U_S and U_{S_h} are the functions given by (9.1).

Remark 9.8. *Using an argument analogous to that in Lemma 6.1, one can use the one-phase problem estimate in Lemma 9.7 to obtain the analogous bound for the two-phase problem*

$$|G(\partial_n^+ U_{S_h}, \partial_n^- U_{S_h}) - G(\partial_n^+ U_S, \partial_n^- U_S)| \leq C_{G,S}|h|.$$

Proof. We shall consider only h such that $|h|$ is no larger than half the distance from S to $\{x_{d+1} = 0\}$, and we focus solely on the case $h > 0$ (the proof is similar for $h < 0$). As in Lemma 5.15, we shall compare U_S to a vertical shift of U_{S_h} . Thus, we consider \tilde{U} defined by

$$\tilde{U}(X, t) := U_{S_h}(X - he_{d+1}, t), \quad \text{defined in } S^+ \cap \{(X, t) \mid X = (x, x_{d+1}), x_{d+1} \geq h\}.$$

Observe that $\tilde{U} = U_S = 0$ on S . On the other hand, taking into account that $\tilde{U} \equiv 1$ on $\{x_{d+1} = h\}$, that $U_S \equiv 1$ on $\{x_{d+1} = 0\}$, and the available Lipschitz bound for U_S in $\{0 \leq x_{d+1} \leq h\}$, we have

$$|\tilde{U} - U_S| \leq C_S h \text{ on } \{x_{d+1} = h\}.$$

Then, applying the maximum principle in $S^+ \setminus \{0 \leq x_{d+1} \leq h\}$, it follows that

$$|\tilde{U} - U_S| \leq C_S h \text{ in } S^+ \setminus \{0 \leq x_{d+1} \leq h\}.$$

Then, using that $\tilde{U} - U_S = 0$ on S and applying Proposition 5.8, it follows that for some constant C depending on S (as well as the dimension and the ellipticity of F),

$$|\partial_n^+(\tilde{U} - U_S)| \leq Ch, \quad \text{on } S.$$

Since $\partial_n^+ \tilde{U}(X, t) = \partial_n^+ U_{S_h}(X - he_{d+1}, t)$, the Lemma is proved. \square

9.2. Correspondence between viscosity solutions of the free boundary evolution and viscosity solutions of the parabolic equation. In this subsection, we are going to show that under the graph assumption for $\partial\{U > 0\}$, viscosity solutions for the free boundary problem correspond to viscosity solutions for the parabolic equations.

First, we show a viscosity solution to the parabolic equation yields a viscosity solution to the free boundary problem (in the sense of Definition 9.3). We recall for the sake of convenience that in this level set context, when U_f solves (1.1), the equation for the normal velocity for U_f becomes

$$\partial_t U_f = G(\partial_n^+ U_f, \partial_n^- U_f) |\nabla U_f|;$$

and for the function f , it is equivalent to

$$\partial_t f = G(I^+(f), I^-(f)) \cdot \sqrt{1 + |\nabla f|^2}. \quad (9.4)$$

(We recall that I^\pm are defined in Section 6)

The first relationship between the two different notions of solution will be made under an additional assumption that f is Lipschitz. Then the next lemma will remove this assumption.

Lemma 9.9. *If f is a globally Lipschitz viscosity subsolution (respectively supersolution) of the equation (9.4) in $\mathbb{R}^d \times [a, b]$, then U_f is a viscosity subsolution (resp. supersolution) in $\Sigma_0 \times [a, b]$ of the evolution (1.1).*

Proof. Let $(X_0, t_0) \in \Gamma_f$ and let S be a test interface in $[t_0 - \delta, t_0]$ touching U_f from above at (X_0, t_0) . We first need to reduce to the case in which there is an intermediate test interface that is a graph of some function over \mathbb{R}^d . We claim there is some ϕ such that $U_f \leq U_\phi \leq U_S$. For $\varepsilon > 0$ define ϕ^ε

$$\phi^\varepsilon(x, t) := \sup \left\{ P_{h, x', t'}(x, t) \mid (h, x', t') \text{ s.t. } D_{P_{h, x', t'}} \subset \{U_S > 0\} \right\}.$$

where

$$P_{h, x', t'}(x, t) := h - \frac{1}{2\varepsilon}|x - x'|^2 - \frac{1}{2\varepsilon}|t - t'|^2.$$

(For convenience, we recall that the notation for the set $D_{P_{h, x', t'}}$ appears in (1.4).) It is clear that $D_{\phi^\varepsilon} \subset \{U_S > 0\}$. Therefore, applying the comparison principle yields

$$U_{\phi^\varepsilon} \leq U_S \text{ everywhere, and } U_{\phi^\varepsilon} = U_S \text{ on } \Gamma_{\phi^\varepsilon} \cap S.$$

The interface $\partial\{U_S > 0\}$ is smooth and touches Γ_f at (X_0, t_0) , since f is Lipschitz, the normal to $\partial\{U_S > 0\}$ cannot be orthogonal to e_{d+1} . This means that if ε is chosen sufficiently small (depending on the smoothness of S) then there is some $x_0 \in \mathbb{R}^d$ such that

$$(X_0, t_0) \in \Gamma_{\phi^\varepsilon} \cap \partial\{U_S > 0\}, \text{ and } X_0 = (x_0, \phi^\varepsilon(x_0, t_0)).$$

In particular, for such ε we have $G(\partial_n^+ U_S, \partial_n^- U_S) \geq G(\partial_n^+ U_{\phi^\varepsilon}, \partial_n^- U_{\phi^\varepsilon})$ and hence $V^\varepsilon \geq V$ at this contact point (where V^ε is the normal velocity of $\partial\{U_{\phi^\varepsilon} > 0\}$). Moreover, from the construction of ϕ^ε and the fact that $U_f \leq U_S$, it follows that

$$f(x, t) \leq \phi^\varepsilon(x, t) \text{ and } f(x_0, t_0) = \phi^\varepsilon(x_0, t_0).$$

(One can see this from the fact that choosing $h = f(x, t)$ gives that $P_{h, x', t'} \geq f^\varepsilon$, where f^ε is the sup-convolution of f , so $f \leq f^\varepsilon \leq \phi^\varepsilon$.) Since ϕ^ε is pointwise $C^{1,1}$ at (x_0, t_0) and f is a viscosity subsolution of (9.4), it follows that (cf. Lemma 8.5)

$$\partial_t \phi^\varepsilon \leq G(I^+(\phi^\varepsilon), I^-(\phi^\varepsilon)) \cdot \sqrt{1 + |\nabla \phi^\varepsilon|^2}, \text{ at } (x_0, t_0).$$

Recalling the level set formulation as it pertains to the defining function that is $\Phi(X, t) = f(x, t) - x_{d+1}$, we see that as the set $\partial\{U_{\phi^\varepsilon} > 0\} = \Gamma_{\phi^\varepsilon}$, the normal velocity V^ε is

$$\frac{\partial_t \phi^\varepsilon}{\sqrt{1 + |\nabla \phi^\varepsilon|^2}} = V^\varepsilon.$$

In other words, in light of the definitions of I^\pm , we see that

$$V^\varepsilon \leq G(\partial_n^+ U_{\phi^\varepsilon}, \partial_n^- U_{\phi^\varepsilon}) \text{ at } (X_0, t_0).$$

After combining all of the above arguments, we have established that

$$V \leq V^\varepsilon \leq G(\partial_n^+ U_{\phi^\varepsilon}, \partial_n^- U_{\phi^\varepsilon}) \leq G(\partial_n^+ U_S, \partial_n^- U_S) \text{ at } (X_0, t_0).$$

This shows that U_f is a viscosity subsolution of the free boundary flow.

If f is a supersolution, the argument is similar, so we only highlight some of the steps: suppose now the test interface S touches U_f from below at a free boundary point (X_0, t_0) , then for $\varepsilon > 0$ we define ϕ_ε by

$$\phi_\varepsilon(x, t) := \inf \left\{ Q_{h, x', t'}(x, t) \mid (h, x', t') \text{ s.t. } \{U_S > 0\} \subset D_{Q_{h, x', t'}} \right\},$$

where

$$Q_{h, x', t'}(x, t) := h + \frac{1}{2\varepsilon}|x - x'|^2 + \frac{1}{2\varepsilon}|t - t'|^2.$$

As before, from the construction of ϕ_ε one can see that $\partial\{U_S > 0\} \subset D_{\phi_\varepsilon}$. The comparison principle then says that

$$U_S \leq U_{\phi_\varepsilon} \text{ everywhere, and } U_{\phi_\varepsilon} = U_S \text{ on } \Gamma_{\phi_\varepsilon} \cap S.$$

The fact that f is Lipschitz and the smoothness $\partial\{U_S > 0\}$ means that the normal to $\partial\{U_S > 0\}$ at the contact point cannot be perpendicular to e_{d+1} . It follows that for all sufficiently small ε there is a $x_0 \in \mathbb{R}^d$ such that $X_0 = (x_0, \phi_\varepsilon(x_0, t_0))$ and

$$\phi_\varepsilon(x, t) \leq f(x, t) \text{ and } \phi_\varepsilon(x_0, t_0) = f(x_0, t_0).$$

Arguing as in the case of a subsolution, one arrives at

$$V \geq G(\partial_n^+ U_S, \partial_n^- U_S) \text{ at } (X_0, t_0).$$

□

Now we can prove the same result, but without a Lipschitz assumption on f .

Lemma 9.10. *If f is a viscosity subsolution (respectively supersolution) of (9.4) in $\mathbb{R}^d \times [a, b]$, then U_f is a viscosity subsolution (resp. supersolution) of (1.1) in $\Sigma_0 \times [a, b]$.*

Proof. Let f be a viscosity subsolution of the problem in $\mathbb{R}^d \times [a, b]$. Let $\varepsilon > 0$ be appropriately small and define the sup-convolution f^ε ; we see f^ε is still a viscosity subsolution and it is also a Lipschitz continuous function. By Lemma 9.9, it follows that U_{f^ε} is a viscosity subsolution of the free boundary problem for every $\varepsilon > 0$.

Let S be a test interface touching U_f from above at (X_0, t_0) , $X_0 = (x_0, f(x_0, t_0))$. For each sufficiently small $\varepsilon > 0$ there is $h_\varepsilon \in \mathbb{R}$ such that (see (9.3)) S_{h_ε} touches f_ε from above at a point of the form $(X_\varepsilon, t_\varepsilon) \in S_{h_\varepsilon}$, $X_\varepsilon = (x_\varepsilon, f_\varepsilon(x_\varepsilon, t_\varepsilon))$. Moreover, the $(X_\varepsilon, t_\varepsilon)$ and h_ε can be chosen so that

$$\lim_{\varepsilon \rightarrow 0} (X_\varepsilon, t_\varepsilon) = (X_0, t_0), \quad \lim_{\varepsilon \rightarrow 0} h_\varepsilon = 0. \quad (9.5)$$

Then, as noted above U_{f_ε} is a viscosity subsolution, and it follows that

$$\partial_t V_{S_{h_\varepsilon}}(X_\varepsilon, t_\varepsilon) \leq G(\partial_n^+ U_{S_{h_\varepsilon}}(X_\varepsilon, t_\varepsilon)).$$

On the other hand, by Lemma 9.7 (and Remark 9.8 for the two-phase case),

$$|\partial_n^+ U_S(X_\varepsilon, t_\varepsilon) - \partial_n^+ U_{S_{h_\varepsilon}}(\tilde{X}_\varepsilon, t_\varepsilon)| \leq Ch_\varepsilon, \quad \tilde{X}_\varepsilon := X_\varepsilon - h_\varepsilon e_{d+1},$$

and it follows that

$$\partial_t V_S(\tilde{X}_\varepsilon, t_\varepsilon) \leq G(\partial_n^+ U_S(\tilde{X}_\varepsilon, t_\varepsilon)) + C|h_\varepsilon|.$$

Now, given that $(\tilde{X}_\varepsilon, t_\varepsilon) \rightarrow (X_0, t_0)$ as $\varepsilon \rightarrow 0$, we conclude that

$$\partial_t V_S(X_0, t_0) \leq G(\partial_n^+ U_S(X_0, t_0)).$$

Here we note that the reason this works is that all of the calculations occur at the level of U_S and $U_{S_{h_\varepsilon}}$, which are both smooth; hence the limiting operations are straightforward. This proves U_f is a subsolution. The argument for a supersolution is entirely analogous and we omit the details. \square

The next proposition is a basic observation about how test functions for the lower dimensional, non-local parabolic problem yield to test functions of the free boundary problem.

Proposition 9.11. *Let ϕ be an admissible test function touching f from above at some $t_0 \in [a, b]$ and $x_0 \in \Sigma_0$. Then, U_ϕ touches U_f from above at (X_0, t_0) where $X_0 = (x_0, f(x_0, t_0))$.*

Proof. First of all, since $f \leq \phi$ everywhere, it follows that

$$D_f \subset D_\phi.$$

Then, from the definition of U_ϕ, U_f it follows that $U_\phi \geq U_f$ on ∂D_f and $U_\phi \geq U_f$ on ∂D_f^- . Applying the comparison principle to each phase yields

$$U_f \leq U_\phi \text{ in } D_f \text{ and } U_f \leq U_\phi \text{ in } D_f^-.$$

This means that $U_f \leq U_\phi$ for all times $t \in [a, t_0]$. Since $f = \phi$ at (x_0, t_0) , we have $U_f = U_\phi = 0$ at time t_0 at the point $(x_0, f(x_0)) = (x_0, \phi(x_0))$, proving the proposition. \square

Lemma 9.12. *Let U be a viscosity subsolution (supersolution) of (1.1) in $\Sigma_0 \times [a, b]$ whose free boundary is given as the graph of some upper semi-continuous f (lower semi-continuous), then f is a viscosity subsolution (supersolution) of (9.4) in $\mathbb{R}^d \times [a, b]$.*

Proof. Let ϕ be a test function which touches f from above at (x_0, t_0) . According to Proposition 9.11, U_ϕ is a test function which touches U from above at (X_0, t_0) , where $X_0 = (x_0, f(x_0, t_0))$. Since U is a viscosity subsolution, it follows that

$$V_{U_\phi} \leq G(\partial_n^+ U_\phi, \partial_n^- U_\phi), \text{ at } (X_0, t_0)$$

This implies that at (X_0, t_0) we have

$$\frac{\partial_t \phi}{\sqrt{1 + |\nabla \phi|^2}} \leq G(I^+(\phi), I^-(\phi)).$$

Since ϕ was arbitrary, it follows f is a viscosity subsolution. The proof for supersolutions is entirely analogous and we omit it. \square

10. PROPAGATION OF THE MODULUS OF CONTINUITY

After all of the build-up of the preceding sections, we can now demonstrate that a modulus of continuity of initial data will be preserved by the fractional parabolic equation. This result is very simple once the comparison theorem for viscosity solutions has been established.

Lemma 10.1. *Let J be as in assumption 7.1, $T > 0$ and $f : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$ a continuous viscosity solution of*

$$\begin{aligned} \partial_t f &= J(f) \text{ in } \mathbb{R}^d \times [0, T], \\ f(x, 0) &= f_0(x) \text{ in } \mathbb{R}^d. \end{aligned}$$

If f_0 is continuous with modulus of continuity $\omega(\cdot)$, then the same will be true of $f(\cdot, t)$ for all $t \in [0, T]$. In particular, we have the estimate

$$|f(x, t) - f(y, t)| \leq \omega(|x - y|).$$

Proof. Let $h \in \mathbb{R}^d$ be fixed, and consider the function,

$$w(x, t) := (\tau_h f)(x, t) - f(x, t).$$

By assumption,

$$\forall x \in \mathbb{R}^d, \quad w(x, 0) \leq \omega(|h|).$$

By translation invariance of J and Lemma ??, w satisfies, in the viscosity sense

$$\partial_t w \leq M_J^+(w).$$

In light of the above, Lemma 8.8 implies that for any $(x, t) \in \mathbb{R}^d \times [0, T]$,

$$w(x, t) \leq \omega(|h|).$$

Since this holds for an arbitrary vector h , the lemma is proved. \square

11. THE PROOFS OF THEOREMS 1.1 AND 1.4

In this section, we collect all of the various facts of the previous sections that combine to prove Theorems 1.1 and 1.4.

11.1. Proof / explanation of Theorem 1.1. To be precise about the equation for the one-phase problem, we mean that for all t , $\partial\{U_f > 0\} = \text{graph}(f(\cdot, t))$, that U_f solves the problem (1.1), in the free boundary viscosity sense of Section 9, and that the normal velocity is given by (5.2)

$$V(x, f(x)) = G(\partial_n U_f(x, f(x))) = G(I(f, x)).$$

Furthermore, the only restriction on f is that $f \geq \delta$ and that f is bounded and continuous on $\mathbb{R}^d \times [0, T]$.

To be precise about the equation for the two-phase problem, we take the analogous set-up, but the normal velocity is prescribed as (6.5),

$$V(x, f(x)) = G(\partial_n^+ U_f(x, f(x)), \partial_n^- U_f(x, f(x))) = H(f, x).$$

The notion of solution is again the free boundary viscosity solution of Section 9. We have a further restriction that $\delta \leq f \leq L - \delta$ and f is continuous on $\mathbb{R}^d \times [0, T]$.

We note that per the assumptions of Section 5, for the one-phase problem, and Section 6 for the two-phase problem, we always assume the free boundary is a *bounded* graph for all times. Section 9 (Lemmas 9.10 and 9.12) contains the proof of (i), that under the graph assumption, the free boundary viscosity solution is equivalent to the parabolic viscosity solution of

$$\begin{cases} \partial_t f = J(f) & \text{in } \mathbb{R}^d \times (0, T) \\ f(\cdot, 0) = f_0 & \text{on } \mathbb{R}^d. \end{cases}$$

Thanks to Theorem 7.10, both

$$J(f) = G(I(f)) \cdot \sqrt{1 + |\nabla f|^2} \quad \text{and} \quad H(f) = G(f) \cdot \sqrt{1 + |\nabla f|^2}$$

are admissible operators for solving the parabolic equation, for respectively the one and two phase problems (for I, H , defined respectively in (5.2), (6.5)).

The proof of the propagation of modulus claimed in Theorem 1.1 (ii) is a consequence of the correspondence of solutions in (i), combined with the result of Lemma 10.1, which is applicable thanks to Theorem 7.10. This was, of course, only possible because of the comparison theorem deduced in Section 8.

The proof of the existence of viscosity solutions claimed in Theorem 1.1 (iii), is a consequence of the correspondence of solutions in (i), with the proof of existence in Section 8.2.

The precise statement of Theorem 1.4 has the operator I defined with equation (5.1), through the formula (5.2). The proof of Lipschitz nature of I in Theorem 1.4, is a consequence of Theorem 5.13. The min-max formula is a result of Theorem 7.5, combined with verification the I satisfies the results of Assumption 7.1, which is in Theorem 7.10.

12. COMMENTS ABOUT OUR ASSUMPTIONS, AND SOME QUESTIONS

There are many assumptions that accompany our Theorems 1.1 and 1.4. Some are for simplicity and some are more nuanced. Here we discuss how some could be removed and how some may be difficult to overcome. At the end, we suggest some lines of open questions that could be useful.

12.1. Translation invariance. In all of the above situations, we have only treated free boundary evolutions in which the equations in the positive and negative phases of U , as well as the free boundary condition, G , are translation invariant. We would like to say that we did this out of reasons of simplicity, but we cannot claim that motivation, as it is not clear how difficult it will be to remove this assumption. The main place it was used in a tangible fashion was in the proof of Lemma 5.15, and this lemma played a fundamental role in most of the major results that followed— in particular Theorem 5.13 and all subsequent results invoking it. Although the translation invariance led to a simpler min-max representation in Section 7, it does not play an essential role there. It does play, however, a substantial role in the arguments of Section 8. This is because (surprisingly) it is still unknown to what extent the uniqueness theory for viscosity solutions of integro-differential equations is applicable. Presumably, one expects the uniqueness theory to parallel that of second order equations, but to date, there are only a handful of works that can handle equations with non-trivial dependence on x ; we defer to the work of Barles-Imbert [6] for the latest results and earlier references. In particular, if one pursues the x -dependent min-max formulation in Section 7, it is not clear if the resulting integro-differential operators will fall into the scope of [6].

12.2. Rotational invariance. In principle, this should only be a technical assumption that is an artifact of our method to establish the Lipschitz continuity of the operator, I , culminating in Theorem 5.13. The interested reader can probe more deeply into our method of proof for Theorem 5.13, where we rotated two domains so that they would intersect in a C^1 fashion. The way we chose to deal with this situation required to rotate the underlying equations, hence needing an invariance assumption in the absence of better estimates. It would be helpful to remove this assumption.

12.3. Graph assumption. As the careful reader will see, we *assumed* in Theorem 1.1 that if the free boundary evolution, $\partial\{U(\cdot, t) > 0\}$ is a graph for all times, then one can deduce a correspondence between the free boundary viscosity solution and the parabolic viscosity solution. This raises the question of whether or not the graph assumption is reasonable? Answering this question is essential to the future use of the methods suggested in this paper. This ties in with the star-shaped assumption that appears for some of the preliminary results in [27]. In [27], they were able to first focus on star-shaped initial free boundaries (which are equivalent to being a graph over some sphere), and then they subsequently extended the regularity results to more general boundaries. It is conceivable that something similar could be done in our context, but it is not obvious how to carry out this procedure at the moment.

12.4. Some Questions. We believe there are at least a few natural questions resulting from the results presented above, and we mention a few of them. This list is by no means complete, but we think these are some of the most interesting questions.

x -dependence. As mentioned above, all ingredients (F_1 , F_2 , G) were taken to be independent of x . What can be said when x dependence is allowed? This may lead to a better understanding of where one may hope to push the uniqueness and regularity theory for viscosity solutions of integro-differential operators like the ones that appear in (7.2).

Anisotropic G . What happens if G is allowed to depend, in an appropriately monotone way, on the full gradients, ∇U_f^+ and ∇U_f^- , instead of simply $\partial_n^+ U_f$ and $\partial_n^- U_f$? Any reasonable ellipticity restriction on G should keep it within the realm of the GCP and would likely enjoy properties similar to those listed in Assumption 7.1. Is this indeed the case?

Lévy measures. Is it possible to find more detailed information about the Lévy measures, μ^{ij} , that appear in Theorem 1.4? Can it be proved that they are in a class for which some regularity theory is known (e.g. [12], [23], [24], or [55])? Are μ^{ij} in a different class that has not yet been studied, but possibly still enjoy certain regularity properties for solutions of $\partial_t f = I(f)$? Answering these questions will be an essential building block in order that the nonlocal theory can shed new light on these free boundary problems.

Graphs over a manifold. In this paper, we made the assumption that the free boundary was a graph over \mathbb{R}^d that evolves with time. Can the results here be extended to include the case where the free boundary, *for short times*, is the evolution of a graph over some submanifold of \mathbb{R}^{d+1} (in this case, the natural submanifold would be that of the initial conditions, $\partial\{U(\cdot, 0) > 0\}$)? This situation is not incompatible with the min-max theory developed in Section 7. Indeed, the results of [37] were proved for operators acting on functions in, e.g. $C^{1,\gamma}(\partial\Omega)$, whenever $\partial\Omega$ is a nice enough hyper-surface. Furthermore, this idea of tracking the free boundary as a graph over a reference manifold is compatible with some of the results in [27], where the preservation of the star-shape property is equivalent to being a graph over a sphere in the radial direction. Again, even if one can justify the graph assumption, this would likely result in nonlocal parabolic equations that have non-trivial x -dependence (in fact, the notion of translation invariance makes no sense in this situation). We note that in the case of rotationally invariant operators in the respective positive and negative phases, this would result in a rotationally invariant integro-differential operator for functions on the sphere—this should be amenable to similar ideas we used for translation invariant equations for functions on \mathbb{R}^d . Studying the operator on a manifold that does not necessarily admit an invariant group of transformations puts the resulting parabolic equation for the free boundary into a class of equations for which existence, uniqueness, regularity, etc... is not well established (i.e. viscosity solutions for nonlocal parabolic equations with x -dependent coefficients, posed over a manifold).

Equations in divergence form. One should note that we did not treat the case when in the set $\{U > 0\}$, we have the equation

$$F(D^2U, \nabla U) = \operatorname{div}(A(x)\nabla U) = 0.$$

In most reasonable circumstances, this equation enjoys the maximum principle, and hence also the free boundary operator will enjoy the GCP. Note that this equation is not translation invariant (otherwise, it is contained in our results). However, the methods and results for the divergence theory are a bit different from those employed herein for the fully nonlinear theory, and so there may be different outcomes. Thus, the question is: how much can be said about the operator, I , under this assumption? It may be possible that within the scope of Section 7, the Lévy measures could be shown to have nice properties with respect to Lebesgue measure on \mathbb{R}^d , akin to some of the existing integro-differential results. The counter-balance to this benefit would be the fact that the integro-differential equation would no longer be translation invariant. However, the known results about Green's functions and Poisson kernels for divergence equations may shed extra light on the situation. It is conceivable that under some extra regularity assumptions on $A(x)$, one may obtain a class of integro-differential operators that enjoys known Schauder-type results, or could be within reach of minor modifications of such existing results.

Problems of Stefan type. If one replaces equation (1.1) with a parabolic problem, then the resulting free boundary problem is known as the Stefan problem. It could be possible that the ideas in Sections 5 and 7 might extend to the Stefan set-up. This would require a new result, similar to that of Section 7 that applied to operators with the *space-time* GCP instead of just the GCP in space alone. One might conjecture that the resulting linear operators in the min-max would be nonlocal in both space and time, not just nonlocal in space, as was proved here. That would mean that the

space-time nonlocal theory could be relevant, such as, e.g. Allen [3] (which treats a more general form of the Caputo-Marchaud time derivative), or even a nonlinear (in time) extension of similar results.

REFERENCES

- [1] Andrew Acker. On the existence of convex classical solutions to a generalized prandtl-batchelor free boundary problem. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 49(1):1–30, 1998.
- [2] Nicholas D Alikakos, Peter W Bates, and Xinfu Chen. Convergence of the cahn-hilliard equation to the hele-shaw model. *Archive for rational mechanics and analysis*, 128(2):165–205, 1994.
- [3] Mark Allen. Holder regularity for nondivergence nonlocal parabolic equations. *arXiv preprint arXiv:1610.10073*, 2016.
- [4] David M Ambrose. Well-posedness of two-phase hele–shaw flow without surface tension. *European Journal of Applied Mathematics*, 15(5):597–607, 2004.
- [5] I. Athanasopoulos, L. Caffarelli, and S. Salsa. Regularity of the free boundary in parabolic phase-transition problems. *Acta Math.*, 176(2):245–282, 1996.
- [6] Guy Barles and Cyril Imbert. Second-order elliptic integro-differential equations: viscosity solutions’ theory revisited. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 25(3):567–585, 2008.
- [7] G. K. Batchelor. On steady laminar flow with closed streamlines at large Reynolds number. *J. Fluid Mech.*, 1:177–190, 1956.
- [8] G. K. Batchelor. A proposal concerning laminar wakes behind bluff bodies at large Reynolds number. *J. Fluid Mech.*, 1:388–398, 1956.
- [9] Garrett Birkhoff and Eduardo Zarantonello. *Jets, wakes, and cavities*. 1957.
- [10] L. Caffarelli, M. G. Crandall, M. Kocan, and A. Świąch. On viscosity solutions of fully nonlinear equations with measurable ingredients. *Comm. Pure Appl. Math.*, 49(4):365–397, 1996.
- [11] Luis Caffarelli and Sandro Salsa. *A geometric approach to free boundary problems*, volume 68 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [12] Luis Caffarelli and Luis Silvestre. Regularity theory for fully nonlinear integro-differential equations. *Comm. Pure Appl. Math.*, 62(5):597–638, 2009.
- [13] Luis Caffarelli and Juan Luis Vazquez. Viscosity solutions for the porous medium equation. In *Proceedings of Symposia in Pure Mathematics*, volume 65, pages 13–26. American Mathematical Society, 1999.
- [14] Luis A. Caffarelli. A Harnack inequality approach to the regularity of free boundaries. III. Existence theory, compactness, and dependence on X . *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 15(4):583–602 (1989), 1988.
- [15] Luis A Caffarelli. The obstacle problem revisited. *Journal of Fourier Analysis and Applications*, 4(4-5):383–402, 1998.
- [16] Luis A. Caffarelli and Xavier Cabré. *Fully nonlinear elliptic equations*, volume 43 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1995.
- [17] Luis A. Caffarelli, David Jerison, and Carlos E. Kenig. Some new monotonicity theorems with applications to free boundary problems. *Ann. of Math. (2)*, 155(2):369–404, 2002.
- [18] Luis A Caffarelli and Juan L Vázquez. A free-boundary problem for the heat equation arising in flame propagation. *Transactions of the American Mathematical Society*, 347(2):411–441, 1995.
- [19] G Caginalp. Stefan and hele-shaw type models as asymptotic limits of the phase-field equations. *Physical Review A*, 39(11):5887, 1989.
- [20] Gunduz Caginalp and Xinfu Chen. Convergence of the phase field model to its sharp interface limits. *European Journal of Applied Mathematics*, 9(4):417–445, 1998.
- [21] Stephen Cameron. Global well-posedness for the 2d muskat problem with slope less than 1. *arXiv preprint arXiv:1704.08401*, 2017.
- [22] Angel Castro, Diego Córdoba, Charles L. Fefferman, Francisco Gancedo, and María López-Fernández. Turning waves and breakdown for incompressible flows. *Proc. Natl. Acad. Sci. USA*, 108(12):4754–4759, 2011.
- [23] Héctor Chang Lara. Regularity for fully non linear equations with non local drift. *arXiv:1210.4242 [math.AP]*, 2012.
- [24] Héctor Chang Lara and Gonzalo Dávila. Regularity for solutions of nonlocal, nonsymmetric equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 29(6):833–859, 2012.
- [25] Héctor A Chang-Lara and Nestor Guillen. From the free boundary condition for hele-shaw to a fractional parabolic equation. *arXiv preprint arXiv:1605.07591*, 2016.
- [26] CH Cheng, Daniel Coutand, and Steve Shkoller. Global existence and decay for solutions of the hele-shaw flow with injection. *arXiv preprint arXiv:1208.6213*, 2012.
- [27] Sunhi Choi, David Jerison, and Inwon Kim. Regularity for the one-phase Hele-Shaw problem from a Lipschitz initial surface. *Amer. J. Math.*, 129(2):527–582, 2007.
- [28] Sunhi Choi, David Jerison, and Inwon Kim. Local regularization of the one-phase Hele-Shaw flow. *Indiana Univ. Math. J.*, 58(6):2765–2804, 2009.
- [29] Frank H Clarke. *Optimization and nonsmooth analysis*, volume 5. Siam, 1990.

- [30] Peter Constantin, Diego Córdoba, Francisco Gancedo, and Robert M. Strain. On the global existence for the Muskat problem. *J. Eur. Math. Soc. (JEMS)*, 15(1):201–227, 2013.
- [31] Diego Córdoba and Francisco Gancedo. Contour dynamics of incompressible 3-D fluids in a porous medium with different densities. *Comm. Math. Phys.*, 273(2):445–471, 2007.
- [32] Philippe Courrege. Sur la forme intégró-différentielle des opérateurs de c_k^∞ dans c satisfaisant au principe du maximum. *Séminaire BreLOT-Choquet-Deny. Théorie du Potentiel*, 10(1):1–38, 1965.
- [33] Michael G. Crandall, Hitoshi Ishii, and Pierre-Louis Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [34] D. De Silva. Free boundary regularity for a problem with right hand side. *Interfaces Free Bound.*, 13(2):223–238, 2011.
- [35] G Duvaut and JL Lions. Les inégalités en mécanique et en physique. *Dunod, Paris*, 1972.
- [36] Joachim Escher, Gieri Simonett, et al. Classical solutions for Hele-Shaw models with surface tension. *Advances in Differential Equations*, 2(4):619–642, 1997.
- [37] Nestor Guillen and Russell W Schwab. Min-max formulas for nonlocal elliptic operators. *arXiv preprint arXiv:1606.08417*, 2016.
- [38] Nestor Guillen and Russell W. Schwab. Neumann homogenization via integro-differential operators. *Discrete and Continuous Dynamical Systems*, 36(7):3677–3703, 2016.
- [39] Mahir Hadžić and Steve Shkoller. Global stability and decay for the classical stefan problem. *Communications on Pure and Applied Mathematics*, 68(5):689–757, 2015.
- [40] Cyril Imbert and Luis Silvestre. An introduction to fully nonlinear parabolic equations. In *An introduction to the Kähler-Ricci flow*, volume 2086 of *Lecture Notes in Math.*, pages 7–88. Springer, Cham, 2013.
- [41] Hitoshi Ishii. Perron’s method for Hamilton-Jacobi equations. *Duke Math. J.*, 55(2):369–384, 1987.
- [42] Hitoshi Ishii. On uniqueness and existence of viscosity solutions of fully nonlinear second-order elliptic PDEs. *Comm. Pure Appl. Math.*, 42(1):15–45, 1989.
- [43] R. Jensen, P.-L. Lions, and P. E. Souganidis. A uniqueness result for viscosity solutions of second order fully nonlinear partial differential equations. *Proc. Amer. Math. Soc.*, 102(4):975–978, 1988.
- [44] Robert Jensen. The maximum principle for viscosity solutions of fully nonlinear second order partial differential equations. *Arch. Rational Mech. Anal.*, 101(1):1–27, 1988.
- [45] David Jerison and Inwon Kim. The one-phase Hele-Shaw problem with singularities. *J. Geom. Anal.*, 15(4):641–667, 2005.
- [46] Inwon C. Kim. Uniqueness and existence results on the Hele-Shaw and the Stefan problems. *Arch. Ration. Mech. Anal.*, 168(4):299–328, 2003.
- [47] Inwon C. Kim and Norbert Pořár. Viscosity solutions for the two-phase Stefan problem. *Comm. Partial Differential Equations*, 36(1):42–66, 2011.
- [48] David Kinderlehrer and Guido Stampacchia. *An introduction to variational inequalities and their applications*, volume 31. Siam, 1980.
- [49] A. Kiselev, F. Nazarov, and A. Volberg. Global well-posedness for the critical 2D dissipative quasi-geostrophic equation. *Invent. Math.*, 167(3):445–453, 2007.
- [50] J.-M. Lasry and P.-L. Lions. A remark on regularization in Hilbert spaces. *Israel J. Math.*, 55(3):257–266, 1986.
- [51] Stephan Luckhaus. Solutions for the two-phase stefan problem with the gibbs–thomson law for the melting temperature. *European Journal of Applied Mathematics*, 1(2):101–111, 1990.
- [52] Arshak Petrosyan, Henrik Shahgholian, and Nina N Ural’ceva. *Regularity of free boundaries in obstacle-type problems*, volume 136. American Mathematical Soc., 2012.
- [53] Jan Prüss, Jürgen Saal, and Gieri Simonett. Existence of analytic solutions for the classical stefan problem. *Mathematische Annalen*, 338(3):703–755, 2007.
- [54] Jan Prüss and Gieri Simonett. *Moving interfaces and quasilinear parabolic evolution equations*, volume 105. Springer, 2016.
- [55] Russell W. Schwab and Luis Silvestre. Regularity for parabolic integro-differential equations with very irregular kernels. *Anal. PDE*, 9(3):727–772, 2016.
- [56] Michael Siegel, Russel E. Caffisch, and Sam Howison. Global existence, singular solutions, and ill-posedness for the Muskat problem. *Comm. Pure Appl. Math.*, 57(10):1374–1411, 2004.
- [57] Luis Silvestre. On the differentiability of the solution to the Hamilton-Jacobi equation with critical fractional diffusion. *Adv. Math.*, 226(2):2020–2039, 2011.
- [58] Neil S. Trudinger. Hölder gradient estimates for fully nonlinear elliptic equations. *Proc. Roy. Soc. Edinburgh Sect. A*, 108(1-2):57–65, 1988.
- [59] Juan Luis Vázquez. *The porous medium equation: mathematical theory*. Oxford University Press, 2007.
- [60] Lihe Wang. On the regularity theory of fully nonlinear parabolic equations. I. *Comm. Pure Appl. Math.*, 45(1):27–76, 1992.

CENTRO DE INVESTIGACIÓN EN MATEMÁTICAS, A.C., JALISCO S/N, COL. VALENCIANA CP: 36023 GUANAJUATO, GTO, MÉXICO, APARTADO POSTAL 402, CP 36000.

Email address: `hector.chang@cimat.mx`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, AMHERST, MA 90095

Email address: `nguillen@math.umass.edu`

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, 619 RED CEDAR ROAD, EAST LANSING, MI 48824

Email address: `rschwab@math.msu.edu`