

## Notes on the complex exponential and sine functions (§1.5)

**I. Periodicity of the imaginary exponential.** Recall the definition: if  $\theta \in \mathbb{R}$  then  $e^{i\theta} \stackrel{\text{def}}{=} \cos \theta + i \sin \theta$ . Clearly  $e^{i(\theta+2\pi)} = e^{i\theta}$  (because of the  $2\pi$  periodicity of the sine and cosine functions of ordinary calculus). It's also clear—from drawing a picture of  $e^{i\theta}$  on the unit circle, say—that  $2\pi$  is the “minimal period” of the imaginary exponential, in the sense that if  $a \in \mathbb{R}$  has the property that  $e^{i(\theta+a)} = e^{i\theta}$  for every  $\theta \in \mathbb{R}$  then  $a = 2\pi n$  for some integer  $n$ .

**II. Periodicity of complex the exponential.** Recall the definition: if  $z = x + iy$  where  $x, y \in \mathbb{R}$ , then

$$e^z \stackrel{\text{def}}{=} e^x e^{iy} = e^x (\cos y + i \sin y).$$

It's clear from this definition and the periodicity of the imaginary exponential (§I) that  $e^{z+2\pi i} = e^z$ , i.e.: “The complex exponential function is periodic with period  $2\pi i$ .”

The first thing we want to show in these notes is that the period  $2\pi i$  is “minimal” in the same sense that  $2\pi$  is the minimal period for the imaginary exponential (and for the ordinary sine and cosine).

THE “MINIMAL PERIOD THEOREM” FOR THE COMPLEX EXPONENTIAL. *If  $\alpha \in \mathbb{C}$  has the property!*

$$(1) \quad e^{z+\alpha} = e^z \quad \text{for all } z \in \mathbb{C},$$

*then  $\alpha = 2\pi ni$  for some integer  $n$ !*

PROOF. If we set  $z = 0$  in (1) we see that  $e^\alpha = e^0 = 1$ . Now write  $\alpha$  in cartesian form:  $\alpha = a + ib$  where  $a, b \in \mathbb{R}$ . Then  $1 = e^\alpha = e^a e^{ib}$ . Take absolute values on both sides of this last equation to obtain  $1 = e^a$ , so, (because in this last equation we are dealing with the ordinary exponential of calculus)  $a = 0$ . Thus  $\alpha = ib$ , hence our previous equation  $e^\alpha = 1$  becomes:  $e^{ib} = 1$ . It follows from the work of §I that  $b = 2\pi n$  for some integer  $n$ . Thus  $\alpha = 2\pi in$ , as promised.  $\square$

COROLLARY.  $e^\alpha = 1 \iff \alpha = 2\pi ni$  for some integer  $n$ !

PROOF. If  $e^\alpha = 1$  then for each  $z \in \mathbb{C}$ :

$$e^{z+\alpha} = e^z e^\alpha = e^z$$

so  $\alpha$  is a period of the complex exponential, and hence, by the Theorem, is an integer multiple of  $2\pi i$ .  $\square$

**III. Univalence<sup>1</sup> of the complex exponential.** *The complex exponential is univalent in any open horizontal strip of width  $2\pi$  !or less!!*

*Remark!* Width  $2\pi$  is the best we can hope for by the periodicity of the complex exponential noted in §II above.

PROOF OF THEOREM. We'll do a little better, and show that: *if*

$$(2) \quad e^z = e^w \text{ with } |\operatorname{Im} z - \operatorname{Im} w| < 2\pi,$$

*then  $z = w$ !*

Thus, for example, if  $a \in \mathbb{R}$  and  $S$  is the (non-open) strip  $\{z \in \mathbb{C} : a < \operatorname{Im} z \leq a + 2\pi\}$ , then the complex exponential is univalent on  $S$ . Also, if  $S$  is any open ribbon-shaped region of vertical width  $2\pi$  or less (draw a picture!), then the complex exponential is univalent on  $S$ .

So suppose  $z$  and  $w$  are complex numbers that satisfy condition (2). We wish to show  $z = w$ . Multiply both sides of  $e^z = e^w$  by  $e^{-w}$  and use the addition law for the complex exponential to get  $e^{z-w} = 1$ , whereupon the Corollary to the “Minimal Period Theorem” of §II insures that  $z - w = 2\pi ni$  for some integer  $n$ . Thus  $\operatorname{Im} z - \operatorname{Im} w = 2\pi n$ , hence  $|\operatorname{Im} z - \operatorname{Im} w| = 2\pi|n|$ . But our hypothesis is that  $|\operatorname{Im} z - \operatorname{Im} w| < 2\pi$ , hence  $n = 0$ , whereupon  $z = w$ .  $\square$

**III. Zeros of the complex sine function.** Recall that the complex sine function is defined, for  $z \in \mathbb{C}$ , as:

$$\sin z \stackrel{\text{def}}{=} \frac{e^{iz} - e^{-iz}}{2i}.$$

The goal of this section is to show that this extension of the usual sine function of calculus to the complex plane does not add any new zeros.

THEOREM.  $\sin z = 0 \iff z = n\pi$  for some integer  $n$ !

PROOF. By trigonometry we know that  $\sin \pi n = 0$  for any integer  $n$ , so what's at stake here is the converse: if  $\sin z = 0$  then  $z = \pi n$  for some integer  $n$ .

Well,  $\sin z = 0$  implies that  $e^{iz} = e^{-iz}$ , so by multiplying both sides by  $e^{iz}$  and using the addition formula for the complex exponential, we see that  $e^{i2z} = 1$ , whereupon, by §I, there's an integer  $n$  such that  $2z = 2\pi n$ , i.e.,  $z = \pi n$ .  $\square$

**IV. Periodicity of the complex sine function.** *The minimal period of the complex sine function is  $2\pi$ !*

PROOF. We know that the complex sine function has period  $2\pi$  (because of the  $2\pi i$  periodicity of the complex exponential). The important assertion here is that if, for some complex number  $\alpha$ ,

$$(3) \quad \sin(z + \alpha) = \sin z \quad \text{for all } z \in \mathbb{C},$$

---

<sup>1</sup>Here “univalence” means “one-to-one-ness”. Also, I'll use “univalent” to mean “one-to-one”.

then  $\alpha$  is an integer multiple of  $2\pi$ .

So suppose we have (3) for some  $\alpha \in \mathbb{C}$ . Then upon setting  $z = 0$  we see that  $\sin \alpha = 0$ , hence by §III we know that  $\alpha$  is an integer multiple of  $\pi$ . We wish to show that this integer is *even*. In any case, we now know  $\alpha$  is *real*. Now set  $z = \pi/2$  in (3). Then  $\sin(\alpha + \frac{\pi}{2}) = \sin \frac{\pi}{2} = 1$ , so (since  $\alpha$  is real, hence we're now operating in the realm of ordinary trigonometry):

$$\alpha + \frac{\pi}{2} = \frac{\pi}{2} + 2\pi n \quad \text{for some integer } n$$

whereupon  $\alpha = 2\pi n$ , as promised. □

**V. Univalence of the complex sine function** (cf. page 50). *The complex sine is univalent on vertical the strip*

$$V \stackrel{\text{def}}{=} \{z \in \mathbb{C} : 0 < \operatorname{Re} z < \frac{\pi}{2}\}.$$

PROOF. Suppose  $z, w \in V$  and  $\sin z = \sin w$ .

*To show!*  $z = w$ .

Well, from the definition of the complex sine we know that

$$e^{iz} - e^{i(-z)} = e^{iw} - e^{i(-w)},$$

so upon rearranging to get “minus powers” on the same side of the equation:

$$(4) \quad e^{iz} - e^{iw} = e^{i(-z)} - e^{i(-w)} = e^{i(-z)}e^{i(-w)} (e^{iw} - e^{iz}).$$

So either

$$(5) \quad e^{iz} - e^{iw} = 0$$

or we can divide both sides of (4) by  $e^{iz} - e^{iw}$  to yield:

$$(6) \quad -1 = e^{i(-z)}e^{i(-w)} = e^{-i(z+w)}.$$

Suppose it's (5) that's true. Then  $e^{iz} = e^{iw}$  so by a now-familiar argument (using the addition formula for the complex exponential, and the result of §II),  $z = w + 2\pi n$  for some integer  $n$ , i.e.  $|\operatorname{Re} z - \operatorname{Re} w| = 2\pi|n|$ . But  $|\operatorname{Re} z - \operatorname{Re} w| < \pi$ , so  $n = 0$ , hence  $z = w$ .

*I claim !!! can't happen!* If it did, then we'd have  $e^{i\pi} = -1 = e^{-i(z+w)}$ , so by a now-familiar argument, we'd get

$$1 = e^{-i(z+w+\pi)},$$

whereupon the “Corollary” in §I would guarantee that  $z + w + \pi = 2\pi n$  for some integer  $n$ , i.e. that  $z + w$  would be an odd (possibly negative) multiple of  $\pi$ . Thus the same would be true of  $\operatorname{Re} z + \operatorname{Re} w$ : it would be an odd multiple of  $\pi$ . But since  $z, w \in V$  we know  $0 < \operatorname{Re} z + \operatorname{Re} w < \pi$ , so no such “odd multiple” can exist.

Thus it's only (5) that can happen, and so our theorem is proved. □