

*A novel high order method
for elliptic interface problems*

Guowei Wei


Department of Mathematics

Department of Electrical & Computer Engineering

Michigan State University

<http://www.math.msu.edu/~wei>

June, 15 2007



Major mathematical schemes for irregular interfaces and complex domains


- Immersed boundary method (**IBM**, Peskin, 1977) (1st order)
- Immersed interface method (**IIM**, LeVeque and Li, 1994) (2nd order)
- Ghost fluid method (**GFM**, Fedkiw and Osher, 2000) (1st order)

What do we do?


Solving the Poisson equation and the Helmholtz equation with interfaces

- **Complex irregular 3D interfaces without grid generation**
- **The highest order accuracy available (4th-16th orders)**
- **The highest order accuracy for sharp-edged interfaces**
- **Efficient for interface and high frequency waves**
- **Efficient for multiply connected topology**


3D missile geometry





Second order accuracy






Binding with Heme






CYTOCHROME C551





Second order accuracy



Fourth order accuracy



Matched interface & boundary (MIB)



Fourth order accuracy





Mesh	Error	Order
0.2	7.09E-2	-
0.1	3.67E-3	4.27
0.05	2.00E-4	4.20

$$u^- = 10(x + y + z) + 1$$



$$u^+ = 10 \cos(kx) \cos(ky) \cos(kz) + 20$$

$$k = 3, \quad \beta^- = z + 15, \quad \beta^+ = (x + y)/2 + 10$$



Fourth order accuracy



Fourth order accuracy






Matched interface & boundary (MIB)






Sixth order accuracy

Mesh size	Error	Order
0.2	5.3E-7	-
0.1	5.4E-9	6.6
0.05	5.3E-11	6.7



**6th order
accuracy**















**4th order
accuracy**



**2nd order
accuracy**





References of our matched interface &

boundary (MIB)

Maxwell's Equations (straight interfaces, 16th order, Zhao & Wei, JCP 2004)

2D Elliptic equations, (curved interfaces, 6th order, Zhou, Zhao, Feig & Wei, JCP 2006)

2D Sharp-edged interfaces, (2nd order, Yu, Zhou & Wei, JCP 2006)

3D Sharp-edged interfaces, (2nd order, Yu, Geng & Wei, JCP 2007)

3D arbitrary interfaces (4th & 6th orders, Yu & Wei, JCP 2007)

3D singular charges & sharp-edged interfaces (2nd order, Geng, Yu & Wei, 2007)

Our team

Z. Chen G.W. Wei S. Zhao D. Chen

W.H. Geng S.Y. Yu Y.H. Sun Y.C. Zhou

April, 2006