

## § 3.4 limits at Infinity.

Key points: ① horizontal/vertical asymptotes;  $\lim_{x \rightarrow \pm\infty} f(x) = L$  and  $\lim_{x \rightarrow a^\pm} f(x) = \pm\infty$ .

②  $\frac{1}{0^\pm} = \pm\infty$ ,  $\frac{1}{\pm\infty} = 0$ ,  $\infty^{\text{positive power}} = \infty$ ,  $\infty^{\text{negative power}} = 0$ .

③ highest term (leading term) rule for  $\lim_{x \rightarrow \pm\infty}$ .

• Def:  $\lim_{\substack{x \rightarrow \infty \\ (x \rightarrow -\infty)}} f(x) = L$  means as  $x$  approaches infinity (as  $x$  gets arbitrarily large) ( $+\infty$  or  $-\infty$ )  $f(x)$  approaches  $L$ . (positive or negative)

If  $L$  is finite,  $y=L$  is called a horizontal asymptote of  $y=f(x)$ .

Recall: If  $\lim_{x \rightarrow a^\pm} f(x) = \pm\infty$ ,  $x=a$  is called a vertical asymptote of  $y=f(x)$ . (Sec 1.5, lecture week 1, page 5).

•  $x \rightarrow \infty$  can be treated as "finite numbers" following the rules below:


①  $\lim_{x \rightarrow \pm\infty} \frac{1}{x} = 0 \iff \frac{1}{\pm\infty} = 0$ . In s.l.s, we take  $\lim_{x \rightarrow 0^\pm} \frac{1}{x} = \pm\infty \iff \frac{1}{0^\pm} = \pm\infty$ .

②  $x^{\text{positive power}}$  approaches  $\infty$  as  $x$  approaches  $\infty$ :  $\lim_{x \rightarrow \infty} \sqrt{x} = \infty$ ,  $\lim_{x \rightarrow \infty} x = \infty$ ,  $\lim_{x \rightarrow \infty} x^{\frac{3}{2}} = \infty$ ,  $\lim_{x \rightarrow \infty} x^2 = \infty$ .  
 $x^{\text{negative power}} = \frac{1}{x^{\text{positive power}}} \xrightarrow{x \rightarrow \infty} 0$ :  $\lim_{x \rightarrow \infty} x^{-\frac{1}{2}} = \lim_{x \rightarrow \infty} \frac{1}{\sqrt{x}} = 0$ ,  $\lim_{x \rightarrow \infty} x^{-5} = \lim_{x \rightarrow \infty} \frac{1}{x^5} = 0, \dots$

eg. 1.  $y = 3 + \frac{2}{x-1}$ .  $\lim_{x \rightarrow \pm\infty} 3 + \frac{2}{x-1} = 3 + \frac{2}{\pm\infty} = 3$

(sec 1.5  $\Rightarrow$ )  $\lim_{x \rightarrow 1^+} 3 + \frac{2}{x-1} = \infty$ ,  $\lim_{x \rightarrow 1^-} 3 + \frac{2}{x-1} = -\infty$ .

$y=3$  is a horizontal asymptote and  $x=1$  is a vertical asymptote.



Remark:  $\frac{\infty}{\infty}$  or  $\infty - \infty$  is indeterminate, we have to do some algebra changes first.

• Highest term (leading term) rule: In order to evaluate the limits for a ratio of power functions, we only need to keep the highest order terms in the numerator and the denominator and DROP ALL THE LOWER ORDER TERMS.

eg. 2.  $\lim_{x \rightarrow \infty} \frac{2-3x^2}{3+2x+5x^2} = \lim_{x \rightarrow \infty} \frac{-3x^2}{5x^2} = \lim_{x \rightarrow \infty} \frac{-3}{5} = -\frac{3}{5}$ .  $y = -\frac{3}{5}$  horizontal asymptote.

Remark:  $-3x^2$  is the highest term in the numerator;  $5x^2$  is the highest term in the denominator.

eg. 3. (More examples about highest term rule).

$$\lim_{x \rightarrow \infty} \frac{-7x + \sqrt{x}}{x^3 + 2x} = \lim_{x \rightarrow \infty} \frac{-7x}{x^3} = \lim_{x \rightarrow \infty} \frac{-7}{x^2} = \left( \frac{-7}{\infty} \right) = 0$$

$$\lim_{x \rightarrow \infty} \frac{2 + 3 \cdot x^{\frac{3}{2}}}{1 - \sqrt{x}} = \lim_{x \rightarrow \infty} \frac{3 \cdot x^{\frac{3}{2}}}{-x^{\frac{1}{2}}} = \lim_{x \rightarrow \infty} -3 \cdot x^1 = -\infty. \quad \text{Remark: } \frac{x^a}{x^b} = x^{a-b} = \frac{1}{x^{b-a}}$$

$$\lim_{x \rightarrow \infty} \frac{5x}{3-2x} = \lim_{x \rightarrow \infty} \frac{5x}{-2x} = -\frac{5}{2}. \quad y = -\frac{5}{2} \text{ is the horizontal asymptote.}$$

Remark: Highest order rule is only applied to  $x \rightarrow \infty$ .  $x = \frac{3}{2}$  is vertical asymptote

$$\lim_{x \rightarrow (\frac{3}{2})^+} \frac{5x}{3-2x} \quad \text{Direct plug in} \quad \frac{5 \cdot \frac{3}{2}}{3 - 2 \cdot \frac{3}{2}} = \frac{\text{finite number}}{0^-} = -\infty$$

Remark: Highest order rule has following product form.

(  $x > \frac{3}{2} \Rightarrow 3-2x < 0$  )  
negative sign comes from  $x \rightarrow (\frac{3}{2})^+$

$$\lim_{x \rightarrow \infty} \frac{(2-6x) \cdot (x^2+1)}{(3x+1) \cdot (2x^2-x)} = \lim_{x \rightarrow \infty} \frac{(-6x) \cdot x^2}{3x \cdot 2x^2} \cdot \text{Pick the highest term in each bracket.}$$

$$= \lim_{x \rightarrow \infty} \frac{-6x^3}{6x^3} = -1.$$

Remark: The formal argument for highest term rule: Pull out the highest order terms.

eg. 4 (Re-prove eg. 2).

$$\lim_{x \rightarrow \infty} \frac{2-3x^2}{3+2x+5x^2} = \lim_{x \rightarrow \infty} \frac{x^2 \cdot (\frac{2}{x^2} - 3)}{x^2 \cdot (\frac{3}{x^2} + \frac{2x}{x^2} + 5)} = \frac{0-3}{0+0+5} = -\frac{3}{5}.$$

Hints for WW.

\*7, \*8: Vertical asymptote. See more examples in §1.5, 1.6. Lec Notes. Week 1: Page 5-6.

\*5: Conjugation for root:  $\lim_{x \rightarrow \infty} \sqrt{x^2+3x} - x = \lim_{x \rightarrow \infty} \frac{(\sqrt{x^2+3x}-x)(\sqrt{x^2+3x}+x)}{\sqrt{x^2+3x}+x} = \lim_{x \rightarrow \infty} \frac{x^2+3x-x^2}{\sqrt{x^2+3x}+x}$

$$\lim_{x \rightarrow \infty} \sqrt{4x+1} - 4x = \lim_{x \rightarrow \infty} \frac{(\sqrt{4x+1}-4x)(\sqrt{4x+1}+4x)}{\sqrt{4x+1}+4x}$$

$$= \lim_{x \rightarrow \infty} \frac{4x+1-16x^2}{\sqrt{4x+1}+4x} = \lim_{x \rightarrow \infty} \frac{-16x^2}{4x}$$

$$= \lim_{x \rightarrow \infty} -4x = -\infty$$

$$= \lim_{x \rightarrow \infty} \frac{3x}{\sqrt{x^2+3x}+x} \quad \left[ \begin{array}{l} \text{highest order} \\ \text{rule for} \\ x^2+3x. \end{array} \right]$$

$$= \lim_{x \rightarrow \infty} \frac{3x}{x+x} = \frac{3}{2}.$$

\*6. (Squeeze theorem 3/6)

$$\frac{-1+x}{x} \leq \frac{\sin x + x}{x} \leq \frac{1+x}{x} \quad \text{since } -1 \leq \sin x \leq 1. \quad \lim_{x \rightarrow \infty} \frac{1+x}{x} = 1, \quad \lim_{x \rightarrow \infty} \frac{-1+x}{x} = 1 \Rightarrow \lim_{x \rightarrow \infty} \frac{\sin x + x}{x} = 1.$$

## §3.5. Curve Sketching

- key parts:
- ① Polynomial long division
  - ② Slant asymptote for rational functions.
  - ③ (Curve sketching) combination of 3.3, 3.4, 3.5.

eg0. • Divide 17 by 5, we have  $17 = 3 \cdot 5 + 2$

$$\begin{array}{r} 3 \leftarrow \text{quotient} \\ 5 \overline{) 17} \\ \underline{15} \\ 2 \leftarrow \text{remainder} \end{array}$$

• Divide  $x^2 + 2x - 4$  by  $x - 1$ ,  $x^2 + 2x - 4 = q(x) \cdot (x - 1) + r(x)$

• Find the quotient  $q(x)$  and remainder  $r(x)$  by polynomial long division.

$$\begin{array}{r} x+3 \leftarrow q(x) \\ x-1 \overline{) x^2+2x-4} \\ \underline{x^2-x} \\ 3x-4 \\ \underline{3x-3} \\ -1 \leftarrow r(x) \end{array}$$

i.e.

$$x^2 + 2x - 4 = (x + 3) \cdot (x - 1) - 1.$$

• Consider the ratio  $\frac{17}{5} = \frac{3 \cdot 5 + 2}{5} = 3 + \frac{2}{5}$

• Consider the ratio of polynomials:  $\frac{x^2 + 2x - 4}{x - 1} = \frac{(x + 3) \cdot (x - 1) - 1}{x - 1} = x + 3 - \frac{1}{x - 1}$ .  
(Rational functions)

• Slant asymptote: If  $f(x)$  approaches a line  $y = m \cdot x + b$  as  $x$  approaches infinity, then  $y = mx + b$  is the SLANT ASYMPTOTE of  $f(x)$ .

eg1:  $f(x) = \frac{x^2 + 2x - 4}{x - 1} = \boxed{x + 3} - \frac{1}{x - 1}$ .  $f(x)$  approaches  $y = x + 3$  as  $x \rightarrow \infty$

since  $\lim_{x \rightarrow \infty} (f(x) - (x + 3)) = -\frac{1}{x - 1} \rightarrow 0$  as  $x \rightarrow \infty$ .

i.e.  $y = x + 3$  is the slant asymptote of  $f(x)$ .

• Conclusion: If a rational function can be written as  $f(x) = m \cdot x + b + \frac{r(x)}{d(x)}$  via polynomial long division, then  $y = mx + b$  is the slant asymptote of  $y = f(x)$ .

eg.2. let  $f(x) = \frac{4x^2}{2x-5}$ . Find all the asymptotes (vertical/horizontal/slant) of  $f(x)$ .

• Vertical:  $x = \frac{5}{2}$  since  $\lim_{x \rightarrow (\frac{5}{2})^+} \frac{4x^2}{2x-5} = \infty$  (or  $\lim_{x \rightarrow (\frac{5}{2})^-} \frac{4x^2}{2x-5} = -\infty$ )

• Horizontal: None.  $\lim_{x \rightarrow \pm\infty} \frac{4x^2}{2x-5} \stackrel{\text{highest term}}{\sim} \lim_{x \rightarrow \pm\infty} \frac{4x^2}{2x} = \lim_{x \rightarrow \pm\infty} 2x = \pm\infty$  (Not finite)


• Slant:  $y = 2x + 5$ . Poly-long Division: 
$$\begin{array}{r} 2x + 5 \\ 2x - 5 \overline{) 4x^2 + 0x + 0} \\ \underline{4x^2 - 10x} \phantom{+ 0} \\ 10x + 0 \\ \underline{10x - 25} \\ 25 \end{array}$$
 
$$4x^2 = \underbrace{(2x+5)(2x-5)}_{\text{quotient}} + \underbrace{25}_{\text{remainder}}$$
  
 since  $\frac{4x^2}{2x-5} = 2x + 5 + \frac{25}{2x-5}$ . 
$$\frac{4x^2}{2x-5} = \underbrace{2x+5}_{\text{slant asymp.}} + \frac{25}{2x-5}$$

eg.3. (Curve sketching: combination of 3.3-3.5, fib)


sketch the graph of  $y=f(x)$  such that

- $f$  is continuous and has SLANT asymptote  $y=x$ .
- $f' > 0$  for  $x \in (-\infty, -1) \cup (3, +\infty)$ ,  $f' < 0$  for  $x \in (-1, 3)$
- $f'' > 0$  for  $x \in (-\infty, 3) \cup (1, \infty)$ ,  $f'' < 0$  for  $x \in (-3, 1)$

(The answer is not unique)  
 All the following three are qualified answers.



- Local maximum of  $f$  occurs at  $x = -1$ .
- Local minimum of  $f$  occurs at  $x = 3$ .
- Inflection points of  $f$  are  $x = 3$ ,  $x = 1$ .



eg 4. Suppose  $f(x) = \frac{x}{x^2+1}$ ,  $f'(x) = \frac{1-x^2}{(x^2+1)^2}$ ,  $f''(x) = \frac{2(x^3-3x)}{(x^2+1)^3}$

(a).  $f$  is an odd function whose graph is symmetric with respect to the origin.

Reason:  $f(-x) = \frac{-x}{(-x)^2+1} = -\left[\frac{x}{x^2+1}\right] = -f(x)$ . Remark:  $f$  is even if  $f(-x) = f(x)$


$f$  is odd if  $f(-x) = -f(x)$ .

(b). Interval of increasing/decreasing and local extrema.

$f'(x) = \frac{1-x^2}{(x^2+1)^2} = \frac{(1-x)(1+x)}{(x^2+1)^2} = 0 \Rightarrow x = -1, 1$  (two critical points)

(defined for all  $x$ )

$-1, 1$  divide  $(-\infty, +\infty)$  into



Increasing:  $[-1, 1]$  where  $f' > 0$

Decreasing:  $(-\infty, -1) \cup (1, +\infty)$  where  $f' < 0$

local maximum occurs at  $x = 1$ , local minimum occurs at  $x = -1$ .


(c). Concavity:  $f''(x) = \frac{2x \cdot (x^2-3)}{(x^2+1)^3} = \frac{2x \cdot (x+\sqrt{3}) \cdot (x-\sqrt{3})}{(x^2+1)^3} = 0$

$\Rightarrow x = 0, x = -\sqrt{3}, x = \sqrt{3}$

Concave up:  $(-\sqrt{3}, 0) \cup (\sqrt{3}, +\infty)$  ( $f'' > 0$ )


Concave down:  $(-\infty, -\sqrt{3}) \cup (0, \sqrt{3})$  ( $f'' < 0$ )

Inflection points:  $x = -\sqrt{3}, 0, \sqrt{3}$ .



Hints for WW 5.  $f(x) = \frac{x^3}{x^2-9}$ ,  $f'(x) = \frac{x^4-27x^2}{(x^2-9)^2}$ ,  $f''(x) = \frac{18x \cdot (x^2+27)}{(x^2-9)^3}$


(You can seek help from Wolfram/Alpha for the expression of  $f'$ ).



Increasing:  $(-\infty, -\sqrt{27}] \cup (\sqrt{27}, +\infty)$

Decreasing:  $(-\sqrt{27}, -3) \cup (-3, 3) \cup (3, \sqrt{27}]$

Rank:  $-3, 3$  are not in the domain.



Concave up:  $(-3, 0) \cup (3, +\infty)$

Concave down:  $(-\infty, -3) \cup (0, 3)$

Inflection:  $x = 0$ . ( $x = \pm 3$  not in the domain)

★ ★ ★ eg. 5. Analyze  $f(x) = 2x - 3x^{\frac{2}{3}}$  (related to wu 2).

① Domain of  $f: (-\infty, \infty)$ .  $\Rightarrow$   $f$  has no vertical asymptote

②  $f$  has no horizontal asymptote since  $\lim_{x \rightarrow \pm\infty} 2x - 3x^{\frac{2}{3}} = \lim_{x \rightarrow \pm\infty} x \left[ 2 - \frac{3}{x^{\frac{1}{3}}} \right]$

③  $f'(x) = (2x - 3x^{\frac{2}{3}})' = \boxed{2 - 2x^{-\frac{1}{3}}}$   $= \pm\infty \cdot (2 - 0) = \pm\infty$

④ Critical points of  $f$ : (Hint: critical pts  $\Leftrightarrow f'$  D.N.E or  $f'=0$ )

$f'(x) = 2 - \frac{2}{x^{\frac{1}{3}}}$  D.N.E  $\Leftrightarrow$  Denominator is zero  $\Leftrightarrow x^{\frac{1}{3}} = 0 \Rightarrow x = 0$

$f'(x) = 2 - \frac{2}{x^{\frac{1}{3}}} = 0 \Rightarrow 2 = \frac{2}{x^{\frac{1}{3}}} \Rightarrow x^{\frac{1}{3}} = 1 \Rightarrow x = 1$

Critical points are  $x=0$  and  $x=1$ .

⑤ Increasing/Decreasing Intervals: (determined by the signs of  $f'$ ).

critical points 0, 1 divide  $(-\infty, \infty)$  into  $\frac{(-\infty, 0) \quad 0 \quad (0, 1) \quad 1 \quad (1, +\infty)}$


$x < 0$ ,  $f'(x) > 0$  ( $f'(1) = 4 > 0$ )  $f' \quad +++ \quad --- \quad +++$

$0 < x < 1$ ,  $0 < x^{\frac{1}{3}} < 1 \Rightarrow 2 - \frac{2}{x^{\frac{1}{3}}} < 0$ ,  $f' < 0$

$x > 1$ ,  $x^{\frac{1}{3}} > 1 \Rightarrow 2 - \frac{2}{x^{\frac{1}{3}}} > 0$ ,  $f' > 0$ .

Increasing Interval(s):  $(-\infty, 0) \cup (1, +\infty)$ . Decreasing Interval(s):  $[0, 1]$

⑥ local maximum: attained at  $x=0$  (increasing  $\rightarrow$  decreasing )

local minimum: attained at  $x=1$  (decreasing  $\rightarrow$  increasing )

⑦ Concavity and inflection points:  $f''(x) = (2 - 2x^{-\frac{1}{3}})' = 0 - 2 \cdot (-\frac{1}{3})x^{-\frac{2}{3}} = \frac{2}{3} \cdot x^{-\frac{2}{3}}$

Note that  $f'' = \frac{2}{3} \cdot \frac{1}{(x^{\frac{1}{3}})^2}$  is forever positive except 0 since  $\square^2 > 0$

$f$  is concave up on  $(-\infty, 0) \cup (0, +\infty)$

and concave down nowhere

and no inflection points.