

§ 4.5. Substitution Method

Key points: ① Differential Notation: If $u = g(x)$, then $du = g'(x) \cdot dx$.

$$\textcircled{2} \quad u\text{-sub: } \int \underbrace{f(g(x)) \cdot g'(x) \cdot dx}_{du} \xrightarrow{\substack{u=g(x) \\ du=g'(x)dx}} \int f(u) \cdot du$$

• Goal of U-Substitution Method: By changing the variable x into u , we convert the integral into one of those five basic integrals which we can deal with.

• Basic integrals: $\int x^n dx = \frac{1}{n+1} x^{n+1} + C$, $\int \sin x dx = -\cos x + C$, $\int \cos x dx = \sin x + C$

$$\int \sec^2 x dx = \tan x + C, \quad \int \sec x \cdot \tan x dx = \sec x + C.$$

① Differential Notation and Composition of functions.

e.g. 1. $u = x^2 + 1, \quad \frac{du}{dx} = (x^2 + 1)' = 2x \Rightarrow \boxed{du = 2x \cdot dx}$

e.g. 2. If $f(x) = \sqrt{x}$, ~~then~~ $u = x^2 + 1$, then $f(u) = \sqrt{u} = \sqrt{x^2 + 1}$

② U-Sub method for indefinite integral

e.g. 3. Evaluate $\int \sqrt{u} \cdot du \stackrel{n=\frac{1}{2}}{=} \frac{1}{\frac{1}{2}+1} \cdot u^{\frac{1}{2}+1} + C = \boxed{\frac{2}{3} \cdot u^{\frac{3}{2}} + C}$

e.g. 4. $\int \sqrt{x^2 + 1} \cdot 2x \cdot dx$.

Set $u = x^2 + 1$, According to egl. $du = 2x \cdot dx$.

$$= \int \sqrt{u} \cdot du.$$

Plug in u and du . By substituting u and du the integral of x turns into an integral of u with simpler form, which we evaluate in egl. 3.

$$= \frac{2}{3} \cdot u^{\frac{3}{2}} + C$$

$$= \frac{2}{3} (x^2 + 1)^{\frac{3}{2}} + C$$

Replace u by $x^2 + 1$ in the last step.

- The key of u-sub is to find the right substitution u .

More examples:

eg.5. Evaluate $\int x \cdot (2x^2 + 1)^3 dx$. Hint: $u = 2x^2 + 1$ is the ugly part.
(14 final)

$$u = 2x^2 + 1, du = 4x \cdot dx \Rightarrow \boxed{x \cdot dx = \frac{1}{4} \cdot du}$$

$$= \int (2x^2 + 1)^3 \cdot x \cdot dx$$

$$= \int u^3 \cdot \frac{1}{4} \cdot du = \frac{1}{4} \cdot \frac{1}{3+1} \cdot u^{3+1} + C$$

$$= \frac{1}{16} \cdot u^4 + C = \boxed{\frac{1}{16}(2x^2 + 1)^4 + C}$$

eg.6. Evaluate $\int \frac{\sec(\frac{x}{2}) \cdot \tan(\frac{x}{2})}{\sqrt{\sec(\frac{x}{2})}} dx$. Hint: $u = \sec(\frac{x}{2})$ is the ugly part.
(14 final)

$$u = \sec(\frac{x}{2}), du = \sec(\frac{x}{2}) \tan(\frac{x}{2}) \cdot \frac{1}{2} \cdot dx, \text{ chain rule for } (\sec(\frac{x}{2}))'$$

$$\Rightarrow 2du = \sec(\frac{x}{2}) \cdot \tan(\frac{x}{2}) \cdot dx \quad = \sec(\frac{x}{2}) \cdot \tan(\frac{x}{2})$$

$$= \int \frac{2du}{\sqrt{u}} = \int 2 \cdot u^{-\frac{1}{2}} du$$

$$= 2 \cdot \frac{1}{-\frac{1}{2}+1} \cdot u^{-\frac{1}{2}+1} + C$$

$$= 2 \cdot 2 \cdot u^{\frac{1}{2}} + C \quad \underline{\text{back to } x} \quad \boxed{4(\sec(\frac{x}{2}))^{\frac{1}{2}} + C}$$

- Linear substitution and more general form of the five basic integrals.

$$\star \int (ax+b)^n dx = \frac{1}{n+1} \cdot x^{n+1} + C, \int \sin(ax+b) dx = -\frac{1}{a} \cos(ax+b) + C, \int \cos(ax+b) dx = \frac{1}{a} \sin(ax+b) + C$$

$$\int \tan(ax+b) \cdot \sec(ax+b) dx = \frac{1}{a} \sec(ax+b) + C, \int \sec^2(ax+b) dx = \frac{1}{a} \tan(ax+b) + C.$$

eg.7. $\int \sin(1-2x) dx$

$\frac{u=1-2x}{du=-2dx}$ $\int \sin u \cdot \frac{du}{-2} = \frac{1}{2} (-\cos u) + C = \boxed{-\frac{1}{2} \cos(1-2x) + C}$

$\frac{du}{-2} = dx$

③ U-sub for definite integral.

eg.8. Evaluate the definite integral $\int_0^2 \frac{1}{\sqrt{9-4x}} dx$.

Ugly part: $9-4x$. $u=9-4x$, $du=-4dx \Rightarrow dx = \frac{du}{-4}$

Caution: \int_a^b a, b are for x . They also change as we substitute $9-4x$ by u .

$$\begin{array}{c} \int_{x=0}^{x=2} \\ \xrightarrow{u=9-4x} \end{array} \begin{array}{l} u=9-4(2)=1 \\ u=9-4(0)=9 \end{array}, \quad \begin{array}{l} u=1 \\ u=9 \end{array}$$

$$\int_0^2 \frac{1}{\sqrt{9-4x}} dx \xrightarrow{u=9-4x} \int_9^1 \frac{1}{\sqrt{u}} \cdot \frac{du}{-4} \quad \text{Use the flipping trick}$$

$$= \int_1^9 \frac{1}{4} \cdot u^{-\frac{1}{2}} du = \frac{1}{4} \cdot \frac{1}{\frac{1}{2}} u^{\frac{1}{2}} \Big|_1^9$$

$$= \frac{1}{2} \cdot \sqrt{u} \Big|_1^9 = \boxed{\frac{1}{2} \cdot \sqrt{9} - \frac{1}{2} \cdot \sqrt{1}} = \boxed{1}$$

eg.9. Evaluate $\int_0^{\frac{\pi}{2}} 3 \cdot \tan(\frac{x}{2}) \cdot \sec^2(\frac{x}{2}) dx$. Hint: $(\tan(\frac{x}{2}))' = \sec^2(\frac{x}{2})$


$$u = \tan(\frac{x}{2}), \quad du = \frac{1}{2} \sec^2(\frac{x}{2}) dx \quad \xrightarrow{x=\frac{\pi}{2}} \quad \begin{array}{l} u = \tan(\frac{\pi}{2}) = 1 \\ u = \tan(0) = 0 \end{array}$$

$$2du = \sec^2(\frac{x}{2}) dx, \quad \xrightarrow{x=0} \quad \begin{array}{l} u = \tan(\frac{\pi}{2}) = 1 \\ u = \tan(0) = 0 \end{array}$$

$$= \int_0^1 3 \cdot u \cdot 2du = 6 \cdot \frac{1}{2} u^2 \Big|_0^1 = \boxed{3 \cdot 1^2 - 3 \cdot 0^2 = 3}$$

④ Symmetry integral.

If $f(x)$ is odd, i.e., $f(-x) = -f(x)$, then $\int_a^a f(x) dx = 0$



f odd means the graph of f is symmetric about the origin. Then the area above and below x -axis are the same, which will be cancelled out.

eg.10. $f(x) = \sin x \cdot (x^2 + 1)$. $\int_{-8}^8 \sin x \cdot (x^2 + 1) dx = 0$

since $f(-x) = \sin(-x) \cdot ((-x)^2 + 1) = -\sin x \cdot (x^2 + 1) = -f(x)$. ($\sin x$ is odd),

More examples and hints for webwork:

eg 11. (mw8). $\int_8^{11} x \cdot \sqrt{x-7} dx$. Hint: $u=x-7$, $du=dx$.

$$= \int x \cdot \sqrt{u} du$$

$$= \int (u+7) \cdot \sqrt{u} du$$

$$= \int u \cdot u^{\frac{1}{2}} + 7 \cdot u^{\frac{1}{2}} du$$

$$= \int_1^4 u^{\frac{3}{2}} + 7 \cdot u^{\frac{1}{2}} du = \frac{2}{5} u^{\frac{5}{2}} + 7 \cdot \frac{2}{3} u^{\frac{3}{2}} \Big|_1^4$$

$$4^{\frac{5}{2}} = (\sqrt{4})^5 = 32.$$

$$4^{\frac{3}{2}} = (\sqrt{4})^3 = 8$$

$$\begin{aligned} &= \frac{2}{5} \cdot 4^{\frac{5}{2}} + \frac{14}{3} \cdot 4^{\frac{3}{2}} - \left(\frac{2}{5} \cdot 1 + \frac{14}{3} \cdot 1 \right) \\ &= \frac{64}{5} + \frac{112}{3} - \frac{2}{5} - \frac{14}{3} = \boxed{\frac{62}{5} + \frac{98}{5}} \end{aligned}$$

eg 12. (mw6) $\int \frac{1}{x^2} \sin\left(\frac{3}{x}\right) \cdot \cos\left(\frac{3}{x}\right) dx$. Hint:

$$= \int \cos\left(\frac{3}{x}\right) \cdot \frac{1}{x^2} \sin\left(\frac{3}{x}\right) dx$$

$$= \int u \cdot \frac{1}{3} du$$

$$= \frac{1}{3} \cdot \frac{1}{2} u^2 + C$$

$$= \boxed{\frac{1}{6} \cdot (\cos(\frac{3}{x}))^2 + C}$$

$$u = \cos\left(\frac{3}{x}\right).$$

$$du = -\sin\left(\frac{3}{x}\right) \cdot \left(\frac{3}{x}\right)' dx. \quad \text{chain rule.}$$

$$= -\sin\left(\frac{3}{x}\right) \cdot \frac{-3}{x^2} dx \quad \text{inner } \frac{3}{x} = 3x^{-1} \\ (3x^{-1})' = 3 \cdot \frac{-1}{x^2}$$

$$\Rightarrow \frac{1}{3} du = \sin\left(\frac{3}{x}\right) \cdot \frac{1}{x^2} dx$$

eg 13. $\int \frac{x^3}{\sqrt{1+2x^4}} dx$. Ugly part: $u=1+2x^4$, $du=8x^3 dx$

$$\Rightarrow \frac{1}{8} du = x^3 dx$$

$$= \int \frac{1}{\sqrt{u}} \cdot x^3 dx$$

$$= \int \frac{1}{\sqrt{u}} \cdot \frac{1}{8} du = \int u^{-\frac{1}{2}} \cdot \frac{1}{8} du = \frac{1}{-\frac{1}{2}+1} u^{-\frac{1}{2}+1} \cdot \frac{1}{8} + C$$

$$= 2(u)^{\frac{1}{2}} \cdot \frac{1}{8} + C$$

$$= \frac{1}{4} (1+2x^4)^{\frac{1}{2}} + C.$$